

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
All origins <sup>3</sup>												
Both sexes												
2002 .....	847.3	695.0	31.2	17.4	81.4	103.6	202.9	430.1	952.4	2,314.7	5,556.9	14,828.3
2001 .....	848.5	683.4	33.3	17.3	80.7	105.2	203.6	428.9	964.6	2,353.3	5,582.4	15,112.8
2000 .....	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4
1999 .....	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6
Male												
2002 .....	846.6	761.5	35.2	20.0	117.3	142.2	257.5	547.5	1,184.0	2,855.3	6,760.5	16,254.5
2001 .....	846.4	749.8	37.0	19.8	117.0	143.7	259.6	545.1	1,192.7	2,911.5	6,833.0	16,744.8
2000 .....	853.0	806.5	35.9	20.9	114.9	138.6	255.2	542.8	1,230.7	2,979.6	6,972.6	17,501.4
1999 .....	859.2	805.0	37.9	21.5	113.1	139.7	254.9	533.1	1,252.0	3,073.7	7,083.3	17,597.2
Female												
2002 .....	848.0	625.3	27.0	14.7	43.7	64.0	148.8	316.9	738.0	1,864.7	4,757.9	14,209.6
2001 .....	850.4	613.9	29.5	14.6	42.6	66.0	148.2	316.8	754.0	1,890.8	4,760.5	14,429.9
2000 .....	855.0	663.4	28.7	15.0	43.1	63.5	143.2	312.5	772.2	1,921.2	4,814.7	14,719.2
1999 .....	854.9	663.6	30.3	15.6	43.7	64.1	141.8	307.6	777.6	1,952.3	4,825.4	14,731.3
Hispanic												
Both sexes												
2002 .....	302.2	592.7	29.8	15.5	77.2	79.3	145.8	321.3	719.8	1,734.1	4,143.1	10,679.0
2001 .....	306.8	572.7	30.6	14.7	75.6	84.5	154.8	334.4	722.2	1,776.0	4,262.6	11,821.8
2000 .....	303.8	596.3	29.6	15.7	72.8	84.6	157.6	330.2	744.4	1,803.8	4,335.9	11,822.9
1999 .....	305.7	583.8	30.9	15.6	71.3	85.5	158.2	333.2	752.5	1,853.0	4,356.3	12,177.9
Male												
2002 .....	328.7	644.0	34.2	17.4	114.4	112.5	192.5	423.4	937.4	2,193.4	5,043.5	11,674.1
2001 .....	332.9	624.4	33.8	16.6	111.5	118.0	208.5	443.9	923.9	2,242.6	5,258.0	12,888.3
2000 .....	331.3	637.1	31.5	17.9	107.7	120.2	211.0	439.0	965.7	2,287.9	5,395.3	13,086.2
1999 .....	332.6	623.4	32.9	17.8	104.1	120.6	215.1	444.4	974.8	2,368.9	5,379.2	13,485.9
Female												
2002 .....	274.0	539.1	25.3	13.5	34.1	40.0	94.9	219.8	524.3	1,368.7	3,526.4	10,186.0
2001 .....	279.0	518.9	27.2	12.7	33.7	45.2	97.0	226.7	543.0	1,408.0	3,589.8	11,300.5
2000 .....	274.6	553.6	27.5	13.4	31.7	43.4	100.5	223.8	548.4	1,423.2	3,624.5	11,202.8
1999 .....	277.2	542.3	28.7	13.3	32.9	44.9	97.4	224.9	555.8	1,448.8	3,675.7	11,547.3
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	928.8	713.0	31.4	17.8	81.9	108.4	210.3	438.8	967.2	2,347.0	5,610.8	14,958.7
2001 .....	926.2	703.5	33.8	17.7	81.4	108.9	209.2	435.6	979.5	2,383.9	5,628.5	15,199.0
2000 .....	929.6	760.4	32.9	18.4	80.9	104.1	202.9	431.8	1,006.5	2,427.9	5,707.6	15,619.8
1999 .....	929.9	760.7	34.8	19.1	80.5	104.7	201.6	423.3	1,019.2	2,485.0	5,753.4	15,631.0
Male												
2002 .....	928.0	782.6	35.4	20.4	117.3	148.6	266.0	557.0	1,198.1	2,889.1	6,825.9	16,418.0
2001 .....	923.6	773.2	37.6	20.4	117.5	148.5	265.2	551.6	1,207.0	2,944.1	6,888.0	16,866.3
2000 .....	928.1	836.8	36.8	21.5	115.9	141.6	259.2	548.7	1,244.2	3,009.8	7,020.7	17,639.8
1999 .....	932.2	834.9	38.8	22.1	114.6	142.7	258.1	537.3	1,265.6	3,102.8	7,133.0	17,710.4
Female												
2002 .....	929.5	640.1	27.2	15.0	45.3	68.5	155.7	325.2	753.0	1,894.4	4,805.4	14,329.1
2001 .....	928.6	630.4	29.9	14.9	44.1	69.6	154.3	324.0	768.7	1,918.4	4,801.8	14,505.6
2000 .....	931.0	680.1	28.8	15.2	45.0	66.8	147.7	319.2	786.8	1,947.5	4,852.4	14,802.9
1999 .....	927.8	682.9	30.6	16.0	45.4	67.1	146.3	313.7	791.7	1,977.5	4,858.2	14,798.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
Non-Hispanic white												
Both sexes												
2002 .....	997.5	575.9	27.1	16.0	75.8	95.9	191.6	400.1	912.8	2,296.1	5,596.0	15,149.1
2001 .....	991.1	555.4	30.1	16.3	74.1	96.4	188.3	396.0	924.0	2,325.5	5,604.4	15,371.2
2000 .....	993.2	596.5	28.5	17.1	73.6	90.1	181.0	390.5	950.2	2,366.2	5,675.4	15,799.7
1999 .....	990.7	593.1	30.1	17.3	72.4	91.3	177.8	381.5	957.8	2,420.0	5,716.0	15,795.3
Male												
2002 .....	983.9	643.5	30.3	18.3	106.7	130.9	244.9	509.9	1,126.5	2,824.1	6,801.7	16,641.9
2001 .....	975.6	611.6	33.7	18.6	105.1	131.5	241.6	502.6	1,136.3	2,869.4	6,851.5	17,055.3
2000 .....	978.5	658.7	32.4	20.0	103.5	123.0	233.9	497.7	1,170.9	2,930.5	6,977.8	17,853.2
1999 .....	979.6	658.1	33.4	19.9	100.8	124.5	230.0	483.7	1,187.4	3,023.2	7,088.0	17,871.2
Female												
2002 .....	1,010.6	504.8	23.8	13.6	43.8	60.3	138.3	292.1	710.5	1,846.0	4,787.9	14,504.3
2001 .....	1,006.1	496.4	26.3	13.9	41.9	60.9	134.9	291.0	723.5	1,864.1	4,777.3	14,670.6
2000 .....	1,007.3	530.9	24.4	13.9	42.6	56.8	128.1	285.0	742.1	1,891.0	4,819.3	14,971.7
1999 .....	1,001.3	524.6	26.7	14.7	42.9	57.8	125.6	281.0	741.6	1,915.1	4,817.7	14,967.5
Non-Hispanic black												
Both sexes												
2002 .....	792.8	1,280.3	48.4	25.3	116.9	193.9	357.8	782.0	1,592.9	3,204.8	6,584.8	14,386.7
2001 .....	798.1	1,354.4	49.5	24.0	121.4	191.5	365.8	786.9	1,602.5	3,300.7	6,645.3	14,660.8
2000 .....	805.5	1,477.8	51.7	25.0	123.2	194.6	365.1	802.2	1,633.9	3,378.8	6,764.3	14,837.5
1999 .....	812.1	1,510.2	54.6	27.5	125.5	191.1	376.9	796.7	1,681.7	3,454.5	6,805.4	14,839.2
Male												
2002 .....	842.3	1,369.3	55.8	29.8	177.4	274.3	445.7	1,002.9	2,074.5	4,078.9	8,246.2	15,776.9
2001 .....	849.7	1,484.2	55.1	28.6	186.2	267.9	456.3	1,014.6	2,055.4	4,218.9	8,426.8	16,576.2
2000 .....	859.5	1,623.2	56.1	29.1	187.2	270.2	462.7	1,036.2	2,111.4	4,303.9	8,551.0	16,944.8
1999 .....	872.8	1,650.8	61.5	33.1	191.4	267.9	479.5	1,048.1	2,176.7	4,397.8	8,615.3	17,385.3
Female												
2002 .....	748.0	1,187.5	40.8	20.7	56.1	121.3	280.1	592.7	1,206.3	2,584.8	5,644.9	13,855.4
2001 .....	751.2	1,217.8	43.6	19.3	56.8	122.4	285.7	591.5	1,238.2	2,653.0	5,645.1	13,951.4
2000 .....	756.7	1,327.7	47.1	20.7	60.1	126.5	278.8	601.2	1,249.4	2,728.0	5,758.6	14,054.9
1999 .....	757.3	1,365.4	47.4	21.8	60.5	122.1	286.4	581.7	1,285.1	2,789.2	5,792.0	13,887.7
Salmonella infections (A01-A02)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
All origins <sup>3</sup>												
Female												
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
Non-Hispanic white												
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
Hispanic												
Male												
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Shigellosis and amebiasis (A03,A06)</b>												
<b>Non-Hispanic black</b>												
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Certain other intestinal infections (A04,A07-A09)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	0.9	*	*	*	*	*	*	0.1	0.5	2.0	7.0	21.0
2001 .....	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000 .....	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.7	11.8
1999 .....	0.4	*	*	*	*	*	*	0.1	0.2	1.0	3.2	10.3
<b>Male</b>												
2002 .....	0.7	*	*	*	*	*	*	0.1	0.6	2.2	7.0	20.4
2001 .....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.6	12.1
2000 .....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.4
1999 .....	0.3	*	*	*	*	*	*	*	0.2	0.9	3.5	9.5
<b>Female</b>												
2002 .....	1.0	*	*	*	*	*	*	0.1	0.5	1.8	6.9	21.3
2001 .....	0.7	*	*	*	*	*	*	0.1	0.3	1.3	4.5	13.1
2000 .....	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	11.9
1999 .....	0.5	*	*	*	*	*	*	*	0.2	1.0	3.0	10.7
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	5.2	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	1.0	*	*	*	*	*	*	0.1	0.5	2.1	7.1	21.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2001 .....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000 .....	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.8	11.9
1999 .....	0.4	*	*	*	*	*	*	*	0.2	1.0	3.3	10.4
Male												
2002 .....	0.8	*	*	*	*	*	*	0.1	0.6	2.3	7.2	21.0
2001 .....	0.5	*	*	*	*	*	*	*	0.3	1.3	4.7	12.0
2000 .....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.5
1999 .....	0.3	*	*	*	*	*	*	*	0.2	0.9	3.6	9.6
Female												
2002 .....	1.1	*	*	*	*	*	*	0.1	0.5	1.8	7.0	21.6
2001 .....	0.7	*	*	*	*	*	*	0.1	0.4	1.3	4.5	13.2
2000 .....	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	12.1
1999 .....	0.5	*	*	*	*	*	*	*	0.2	1.0	3.1	10.8
Non-Hispanic white												
Both sexes												
2002 .....	1.1	*	*	*	*	*	*	0.2	0.5	2.2	7.5	22.5
2001 .....	0.7	*	*	*	*	*	*	0.1	0.3	1.4	4.7	13.6
2000 .....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.0	12.3
1999 .....	0.5	*	*	*	*	*	*	*	0.2	1.0	3.4	11.0
Male												
2002 .....	0.9	*	*	*	*	*	*	0.2	0.5	2.3	7.5	22.1
2001 .....	0.5	*	*	*	*	*	*	*	0.3	1.4	4.7	12.6
2000 .....	0.5	*	*	*	*	*	*	*	0.3	1.3	4.3	12.0
1999 .....	0.4	*	*	*	*	*	*	*	0.2	1.0	3.8	10.0
Female												
2002 .....	1.3	*	*	*	*	*	*	0.1	0.5	2.0	7.5	22.7
2001 .....	0.9	*	*	*	*	*	*	0.2	0.3	1.3	4.7	14.0
2000 .....	0.7	*	*	*	*	*	*	*	0.3	1.2	3.9	12.5
1999 .....	0.6	*	*	*	*	*	*	*	*	1.0	3.2	11.4
Non-Hispanic black												
Both sexes												
2002 .....	0.4	*	*	*	*	*	*	*	1.1	1.9	3.7	13.3
2001 .....	0.3	*	*	*	*	*	*	*	*	1.3	3.0	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	8.3
1999 .....	0.2	*	*	*	*	*	*	*	*	*	2.8	*
Male												
2002 .....	0.4	*	*	*	*	*	*	*	*	2.9	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.5	*	*	*	*	*	*	*	*	*	3.6	13.4
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Tuberculosis (A16-A19)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	2.6
2001 .....	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.8	1.6	2.8
2000 .....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.0	2.7
1999 .....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	4.0
2001 .....	0.3	*	*	*	*	*	0.1	0.3	0.7	1.2	2.0	4.4
2000 .....	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.8	4.6
1999 .....	0.4	*	*	*	*	*	0.2	0.5	0.8	1.4	2.8	7.1
<b>Female</b>												
2002 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.1
2001 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
2000 .....	0.2	*	*	*	*	*	*	*	0.2	0.7	1.5	2.0
1999 .....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.6	3.0
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	1.8	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	3.6	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	4.5	*
1999 .....	0.4	*	*	*	*	*	*	*	1.2	2.3	5.6	*
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.8	2.5
2001 .....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.7
2000 .....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.8	1.9	2.5
1999 .....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.9	1.9	3.9
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	0.1	0.3	0.6	1.2	2.3	3.8
2001 .....	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	1.9	4.3
2000 .....	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.6	4.3
1999 .....	0.4	*	*	*	*	*	0.2	0.4	0.8	1.3	2.6	6.8
<b>Female</b>												
2002 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	1.9
2001 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.1
2000 .....	0.2	*	*	*	*	*	*	*	*	0.7	1.4	1.8
1999 .....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	2.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Tuberculosis (A16-A19)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.9
2001 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.0
2000 .....	0.2	*	*	*	*	*	*	0.1	0.1	0.5	1.2	2.0
1999 .....	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.3	3.0
<b>Male</b>												
2002 .....	0.2	*	*	*	*	*	*	0.1	0.3	0.8	1.7	2.9
2001 .....	0.2	*	*	*	*	*	*	0.1	0.4	0.6	1.4	2.9
2000 .....	0.2	*	*	*	*	*	*	0.2	*	0.6	1.5	3.6
1999 .....	0.3	*	*	*	*	*	*	0.2	0.4	0.8	1.8	4.9
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	1.1	1.4
2001 .....	0.2	*	*	*	*	*	*	*	*	0.4	0.9	1.7
2000 .....	0.1	*	*	*	*	*	*	*	*	0.3	1.1	1.4
1999 .....	0.2	*	*	*	*	*	*	*	*	0.3	1.0	2.2
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	0.5	*	*	*	*	*	0.4	0.8	1.2	2.6	3.9	*
2001 .....	0.6	*	*	*	*	*	*	0.8	1.5	2.7	3.8	6.9
2000 .....	0.7	*	*	*	*	*	0.5	1.0	1.4	3.2	6.4	*
1999 .....	0.7	*	*	*	*	*	0.4	1.0	1.7	2.6	5.6	8.4
<b>Male</b>												
2002 .....	0.7	*	*	*	*	*	*	1.1	2.0	3.7	6.4	*
2001 .....	0.7	*	*	*	*	*	*	1.4	2.4	4.4	*	*
2000 .....	0.9	*	*	*	*	*	*	1.9	2.5	4.2	11.7	*
1999 .....	0.9	*	*	*	*	*	*	1.8	3.0	4.2	7.8	*
<b>Female</b>												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	3.5	*
2000 .....	0.4	*	*	*	*	*	*	*	*	2.5	3.5	*
1999 .....	0.5	*	*	*	*	*	*	*	*	*	4.4	*
<b>Respiratory tuberculosis (A16)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.5	2.3
2001 .....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4
2000 .....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.7	2.3
1999 .....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.8	3.7
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.9	3.5
2001 .....	0.3	*	*	*	*	*	*	0.3	0.6	0.9	1.7	3.8
2000 .....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.4	4.1
1999 .....	0.4	*	*	*	*	*	0.1	0.4	0.7	1.2	2.5	6.4
<b>Female</b>												
2002 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.8
2001 .....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
All origins <sup>3</sup>												
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.5	1.2	1.6
1999 .....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	2.6
Hispanic												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	5.0	*
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.5	2.2
2001 .....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.2	2.4
2000 .....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.6	2.2
1999 .....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.6	3.5
Male												
2002 .....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	3.3
2001 .....	0.3	*	*	*	*	*	*	0.2	0.6	0.8	1.6	3.8
2000 .....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	2.2	3.8
1999 .....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.1	2.3	6.1
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.4	1.2	1.7
2001 .....	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.8
2000 .....	0.2	*	*	*	*	*	*	*	*	0.4	1.2	1.5
1999 .....	0.2	*	*	*	*	*	*	*	*	0.4	1.2	2.4
Non-Hispanic white												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.6
2001 .....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.7
2000 .....	0.1	*	*	*	*	*	*	0.1	*	0.3	1.1	1.8
1999 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.8
Male												
2002 .....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.2	2.7
2001 .....	0.2	*	*	*	*	*	*	*	0.4	0.4	1.1	2.4
2000 .....	0.2	*	*	*	*	*	*	0.2	*	0.5	1.3	3.2
1999 .....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.6	4.8

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Respiratory tuberculosis (A16)</b>												
Non-Hispanic white												
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
2001 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.4
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
1999 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.9
Non-Hispanic black												
Both sexes												
2002 .....	0.4	*	*	*	*	*	*	0.6	1.0	1.9	3.6	*
2001 .....	0.4	*	*	*	*	*	*	0.6	1.2	2.0	3.0	6.2
2000 .....	0.5	*	*	*	*	*	0.4	0.8	0.9	2.7	5.1	*
1999 .....	0.6	*	*	*	*	*	*	0.7	1.5	2.0	4.8	7.1
Male												
2002 .....	0.6	*	*	*	*	*	*	1.0	1.8	*	6.2	*
2001 .....	0.6	*	*	*	*	*	*	1.2	2.1	3.4	*	*
2000 .....	0.8	*	*	*	*	*	*	1.5	2.1	3.9	9.3	*
1999 .....	0.7	*	*	*	*	*	*	1.3	2.7	3.2	6.9	*
Female												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.4	*	*	*	*	*	*	*	*	*	3.7	*
<b>Other tuberculosis (A17-A19)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
2001 .....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	*
2000 .....	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
1999 .....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
2001 .....	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	*
1999 .....	0.1	*	*	*	*	*	*	0.1	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
Hispanic												
Male												
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
2001 .....	0.1	*	*	*	*	*	*	0.1	*	0.2	0.3	*
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
1999 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
2001 .....	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001 .....	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
2000 .....	0.0	*	*	*	*	*	*	*	*	0.1	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Male												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
Non-Hispanic black												
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Whooping cough (A37)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Scarlet fever and erysipelas (A38,A46)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Meningococcal infection (A39)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	0.1	*	*	0.0	0.1	*	*	*	*	*	*	*
2001 .....	0.1	*	*	0.1	0.1	*	*	*	*	*	*	*
2000 .....	0.1	0.6	*	*	0.1	*	*	0.1	*	*	*	*
1999 .....	0.1	0.8	0.2	0.1	0.1	0.1	0.1	*	*	*	*	*
<b>Male</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	0.1	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Meningococcal infection (A39)</b>												
<b>All origins <sup>3</sup></b>												
<b>Female</b>												
2002 .....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2001 .....	0.1	*	*	0.1	0.2	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	0.1	*	*	0.1	*	*	*	*
1999 .....	0.1	0.8	0.2	0.1	0.1	*	0.1	*	*	*	*	*
<b>Male</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999 .....	0.1	1.0	0.3	*	0.1	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Meningococcal infection (A39)</b>												
Non-Hispanic white												
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Septicemia (A40-A41)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	11.7	7.3	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.7	86.5	203.0
2001 .....	11.3	7.7	0.7	0.2	0.3	0.7	1.8	5.0	12.3	32.8	82.3	205.9
2000 .....	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7
1999 .....	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7
Male												
2002 .....	10.6	7.7	0.5	0.2	0.3	0.8	2.0	5.7	13.7	38.0	99.0	213.9
2001 .....	10.2	8.4	0.8	0.2	0.4	0.7	2.0	5.4	13.2	36.7	95.5	224.3
2000 .....	9.8	7.1	0.7	0.2	0.3	0.8	2.0	5.6	13.0	34.5	91.9	223.0
1999 .....	9.8	9.0	0.6	0.2	0.3	0.7	2.1	5.1	12.4	35.2	91.9	233.5
Female												
2002 .....	12.9	7.0	0.5	0.2	0.3	0.8	1.8	4.7	11.7	31.9	78.2	198.3
2001 .....	12.4	7.0	0.6	0.1	0.3	0.7	1.7	4.7	11.4	29.5	73.7	198.3
2000 .....	12.3	7.3	0.6	0.1	0.2	0.6	1.9	4.2	11.0	28.1	72.9	212.7
1999 .....	12.2	5.8	0.5	0.2	0.3	0.6	1.6	4.1	10.5	28.0	71.3	215.5
Hispanic												
Both sexes												
2002 .....	3.9	7.8	*	*	*	0.5	1.2	3.7	8.4	28.9	63.7	156.3
2001 .....	3.7	6.1	*	*	*	0.3	1.3	3.9	8.2	25.4	62.4	171.3
2000 .....	3.5	8.0	*	*	*	0.5	1.5	3.9	8.5	21.8	59.3	144.7
1999 .....	3.6	5.7	*	*	*	0.4	1.5	3.1	8.0	25.6	61.7	180.3
Male												
2002 .....	3.7	8.0	*	*	*	0.6	1.6	5.2	9.0	30.8	72.2	164.0

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Septicemia (A40-A41)</b>												
Hispanic												
Male												
2001 .....	3.6	7.0	*	*	*	*	1.5	4.7	9.9	29.8	76.0	160.0
2000 .....	3.4	8.4	*	*	*	0.6	1.7	5.0	10.0	26.6	70.3	157.2
1999 .....	3.5	7.0	*	*	*	*	2.0	3.8	10.2	29.4	75.1	181.2
Female												
2002 .....	4.1	7.6	*	*	*	*	0.7	2.3	7.8	27.4	57.9	152.5
2001 .....	3.9	5.2	*	*	*	*	1.1	3.1	6.6	21.9	53.2	176.9
2000 .....	3.5	7.7	*	*	*	*	1.3	2.8	7.2	18.1	51.8	138.5
1999 .....	3.7	*	*	*	*	*	1.0	2.4	6.1	22.5	52.7	179.9
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	12.9	7.1	0.5	0.2	0.3	0.8	2.0	5.3	12.9	34.9	87.3	204.2
2001 .....	12.4	8.1	0.7	0.1	0.4	0.8	1.9	5.1	12.5	33.1	83.0	206.8
2000 .....	12.1	6.9	0.7	0.2	0.3	0.7	2.0	5.0	12.1	31.4	81.0	217.5
1999 .....	12.0	7.9	0.6	0.2	0.3	0.7	1.8	4.7	11.6	31.4	79.8	221.2
Male												
2002 .....	11.6	7.3	0.5	0.3	0.3	0.9	2.0	5.7	14.0	38.2	99.8	215.6
2001 .....	11.2	8.8	0.9	0.1	0.4	0.8	2.1	5.5	13.4	37.0	96.1	226.8
2000 .....	10.7	6.7	0.7	0.2	0.3	0.8	2.0	5.6	13.1	34.8	92.3	224.8
1999 .....	10.7	9.4	0.6	0.2	0.4	0.8	2.1	5.2	12.5	35.4	92.2	234.5
Female												
2002 .....	14.1	6.8	0.5	0.2	0.3	0.8	2.0	4.9	11.9	32.1	79.0	199.3
2001 .....	13.5	7.4	0.6	0.2	0.3	0.8	1.7	4.8	11.8	29.9	74.4	198.5
2000 .....	13.5	7.0	0.6	0.1	0.2	0.7	1.9	4.3	11.2	28.7	73.6	214.5
1999 .....	13.2	6.2	0.5	0.2	0.3	0.7	1.6	4.2	10.8	28.2	71.8	215.9
Non-Hispanic white												
Both sexes												
2002 .....	12.9	5.5	0.5	0.2	0.3	0.7	1.6	4.3	10.7	31.0	81.3	192.6
2001 .....	12.3	5.1	0.7	0.1	0.3	0.6	1.4	4.0	10.4	29.4	77.6	195.6
2000 .....	11.9	4.8	0.5	0.2	0.2	0.5	1.5	3.8	10.0	27.7	74.2	204.3
1999 .....	11.7	5.3	0.5	0.2	0.3	0.6	1.4	3.6	9.5	27.3	73.0	206.0
Male												
2002 .....	11.5	5.6	0.5	0.3	0.3	0.7	1.6	4.6	11.8	34.0	92.1	203.9
2001 .....	11.0	5.8	0.9	*	0.3	0.5	1.7	4.2	11.1	32.5	89.7	213.6
2000 .....	10.4	4.7	0.5	0.2	0.3	0.6	1.6	4.3	10.9	30.9	84.7	211.2
1999 .....	10.2	6.7	0.5	0.2	0.3	0.7	1.5	4.0	9.9	30.5	84.6	218.1
Female												
2002 .....	14.2	5.4	*	0.2	0.3	0.6	1.7	3.9	9.7	28.5	74.0	187.7
2001 .....	13.6	4.4	0.5	*	0.3	0.6	1.2	3.8	9.9	26.8	69.6	188.1
2000 .....	13.4	4.8	0.6	*	*	0.5	1.5	3.3	9.2	24.9	67.2	201.6
1999 .....	13.1	3.8	0.5	0.2	0.2	0.5	1.2	3.3	9.1	24.5	65.4	201.2
Non-Hispanic black												
Both sexes												
2002 .....	16.8	13.1	*	0.3	0.4	2.0	4.6	13.2	33.8	76.0	172.9	370.8
2001 .....	16.4	20.3	0.9	*	0.7	1.9	4.8	13.7	32.6	72.8	161.9	368.6
2000 .....	16.7	16.6	1.1	*	0.6	2.1	4.8	13.9	31.8	72.5	175.1	395.9
1999 .....	16.6	18.6	*	*	0.6	1.6	5.1	13.2	30.8	75.3	169.8	414.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Septicemia (A40-A41)</b>												
Non-Hispanic black												
Male												
2002 .....	15.5	13.9	*	*	*	1.9	5.0	15.0	36.6	87.6	213.7	399.3
2001 .....	15.4	21.4	*	*	0.9	1.9	4.9	15.2	37.4	86.9	196.8	430.8
2000 .....	15.3	16.4	*	*	*	2.4	4.8	16.8	35.9	81.6	208.9	432.1
1999 .....	15.7	20.8	*	*	*	1.7	6.0	15.4	37.9	91.5	202.6	451.6
Female												
2002 .....	18.0	12.3	*	*	*	2.0	4.2	11.6	31.6	67.8	149.9	359.9
2001 .....	17.2	19.1	*	*	*	1.9	4.7	12.5	28.8	62.8	142.4	345.6
2000 .....	17.9	16.9	*	*	*	1.8	4.9	11.5	28.5	66.1	156.1	382.5
1999 .....	17.4	16.4	*	*	*	1.6	4.3	11.3	25.1	63.9	151.4	400.2
<b>Syphilis (A50-A53)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Acute poliomyelitis (A80)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Acute poliomyelitis (A80)</b>												
<b>Non-Hispanic white</b>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic black</b>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Arthropod-borne viral encephalitis (A83-A84,A85.2)</b>												
<b>All origins <sup>3</sup></b>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
All origins <sup>3</sup>												
Female												
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
Non-Hispanic white												
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
Hispanic												
Male												
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
Non-Hispanic black												
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	2.0	*	*	*	0.1	0.2	1.6	5.8	4.0	4.2	4.6	3.9
2001 .....	2.0	*	*	*	0.1	0.2	1.8	5.3	4.0	4.6	4.9	3.8
2000 .....	1.9	*	*	*	*	0.2	1.7	5.0	3.7	4.6	5.3	4.0
1999 .....	1.7	*	*	*	*	0.2	1.6	4.1	3.3	4.8	5.3	4.7
Male												
2002 .....	2.7	*	*	*	*	0.3	2.3	8.9	5.5	4.5	5.3	5.0
2001 .....	2.5	*	*	*	*	0.2	2.5	8.2	5.3	4.8	5.3	4.4
2000 .....	2.4	*	*	*	*	0.2	2.4	7.7	5.0	4.8	6.4	5.7
1999 .....	2.2	*	*	*	*	0.3	2.3	6.3	4.3	5.1	6.6	6.9
Female												
2002 .....	1.4	*	*	*	*	0.2	1.0	2.8	2.7	3.9	4.1	3.3
2001 .....	1.4	*	*	*	*	0.2	1.1	2.5	2.8	4.4	4.6	3.5
2000 .....	1.4	*	*	*	*	0.2	1.1	2.4	2.6	4.5	4.7	3.3
1999 .....	1.3	*	*	*	*	0.2	0.9	2.0	2.4	4.5	4.5	3.9
Hispanic												
Both sexes												
2002 .....	2.0	*	*	*	*	*	1.7	8.0	8.3	8.6	10.5	*
2001 .....	2.1	*	*	*	*	*	2.3	7.8	10.4	9.9	11.0	*
2000 .....	2.0	*	*	*	*	*	2.5	7.0	8.6	11.1	13.0	*
1999 .....	1.9	*	*	*	*	*	2.1	6.8	7.8	12.4	13.5	*
Male												
2002 .....	2.5	*	*	*	*	*	2.5	12.0	11.6	7.6	10.3	*
2001 .....	2.6	*	*	*	*	*	3.1	12.2	14.2	7.0	10.3	*
2000 .....	2.5	*	*	*	*	*	3.8	10.6	11.7	11.0	12.3	*
1999 .....	2.3	*	*	*	*	*	3.3	10.5	9.8	13.0	*	*
Female												
2002 .....	1.4	*	*	*	*	*	0.9	4.0	5.4	9.4	10.7	*
2001 .....	1.7	*	*	*	*	*	1.4	3.5	7.1	12.3	11.5	*
2000 .....	1.5	*	*	*	*	*	1.0	3.3	5.8	11.3	13.5	*
1999 .....	1.5	*	*	*	*	*	0.8	3.3	6.0	12.0	16.3	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	2.0	*	*	*	*	0.2	1.6	5.6	3.6	3.8	4.3	3.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Viral hepatitis (B15-B19)</b>												
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2001 .....	1.9	*	*	*	*	0.2	1.7	5.0	3.5	4.2	4.6	3.6
2000 .....	1.9	*	*	*	*	0.2	1.6	4.8	3.3	4.2	5.0	3.8
1999 .....	1.7	*	*	*	*	0.2	1.5	3.8	3.0	4.3	5.0	4.7
<b>Male</b>												
2002 .....	2.7	*	*	*	*	0.3	2.3	8.6	4.9	4.3	5.1	4.8
2001 .....	2.5	*	*	*	*	0.2	2.4	7.7	4.6	4.6	5.0	4.3
2000 .....	2.4	*	*	*	*	0.2	2.2	7.4	4.4	4.4	6.1	5.4
1999 .....	2.2	*	*	*	*	0.2	2.2	5.9	3.9	4.7	6.4	6.8
<b>Female</b>												
2002 .....	1.4	*	*	*	*	0.2	1.0	2.7	2.4	3.5	3.8	3.2
2001 .....	1.4	*	*	*	*	0.2	1.0	2.3	2.5	3.8	4.3	3.3
2000 .....	1.4	*	*	*	*	0.2	1.0	2.4	2.3	4.0	4.3	3.2
1999 .....	1.2	*	*	*	*	0.2	0.9	1.8	2.1	4.0	4.0	3.9
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	1.9	*	*	*	*	0.2	1.6	5.3	3.0	3.3	4.1	3.5
2001 .....	1.8	*	*	*	*	0.2	1.7	4.6	2.9	3.7	4.3	3.3
2000 .....	1.8	*	*	*	*	0.1	1.6	4.3	2.7	3.7	4.7	3.7
1999 .....	1.6	*	*	*	*	0.2	1.4	3.4	2.5	3.7	4.7	4.5
<b>Male</b>												
2002 .....	2.6	*	*	*	*	0.3	2.3	8.2	4.1	3.9	4.9	4.6
2001 .....	2.5	*	*	*	*	0.2	2.3	7.2	4.0	4.3	5.1	4.4
2000 .....	2.3	*	*	*	*	*	2.1	6.7	3.6	3.8	5.9	5.4
1999 .....	2.1	*	*	*	*	0.2	2.1	5.2	3.4	4.3	6.4	6.5
<b>Female</b>												
2002 .....	1.3	*	*	*	*	*	0.9	2.5	2.0	2.8	3.5	3.1
2001 .....	1.2	*	*	*	*	*	1.0	2.0	2.0	3.1	3.8	2.9
2000 .....	1.3	*	*	*	*	*	1.0	2.0	1.8	3.6	4.0	3.1
1999 .....	1.1	*	*	*	*	0.2	0.8	1.7	1.6	3.3	3.6	3.7
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	2.3	*	*	*	*	0.4	2.0	8.2	7.2	6.0	5.2	*
2001 .....	2.3	*	*	*	*	*	2.2	8.1	6.8	5.6	5.7	*
2000 .....	2.2	*	*	*	*	0.4	2.1	8.0	5.8	5.8	6.3	*
1999 .....	2.1	*	*	*	*	*	2.2	7.0	5.1	6.8	5.9	*
<b>Male</b>												
2002 .....	3.1	*	*	*	*	*	2.6	12.6	10.7	7.0	*	*
2001 .....	2.8	*	*	*	*	*	3.0	12.0	9.0	5.0	*	*
2000 .....	2.9	*	*	*	*	*	3.0	12.3	8.1	7.2	7.7	*
1999 .....	2.6	*	*	*	*	*	3.1	11.8	6.3	5.7	*	*
<b>Female</b>												
2002 .....	1.7	*	*	*	*	*	1.5	4.4	4.4	5.2	5.3	*
2001 .....	1.8	*	*	*	*	*	1.4	4.7	5.0	6.0	6.7	*
2000 .....	1.6	*	*	*	*	*	1.4	4.3	3.9	4.7	5.6	*
1999 .....	1.6	*	*	*	*	*	1.4	2.9	4.1	7.5	6.7	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	4.9	*	*	0.1	0.4	4.6	12.7	11.2	5.1	2.2	0.8	*
2001 .....	5.0	*	*	0.1	0.6	5.3	13.0	10.5	5.2	2.1	0.7	*
2000 .....	5.1	*	*	0.1	0.5	6.1	13.1	11.0	5.1	2.2	0.7	*
1999 .....	5.3	*	0.2	0.2	0.5	6.8	13.8	10.7	4.8	2.2	0.6	*
Male												
2002 .....	7.4	*	*	*	0.4	5.9	18.8	17.7	8.5	3.9	1.4	*
2001 .....	7.6	*	*	0.1	0.5	7.1	19.5	16.8	8.6	3.5	1.5	*
2000 .....	7.9	*	*	0.1	0.5	8.0	19.8	17.8	8.7	3.8	1.3	*
1999 .....	8.2	*	*	0.2	0.5	9.5	21.0	17.5	8.3	3.8	1.0	*
Female												
2002 .....	2.5	*	*	*	0.4	3.3	6.7	4.8	1.9	0.8	0.3	*
2001 .....	2.5	*	*	*	0.6	3.5	6.7	4.4	2.0	0.9	*	*
2000 .....	2.5	*	*	0.1	0.4	4.2	6.5	4.4	1.8	0.8	0.3	*
1999 .....	2.5	*	*	0.2	0.5	4.1	6.7	4.1	1.6	0.8	0.3	*
Hispanic												
Both sexes												
2002 .....	4.8	*	*	*	0.4	3.6	13.4	14.7	9.5	3.4	*	*
2001 .....	5.0	*	*	*	0.5	4.4	13.9	15.5	9.2	4.5	*	*
2000 .....	5.4	*	*	*	0.3	5.6	14.9	16.9	9.5	3.8	*	*
1999 .....	5.8	*	*	*	0.5	6.4	17.3	15.4	8.4	4.7	*	*
Male												
2002 .....	7.2	*	*	*	*	5.1	20.1	23.1	15.0	6.9	*	*
2001 .....	7.6	*	*	*	0.6	6.0	20.8	24.9	14.8	8.6	*	*
2000 .....	8.2	*	*	*	*	7.8	22.7	27.5	15.3	7.2	*	*
1999 .....	8.7	*	*	*	0.6	9.3	26.2	24.2	15.4	9.0	*	*
Female												
2002 .....	2.2	*	*	*	*	1.8	6.2	6.3	4.6	*	*	*
2001 .....	2.3	*	*	*	*	2.5	6.5	6.2	4.1	*	*	*
2000 .....	2.4	*	*	*	*	3.0	6.4	6.6	4.4	*	*	*
1999 .....	2.6	*	*	*	*	3.2	7.9	6.7	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	4.8	*	*	0.1	0.4	4.8	12.5	10.6	4.7	2.1	0.7	*
2001 .....	4.9	*	*	0.1	0.6	5.4	12.7	9.9	4.8	1.9	0.6	*
2000 .....	5.0	*	*	0.1	0.5	6.2	12.7	10.3	4.7	2.0	0.6	*
1999 .....	5.2	*	*	0.2	0.5	6.8	13.2	10.0	4.5	2.0	0.6	*
Male												
2002 .....	7.3	*	*	*	0.4	6.0	18.4	16.9	7.9	3.7	1.3	*
2001 .....	7.5	*	*	0.1	0.5	7.3	19.0	15.8	8.0	3.1	1.3	*
2000 .....	7.7	*	*	*	0.5	7.9	19.1	16.7	8.0	3.4	1.2	*
1999 .....	8.0	*	*	0.2	0.5	9.4	20.1	16.5	7.6	3.4	1.0	*
Female												
2002 .....	2.5	*	*	*	0.5	3.6	6.7	4.7	1.7	0.8	0.3	*
2001 .....	2.5	*	*	*	0.7	3.6	6.6	4.2	1.8	0.9	*	*
2000 .....	2.5	*	*	0.1	0.5	4.4	6.4	4.2	1.6	0.8	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
Non-Hispanic <sup>4</sup>												
Female												
1999 .....	2.5	*	*	0.1	0.6	4.2	6.5	3.9	1.6	0.7	0.3	*
Non-Hispanic white												
Both sexes												
2002 .....	2.1	*	*	*	0.1	1.7	5.8	4.6	2.2	0.8	0.3	*
2001 .....	2.2	*	*	*	0.1	2.1	6.0	4.2	2.1	0.7	0.3	*
2000 .....	2.3	*	*	*	0.1	2.4	6.2	4.6	2.2	0.8	0.3	*
1999 .....	2.4	*	*	*	0.1	3.0	6.4	4.6	2.0	0.8	0.3	*
Male												
2002 .....	3.7	*	*	*	*	2.7	9.8	8.0	4.0	1.5	0.5	*
2001 .....	3.8	*	*	*	*	3.2	10.4	7.5	3.8	1.4	0.6	*
2000 .....	4.0	*	*	*	0.2	3.7	10.4	8.2	4.0	1.5	0.6	*
1999 .....	4.2	*	*	*	0.2	4.9	10.9	8.2	3.6	1.5	0.6	*
Female												
2002 .....	0.6	*	*	*	*	0.8	1.8	1.2	0.5	*	*	*
2001 .....	0.6	*	*	*	*	0.9	1.7	1.0	0.5	*	*	*
2000 .....	0.7	*	*	*	*	1.1	2.0	1.1	0.5	*	*	*
1999 .....	0.7	*	*	*	*	1.2	1.9	1.1	0.4	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	21.3	*	*	*	2.1	21.5	54.0	54.8	26.3	14.6	5.7	*
2001 .....	21.6	*	*	0.4	2.8	24.0	55.0	51.9	28.0	12.8	5.3	*
2000 .....	22.0	*	*	0.4	2.2	27.2	54.5	53.9	26.3	13.2	4.6	*
1999 .....	22.3	*	*	0.6	2.3	28.3	57.2	52.2	26.4	13.2	3.7	*
Male												
2002 .....	30.3	*	*	*	2.0	25.5	75.6	86.2	44.8	26.6	12.3	*
2001 .....	30.8	*	*	*	2.4	30.8	77.8	81.4	47.7	21.4	11.4	*
2000 .....	31.7	*	*	*	2.2	33.4	79.0	86.6	46.4	23.8	8.9	*
1999 .....	32.6	*	*	0.7	1.9	37.1	83.5	85.5	45.6	24.0	6.6	*
Female												
2002 .....	13.2	*	*	*	2.2	17.9	34.9	28.0	11.5	6.0	*	*
2001 .....	13.3	*	*	*	3.2	17.8	34.8	26.6	12.1	6.7	*	*
2000 .....	13.1	*	*	*	2.3	21.7	32.8	25.8	10.1	5.8	*	*
1999 .....	13.0	*	*	*	2.8	20.3	34.1	23.8	11.0	5.6	*	*
Malaria (B50-B54)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malaria (B50-B54)</b>												
<b>All origins <sup>3</sup></b>												
Male												
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Hispanic</b>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic white</b>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malaria (B50-B54)</b>												
<b>Non-Hispanic white</b>												
<b>Male</b>												
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8, A86-B04,B06-B09,B25-B49,B55-B99)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	2.3	5.6	0.4	0.2	0.2	0.4	1.0	2.3	3.3	6.7	12.5	21.7
2001 .....	2.0	4.7	0.4	0.2	0.2	0.4	0.8	1.7	3.0	6.2	10.6	20.1
2000 .....	2.0	4.4	0.5	0.2	0.2	0.4	0.8	1.6	2.9	5.8	11.1	21.2
1999 .....	2.6	5.3	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0
<b>Male</b>												
2002 .....	2.6	6.1	0.5	0.2	0.2	0.5	1.3	3.2	4.2	8.0	15.4	26.8
2001 .....	2.1	4.9	0.5	0.1	0.2	0.4	1.0	2.2	3.7	6.8	13.2	23.6
2000 .....	2.1	4.8	0.5	0.2	0.3	0.4	1.0	1.9	3.3	7.2	13.6	27.6
1999 .....	2.7	6.1	0.6	0.2	0.2	0.6	1.5	2.7	4.2	9.2	17.2	31.9
<b>Female</b>												
2002 .....	2.1	5.0	0.3	0.2	0.2	0.4	0.7	1.4	2.6	5.6	10.6	19.5
2001 .....	1.9	4.6	0.4	0.2	0.2	0.3	0.6	1.2	2.4	5.8	8.9	18.7
2000 .....	1.9	3.9	0.4	0.2	0.2	0.3	0.6	1.2	2.6	4.7	9.5	18.6
1999 .....	2.5	4.5	0.4	0.2	0.2	0.6	0.9	1.9	3.4	6.7	12.3	25.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8, A86-B04,B06-B09,B25-B49,B55-B99)												
Hispanic												
Both sexes												
2002 .....	1.5	4.9	*	*	*	0.7	1.2	3.5	4.3	6.4	11.3	17.7
2001 .....	1.1	5.0	*	*	*	0.5	0.8	2.0	3.0	5.9	10.3	15.0
2000 .....	1.2	3.0	*	*	*	0.4	1.0	1.9	3.2	5.8	11.3	19.2
1999 .....	1.5	4.8	*	*	*	0.6	1.4	2.3	3.8	7.7	14.2	26.0
Male												
2002 .....	1.8	6.3	*	*	*	0.8	1.6	5.2	5.6	7.6	12.3	*
2001 .....	1.3	5.0	*	*	*	0.6	1.1	2.8	4.4	7.6	12.1	*
2000 .....	1.3	*	*	*	*	0.6	1.4	2.1	4.4	5.9	13.3	*
1999 .....	1.5	5.7	*	*	*	0.6	1.9	2.7	3.8	7.9	15.1	*
Female												
2002 .....	1.2	*	*	*	*	0.6	*	1.7	3.2	5.3	10.7	*
2001 .....	1.0	5.0	*	*	*	*	*	1.3	*	4.6	9.1	*
2000 .....	1.0	*	*	*	*	*	*	1.8	*	5.6	9.9	*
1999 .....	1.4	*	*	*	*	*	*	1.9	3.8	7.6	13.5	28.1
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	2.4	5.7	0.4	0.2	0.2	0.4	1.0	2.2	3.3	6.7	12.5	21.8
2001 .....	2.1	4.6	0.5	0.2	0.2	0.3	0.8	1.6	3.0	6.2	10.6	20.3
2000 .....	2.1	4.7	0.4	0.2	0.2	0.3	0.7	1.5	2.9	5.8	11.1	21.2
1999 .....	2.8	5.5	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0
Male												
2002 .....	2.7	5.9	0.5	0.2	0.3	0.4	1.3	3.0	4.0	8.0	15.6	26.8
2001 .....	2.2	4.8	0.5	0.1	0.2	0.4	1.0	2.1	3.6	6.7	13.2	24.0
2000 .....	2.2	5.1	0.5	0.2	0.3	0.3	0.9	1.9	3.2	7.3	13.5	27.8
1999 .....	2.8	6.2	0.6	0.2	0.3	0.6	1.4	2.6	4.2	9.2	17.2	32.3
Female												
2002 .....	2.2	5.4	*	0.2	0.2	0.3	0.7	1.4	2.5	5.6	10.5	19.7
2001 .....	2.1	4.5	0.4	0.2	0.2	0.3	0.6	1.2	2.5	5.8	8.9	18.8
2000 .....	2.0	4.3	0.4	0.2	0.2	0.4	0.6	1.1	2.6	4.7	9.5	18.6
1999 .....	2.7	4.7	0.4	0.2	0.2	0.6	1.0	1.9	3.3	6.7	12.2	24.9
Non-Hispanic white												
Both sexes												
2002 .....	2.5	4.8	0.4	0.2	0.2	0.3	0.8	1.9	2.9	6.6	12.4	22.2
2001 .....	2.1	3.3	0.4	0.1	0.2	0.3	0.7	1.5	2.8	6.0	10.4	20.2
2000 .....	2.1	3.5	0.4	0.2	0.2	0.2	0.6	1.3	2.6	5.6	10.9	21.0
1999 .....	2.7	4.4	0.5	0.2	0.2	0.4	0.9	1.8	3.3	7.4	13.6	26.4
Male												
2002 .....	2.7	4.8	0.4	0.2	0.2	0.3	1.0	2.7	3.5	7.9	15.3	27.2
2001 .....	2.2	3.3	0.4	*	0.2	0.3	0.8	1.9	3.3	6.5	13.1	24.2
2000 .....	2.2	3.9	*	*	0.2	0.2	0.7	1.7	3.0	7.0	13.2	27.0
1999 .....	2.7	5.5	0.6	0.2	0.2	0.4	1.2	2.2	3.8	8.8	16.3	30.8
Female												
2002 .....	2.3	4.9	*	0.2	0.2	0.3	0.5	1.2	2.3	5.5	10.5	20.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8, A86-B04,B06-B09,B25-B49,B55-B99)												
Non-Hispanic white												
Female												
2001 .....	2.1	3.4	*	0.2	0.2	0.2	0.5	1.0	2.3	5.7	8.7	18.6
2000 .....	2.0	3.1	*	0.2	0.2	0.2	0.5	1.0	2.3	4.5	9.5	18.6
1999 .....	2.6	3.3	*	*	0.2	0.4	0.6	1.5	2.9	6.3	11.8	24.7
Non-Hispanic black												
Both sexes												
2002 .....	2.6	9.0	*	*	0.4	0.9	2.2	4.0	5.9	7.4	13.8	21.8
2001 .....	2.3	10.0	*	*	0.4	0.9	1.6	3.0	5.2	8.3	11.3	20.9
2000 .....	2.3	10.4	*	*	0.4	1.0	1.7	3.0	4.9	8.0	13.3	24.2
1999 .....	3.6	10.0	1.0	*	0.4	1.7	3.1	5.6	7.3	11.9	20.9	34.3
Male												
2002 .....	3.0	10.0	*	*	*	1.1	2.6	5.6	8.7	9.1	17.0	28.4
2001 .....	2.5	11.5	*	*	*	*	2.1	4.0	6.4	9.8	12.6	*
2000 .....	2.6	10.6	*	*	*	0.9	2.2	4.0	5.8	10.5	17.9	31.7
1999 .....	3.7	10.0	*	*	*	1.9	3.1	6.3	7.9	14.1	29.4	46.4
Female												
2002 .....	2.2	7.9	*	*	*	*	1.8	2.6	3.7	6.2	11.9	19.2
2001 .....	2.1	8.4	*	*	*	1.0	1.2	2.1	4.3	7.2	10.6	20.5
2000 .....	2.1	10.2	*	*	*	1.0	1.3	2.2	4.1	6.2	10.8	21.4
1999 .....	3.5	10.0	*	*	*	1.5	3.0	4.9	6.7	10.4	16.1	29.8
Malignant neoplasms (C00-C97)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	193.2	1.8	2.6	2.6	4.3	9.7	35.8	123.8	351.1	792.1	1,311.9	1,723.9
2001 .....	194.4	1.6	2.7	2.5	4.3	10.1	36.8	126.5	356.5	802.8	1,315.8	1,765.6
2000 .....	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4
1999 .....	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8
Male												
2002 .....	203.8	2.0	2.7	2.9	4.9	9.2	31.5	128.0	399.8	964.8	1,711.3	2,491.1
2001 .....	205.3	1.5	2.9	2.5	5.0	9.3	32.6	130.3	405.2	984.6	1,727.1	2,613.6
2000 .....	207.2	2.6	3.0	2.7	5.1	9.2	32.7	130.9	415.8	1,001.9	1,760.6	2,710.7
1999 .....	208.9	1.9	2.9	2.7	5.2	9.7	33.3	131.7	428.4	1,019.4	1,767.0	2,722.2
Female												
2002 .....	183.0	1.6	2.4	2.4	3.6	10.2	40.0	119.8	306.0	648.5	1,046.7	1,391.1
2001 .....	183.9	1.8	2.5	2.4	3.5	10.9	41.0	122.7	311.5	652.2	1,045.4	1,410.7
2000 .....	186.2	2.3	2.5	2.2	3.6	10.4	40.4	124.2	321.3	663.6	1,058.5	1,456.4
1999 .....	185.6	1.7	2.5	2.3	3.7	10.3	40.9	123.7	325.0	669.5	1,048.5	1,436.5
Hispanic												
Both sexes												
2002 .....	59.7	*	2.3	2.9	4.5	6.8	23.2	78.2	215.1	496.0	894.9	1,308.8
2001 .....	60.5	*	2.8	2.5	4.3	7.2	25.6	82.8	223.4	504.0	913.9	1,363.7
2000 .....	59.9	*	2.5	2.9	4.2	7.3	25.2	82.1	221.0	516.2	953.5	1,383.5
1999 .....	59.6	*	2.5	2.7	4.1	7.4	24.4	84.9	225.1	520.5	926.8	1,399.6

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms (C00-C97)</b>												
Hispanic												
Male												
2002 .....	61.2	*	2.2	2.9	5.2	6.3	18.4	78.4	254.3	622.3	1,190.8	1,869.0
2001 .....	62.2	*	2.9	2.5	5.5	6.3	21.7	81.5	253.5	642.8	1,258.3	1,967.4
2000 .....	61.3	*	2.9	3.5	4.5	6.9	20.1	79.4	253.1	651.2	1,306.4	2,049.7
1999 .....	61.2	*	2.3	2.8	4.6	6.6	19.8	84.6	268.1	668.9	1,252.9	1,945.3
Female												
2002 .....	58.1	*	2.4	2.9	3.6	7.5	28.4	78.0	179.8	395.6	692.2	1,031.2
2001 .....	58.7	*	2.7	2.5	2.9	8.3	29.8	84.2	196.7	394.5	681.2	1,068.6
2000 .....	58.5	*	2.2	2.3	3.7	7.8	30.7	84.7	192.5	410.0	716.5	1,056.5
1999 .....	58.0	*	2.7	2.6	3.5	8.3	29.4	85.1	187.2	404.3	709.8	1,136.5
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	213.4	1.7	2.6	2.5	4.2	10.3	37.6	128.0	360.9	810.5	1,329.2	1,736.8
2001 .....	213.9	1.6	2.7	2.4	4.2	10.6	38.2	130.3	365.9	820.5	1,331.4	1,776.6
2000 .....	215.6	2.5	2.7	2.4	4.4	10.3	37.9	131.2	376.6	832.8	1,348.9	1,831.1
1999 .....	215.5	1.7	2.7	2.4	4.5	10.4	38.6	131.0	384.6	843.2	1,344.5	1,816.1
Male												
2002 .....	226.7	1.9	2.8	2.8	4.8	9.9	33.5	132.6	410.0	985.1	1,733.2	2,513.2
2001 .....	227.2	1.6	2.9	2.5	4.9	9.9	34.0	134.6	415.3	1,003.8	1,745.1	2,634.8
2000 .....	228.7	2.6	2.9	2.5	5.2	9.7	34.3	135.1	426.6	1,020.2	1,776.0	2,732.4
1999 .....	229.9	1.6	3.0	2.6	5.3	10.3	35.0	135.4	438.8	1,036.8	1,783.7	2,747.5
Female												
2002 .....	200.8	1.5	2.4	2.2	3.6	10.6	41.6	123.5	315.4	664.7	1,061.5	1,401.8
2001 .....	201.2	1.6	2.5	2.3	3.6	11.3	42.4	126.0	320.0	668.1	1,059.8	1,419.6
2000 .....	203.1	2.4	2.6	2.2	3.6	10.9	41.4	127.3	330.3	678.1	1,070.6	1,466.6
1999 .....	201.9	1.8	2.5	2.2	3.7	10.5	42.1	126.8	334.5	684.2	1,059.4	1,443.2
Non-Hispanic white												
Both sexes												
2002 .....	230.9	1.6	2.6	2.5	4.3	10.0	36.2	122.3	353.0	809.2	1,328.4	1,737.9
2001 .....	230.6	1.8	2.9	2.4	4.3	10.3	36.5	124.6	358.3	815.9	1,326.5	1,775.2
2000 .....	232.2	2.6	2.8	2.4	4.3	10.1	35.6	125.1	369.3	827.7	1,342.9	1,826.6
1999 .....	231.5	1.6	3.0	2.3	4.3	10.2	36.6	124.8	375.0	837.1	1,336.4	1,808.3
Male												
2002 .....	243.8	2.0	2.8	2.7	4.9	9.8	32.5	125.9	395.5	975.3	1,716.5	2,507.7
2001 .....	243.4	1.7	3.1	2.4	4.9	10.0	32.6	127.3	401.7	988.2	1,721.8	2,616.8
2000 .....	244.4	2.9	3.1	2.6	5.0	9.7	32.3	127.2	412.0	1,002.1	1,750.2	2,714.1
1999 .....	245.1	1.7	3.3	2.6	5.0	10.1	33.5	126.7	422.3	1,019.3	1,757.7	2,721.3
Female												
2002 .....	218.5	*	2.3	2.3	3.6	10.3	39.9	118.7	312.8	667.7	1,068.3	1,405.4
2001 .....	218.4	1.8	2.8	2.4	3.6	10.7	40.4	121.9	317.3	669.7	1,064.4	1,425.1
2000 .....	220.6	2.4	2.4	2.3	3.5	10.5	38.9	123.0	328.9	681.0	1,075.3	1,468.7
1999 .....	218.4	*	2.7	2.1	3.6	10.3	39.7	123.0	330.5	684.6	1,060.5	1,444.2
Non-Hispanic black												
Both sexes												
2002 .....	171.5	*	3.0	2.6	4.1	12.8	51.5	186.0	494.6	977.2	1,537.4	1,904.3
2001 .....	172.8	*	2.0	2.7	4.4	13.4	52.8	188.7	494.5	1,005.6	1,559.5	1,947.5
2000 .....	174.9	*	2.8	2.3	5.3	12.3	55.3	196.7	505.2	1,019.3	1,582.8	2,015.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms (C00-C97)</b>												
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
1999 .....	176.9	*	2.0	2.8	5.6	12.6	55.3	196.8	530.5	1,033.6	1,592.9	2,024.0
<b>Male</b>												
2002 .....	187.8	*	2.7	3.1	4.6	11.7	44.3	202.3	621.8	1,293.5	2,247.6	3,008.4
2001 .....	190.8	*	2.4	3.0	5.1	10.9	45.8	209.6	614.1	1,354.7	2,310.5	3,238.5
2000 .....	194.9	*	2.6	2.6	6.4	10.5	49.6	219.0	637.4	1,382.7	2,372.3	3,290.1
1999 .....	197.4	*	2.0	2.9	6.6	11.9	48.6	225.8	670.9	1,387.0	2,377.4	3,361.1
<b>Female</b>												
2002 .....	156.7	*	3.2	2.0	3.6	13.9	57.8	172.1	392.5	752.8	1,135.6	1,482.4
2001 .....	156.5	*	*	2.3	3.8	15.7	59.0	170.7	398.3	759.4	1,137.8	1,469.3
2000 .....	156.9	*	3.0	1.9	4.2	14.0	60.4	177.5	398.8	763.6	1,138.4	1,541.6
1999 .....	158.3	*	2.0	2.7	4.8	13.3	61.3	172.0	417.9	784.2	1,153.6	1,524.4
<b>Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	2.7	*	*	*	0.1	0.1	0.6	2.7	6.8	10.4	13.5	18.7
2001 .....	2.7	*	*	*	0.1	0.1	0.6	2.7	6.9	10.4	14.2	18.5
2000 .....	2.7	*	*	*	0.1	0.1	0.6	2.8	6.8	10.7	13.2	19.4
1999 .....	2.7	*	*	*	0.1	0.1	0.6	2.9	6.8	10.3	14.3	19.3
<b>Male</b>												
2002 .....	3.7	*	*	*	*	0.2	0.8	4.4	10.9	15.9	19.6	26.9
2001 .....	3.7	*	*	*	*	0.2	0.9	4.4	11.3	16.0	20.6	25.8
2000 .....	3.6	*	*	*	*	0.1	1.0	4.5	10.9	16.0	19.2	27.1
1999 .....	3.6	*	*	*	0.1	0.1	0.9	4.6	10.8	15.8	20.9	27.1
<b>Female</b>												
2002 .....	1.7	*	*	*	*	*	0.4	1.1	3.0	5.8	9.5	15.1
2001 .....	1.8	*	*	*	*	0.1	0.4	1.2	2.8	5.8	10.0	15.4
2000 .....	1.8	*	*	*	*	0.1	0.3	1.3	2.9	6.3	9.2	16.3
1999 .....	1.8	*	*	*	*	*	0.4	1.3	3.1	5.8	9.9	16.1
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	0.7	*	*	*	*	*	*	1.3	3.3	5.3	9.0	16.1
2001 .....	0.8	*	*	*	*	*	*	1.6	3.7	6.9	10.5	19.4
2000 .....	0.8	*	*	*	*	*	*	1.3	4.4	6.8	9.5	18.6
1999 .....	0.8	*	*	*	*	*	*	1.4	4.0	6.9	13.1	14.0
<b>Male</b>												
2002 .....	1.1	*	*	*	*	*	*	2.1	5.7	9.7	14.8	*
2001 .....	1.2	*	*	*	*	*	*	2.4	6.3	11.6	17.4	*
2000 .....	1.2	*	*	*	*	*	*	2.2	7.7	12.2	13.8	40.3
1999 .....	1.2	*	*	*	*	*	*	2.3	7.2	13.6	18.2	*
<b>Female</b>												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.5	*	*	*	*	*	*	*	*	3.1	*	*
2000 .....	0.4	*	*	*	*	*	*	*	*	*	6.6	*
1999 .....	0.5	*	*	*	*	*	*	*	*	*	9.7	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)</b>												
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	3.0	*	*	*	0.1	0.2	0.6	2.9	7.1	10.7	13.7	18.7
2001 .....	3.0	*	*	*	0.1	0.2	0.7	2.8	7.1	10.6	14.3	18.4
2000 .....	2.9	*	*	*	0.1	0.1	0.7	3.0	6.9	10.8	13.3	19.5
1999 .....	2.9	*	*	*	0.1	0.1	0.7	3.0	7.0	10.5	14.3	19.4
<b>Male</b>												
2002 .....	4.1	*	*	*	*	0.2	0.8	4.6	11.3	16.2	19.8	26.7
2001 .....	4.1	*	*	*	*	0.2	1.0	4.6	11.6	16.2	20.6	25.8
2000 .....	3.9	*	*	*	*	0.1	1.0	4.6	11.1	16.1	19.4	26.5
1999 .....	4.0	*	*	*	0.1	0.2	1.0	4.8	11.0	15.8	21.0	27.3
<b>Female</b>												
2002 .....	1.9	*	*	*	*	*	0.4	1.2	3.2	6.1	9.7	15.3
2001 .....	1.9	*	*	*	*	0.1	0.4	1.2	2.9	6.0	10.2	15.3
2000 .....	2.0	*	*	*	*	0.1	0.4	1.3	3.0	6.5	9.3	16.6
1999 .....	2.0	*	*	*	*	*	0.4	1.4	3.2	6.0	9.9	16.2
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	3.1	*	*	*	*	0.1	0.5	2.5	6.5	10.6	13.9	18.9
2001 .....	3.1	*	*	*	*	0.1	0.5	2.5	6.5	10.2	14.5	18.9
2000 .....	3.0	*	*	*	*	0.1	0.6	2.5	6.1	10.4	13.3	19.8
1999 .....	3.0	*	*	*	*	0.1	0.5	2.4	6.3	10.1	14.2	19.9
<b>Male</b>												
2002 .....	4.1	*	*	*	*	0.2	0.7	4.0	10.2	15.7	19.9	26.8
2001 .....	4.1	*	*	*	*	0.2	0.8	4.0	10.4	15.3	20.5	26.4
2000 .....	3.9	*	*	*	*	*	0.8	3.8	9.7	15.0	18.9	27.4
1999 .....	3.9	*	*	*	*	*	0.8	3.8	9.7	15.1	20.3	27.5
<b>Female</b>												
2002 .....	2.1	*	*	*	*	*	0.3	1.0	2.9	6.2	9.8	15.5
2001 .....	2.1	*	*	*	*	*	0.3	1.0	2.8	5.9	10.5	15.7
2000 .....	2.1	*	*	*	*	*	0.4	1.2	2.8	6.5	9.6	16.8
1999 .....	2.1	*	*	*	*	*	0.3	1.1	3.0	5.9	10.1	16.8
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	2.9	*	*	*	*	*	1.1	5.6	12.4	12.9	12.1	16.0
2001 .....	3.0	*	*	*	*	*	1.3	5.3	12.9	15.7	13.9	13.1
2000 .....	3.1	*	*	*	*	*	1.1	6.1	13.7	15.7	15.1	14.9
1999 .....	3.3	*	*	*	*	*	1.2	7.3	13.6	15.4	16.8	14.5
<b>Male</b>												
2002 .....	4.4	*	*	*	*	*	1.6	9.8	22.5	23.0	18.8	26.3
2001 .....	4.6	*	*	*	*	*	1.7	8.9	23.9	27.3	23.4	*
2000 .....	4.9	*	*	*	*	*	1.8	10.7	24.7	28.6	29.0	*
1999 .....	5.0	*	*	*	*	*	1.8	12.3	23.5	25.4	31.9	25.0
<b>Female</b>												
2002 .....	1.5	*	*	*	*	*	0.7	2.1	4.3	5.8	8.3	12.1
2001 .....	1.5	*	*	*	*	*	0.9	2.1	4.0	7.5	8.6	10.7
2000 .....	1.5	*	*	*	*	*	*	2.1	4.9	6.6	7.3	14.4
1999 .....	1.7	*	*	*	*	*	0.7	3.0	5.7	8.4	8.4	10.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of esophagus (C15)</b>												
<b>All origins <sup>3</sup></b>												
Both sexes												
2002 .....	4.4	*	*	*	*	*	0.7	3.5	10.4	20.2	26.6	24.6
2001 .....	4.4	*	*	*	*	0.1	0.7	3.6	10.1	20.6	26.3	25.8
2000 .....	4.3	*	*	*	*	0.1	0.7	3.6	10.8	20.0	25.6	25.9
1999 .....	4.3	*	*	*	*	0.1	0.7	3.5	10.6	20.1	24.4	26.8
Male												
2002 .....	6.9	*	*	*	*	*	1.1	6.0	18.2	35.6	48.3	43.4
2001 .....	6.9	*	*	*	*	0.1	1.1	6.3	17.6	36.0	47.4	48.1
2000 .....	6.7	*	*	*	*	0.2	1.2	6.3	18.6	34.1	45.8	47.8
1999 .....	6.6	*	*	*	*	0.1	1.1	6.1	18.3	35.3	44.4	48.0
Female												
2002 .....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.4	12.2	16.5
2001 .....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.9	12.4	16.5
2000 .....	2.1	*	*	*	*	*	0.1	1.0	3.5	8.3	12.4	17.1
1999 .....	2.0	*	*	*	*	*	0.2	0.9	3.6	7.6	11.5	18.3
<b>Hispanic</b>												
Both sexes												
2002 .....	1.1	*	*	*	*	*	0.4	1.5	4.1	10.2	15.2	23.3
2001 .....	1.1	*	*	*	*	*	*	1.9	5.2	8.4	18.2	20.0
2000 .....	1.1	*	*	*	*	*	*	1.2	5.0	11.1	19.4	19.9
1999 .....	1.1	*	*	*	*	*	*	1.5	5.2	9.2	18.1	26.7
Male												
2002 .....	1.6	*	*	*	*	*	*	2.7	8.0	18.3	27.1	36.8
2001 .....	1.7	*	*	*	*	*	*	3.6	8.5	16.0	30.4	41.9
2000 .....	1.6	*	*	*	*	*	*	2.2	8.0	19.4	35.4	40.3
1999 .....	1.7	*	*	*	*	*	*	2.7	10.2	16.3	32.8	49.6
Female												
2002 .....	0.4	*	*	*	*	*	*	*	*	3.6	7.0	16.6
2001 .....	0.5	*	*	*	*	*	*	*	2.2	*	10.0	*
2000 .....	0.5	*	*	*	*	*	*	*	2.4	4.6	8.6	*
1999 .....	0.4	*	*	*	*	*	*	*	*	3.6	8.3	*
<b>Non-Hispanic <sup>4</sup></b>												
Both sexes												
2002 .....	4.9	*	*	*	*	*	0.7	3.7	10.8	20.8	27.2	24.5
2001 .....	4.9	*	*	*	*	0.1	0.7	3.7	10.4	21.3	26.6	26.0
2000 .....	4.8	*	*	*	*	0.1	0.7	3.8	11.2	20.4	25.8	26.1
1999 .....	4.7	*	*	*	*	0.1	0.7	3.6	11.0	20.6	24.6	26.8
Male												
2002 .....	7.8	*	*	*	*	*	1.2	6.3	18.9	36.6	49.3	43.5
2001 .....	7.7	*	*	*	*	0.1	1.3	6.5	18.2	37.1	48.1	48.4
2000 .....	7.5	*	*	*	*	0.2	1.3	6.6	19.4	34.8	46.1	47.8
1999 .....	7.4	*	*	*	*	0.2	1.2	6.4	18.8	36.2	44.7	48.0
Female												
2002 .....	2.2	*	*	*	*	*	0.3	1.1	3.4	7.6	12.5	16.3
2001 .....	2.2	*	*	*	*	*	0.3	1.1	3.2	8.2	12.5	16.7
2000 .....	2.3	*	*	*	*	*	0.1	1.0	3.6	8.5	12.6	17.3
1999 .....	2.2	*	*	*	*	*	0.2	0.9	3.8	7.8	11.6	18.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of esophagus (C15)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	5.3	*	*	*	*	*	0.7	3.6	10.4	20.5	27.3	25.1
2001 .....	5.1	*	*	*	*	0.1	0.8	3.5	9.7	20.5	26.8	26.5
2000 .....	5.1	*	*	*	*	0.1	0.7	3.5	10.4	20.0	25.5	26.7
1999 .....	4.9	*	*	*	*	0.1	0.7	3.3	10.2	20.0	24.4	27.2
<b>Male</b>												
2002 .....	8.4	*	*	*	*	*	1.3	6.3	18.2	36.3	49.9	44.7
2001 .....	8.1	*	*	*	*	*	1.3	6.1	17.0	36.0	48.6	49.0
2000 .....	8.0	*	*	*	*	0.2	1.4	6.3	18.3	34.3	45.9	49.4
1999 .....	7.7	*	*	*	*	0.2	1.2	6.0	17.5	35.4	44.4	48.0
<b>Female</b>												
2002 .....	2.3	*	*	*	*	*	0.2	0.9	3.0	7.1	12.2	16.6
2001 .....	2.3	*	*	*	*	*	0.2	1.0	2.7	7.5	12.4	17.1
2000 .....	2.3	*	*	*	*	*	*	0.8	3.0	8.0	12.0	17.5
1999 .....	2.2	*	*	*	*	*	0.2	0.8	3.3	7.1	11.3	18.9
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	4.3	*	*	*	*	*	1.0	5.2	17.7	28.9	29.9	20.6
2001 .....	4.7	*	*	*	*	*	1.0	6.1	18.9	32.8	30.0	24.0
2000 .....	4.7	*	*	*	*	*	0.8	6.8	20.2	28.6	34.1	22.6
1999 .....	4.7	*	*	*	*	*	1.0	6.6	20.3	30.2	31.4	23.6
<b>Male</b>												
2002 .....	6.5	*	*	*	*	*	1.5	8.4	30.7	49.8	52.2	38.3
2001 .....	7.2	*	*	*	*	*	1.6	10.5	33.0	56.2	55.5	47.4
2000 .....	7.0	*	*	*	*	*	1.2	11.5	34.8	47.6	58.9	38.7
1999 .....	7.2	*	*	*	*	*	1.7	11.5	35.3	51.1	57.2	49.9
<b>Female</b>												
2002 .....	2.4	*	*	*	*	*	*	2.4	7.2	14.0	17.2	13.8
2001 .....	2.4	*	*	*	*	*	*	2.3	7.5	16.2	15.7	15.4
2000 .....	2.6	*	*	*	*	*	*	2.7	8.5	15.2	20.1	16.6
1999 .....	2.4	*	*	*	*	*	*	2.4	8.3	15.5	17.0	13.8
<b>Malignant neoplasm of stomach (C16)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	4.2	*	*	*	0.1	0.3	1.0	2.8	6.9	15.4	28.7	47.6
2001 .....	4.3	*	*	*	0.1	0.3	1.0	2.9	7.0	16.5	30.2	45.4
2000 .....	4.5	*	*	*	*	0.3	1.1	2.9	7.4	16.6	31.0	52.7
1999 .....	4.6	*	*	*	*	0.3	1.1	3.0	7.8	17.1	31.6	50.1
<b>Male</b>												
2002 .....	5.0	*	*	*	*	0.4	1.1	3.7	9.5	22.5	40.6	67.2
2001 .....	5.1	*	*	*	*	0.4	1.1	3.8	10.2	23.2	43.1	62.6
2000 .....	5.3	*	*	*	*	0.3	1.3	3.8	10.5	23.7	43.9	75.6
1999 .....	5.5	*	*	*	*	0.4	1.3	4.2	11.0	24.2	46.5	71.3
<b>Female</b>												
2002 .....	3.5	*	*	*	0.1	0.3	0.9	1.9	4.5	9.5	20.8	39.1
2001 .....	3.5	*	*	*	*	0.3	0.9	2.0	4.1	10.9	21.7	38.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
All origins <sup>3</sup>												
Female												
2000 .....	3.7	*	*	*	*	0.3	0.9	2.1	4.6	10.8	22.6	43.4
1999 .....	3.7	*	*	*	*	0.3	1.0	1.9	4.9	11.3	21.9	41.6
Hispanic												
Both sexes												
2002 .....	3.1	*	*	*	*	0.5	1.7	4.2	11.0	21.9	47.2	72.1
2001 .....	3.2	*	*	*	*	0.6	1.8	5.4	11.3	26.4	45.6	63.8
2000 .....	3.1	*	*	*	*	0.4	2.0	4.8	9.8	28.4	50.2	65.7
1999 .....	3.1	*	*	*	*	0.6	1.7	4.1	11.7	26.2	49.8	73.7
Male												
2002 .....	3.4	*	*	*	*	0.7	1.5	4.7	13.7	32.9	61.6	107.1
2001 .....	3.6	*	*	*	*	*	1.9	6.2	15.0	37.1	60.4	95.2
2000 .....	3.6	*	*	*	*	*	2.0	5.0	13.7	43.3	70.8	98.8
1999 .....	3.5	*	*	*	*	0.7	1.6	5.2	16.4	35.6	73.5	75.5
Female												
2002 .....	2.7	*	*	*	*	*	1.9	3.8	8.7	13.2	37.4	54.7
2001 .....	2.8	*	*	*	*	0.7	1.7	4.6	8.1	18.0	35.7	48.4
2000 .....	2.7	*	*	*	*	*	2.1	4.6	6.3	16.8	36.3	49.5
1999 .....	2.7	*	*	*	*	*	1.8	3.0	7.6	18.8	34.0	72.8
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	4.4	*	*	*	0.1	0.3	0.9	2.6	6.5	14.9	27.7	46.3
2001 .....	4.5	*	*	*	*	0.2	0.9	2.6	6.6	15.8	29.4	44.7
2000 .....	4.7	*	*	*	*	0.3	0.9	2.7	7.2	15.8	30.1	52.0
1999 .....	4.7	*	*	*	*	0.3	1.0	2.9	7.5	16.5	30.8	49.2
Male												
2002 .....	5.3	*	*	*	*	0.3	1.0	3.6	9.1	21.7	39.4	65.0
2001 .....	5.4	*	*	*	*	0.3	1.0	3.6	9.8	22.3	42.1	61.0
2000 .....	5.5	*	*	*	*	0.3	1.2	3.7	10.3	22.4	42.6	74.5
1999 .....	5.7	*	*	*	*	0.3	1.2	4.0	10.6	23.4	45.3	71.0
Female												
2002 .....	3.5	*	*	*	*	0.3	0.7	1.7	4.1	9.2	19.9	38.3
2001 .....	3.6	*	*	*	*	0.2	0.8	1.7	3.7	10.4	21.0	37.9
2000 .....	3.8	*	*	*	*	0.3	0.7	1.8	4.4	10.4	21.9	42.9
1999 .....	3.8	*	*	*	*	0.3	0.9	1.8	4.7	10.8	21.3	40.5
Non-Hispanic white												
Both sexes												
2002 .....	4.1	*	*	*	*	0.2	0.7	2.1	5.4	12.6	24.1	40.7
2001 .....	4.2	*	*	*	*	0.2	0.7	2.1	5.6	13.5	25.8	40.0
2000 .....	4.4	*	*	*	*	0.2	0.7	2.2	6.1	13.5	26.1	46.3
1999 .....	4.4	*	*	*	*	0.2	0.7	2.3	6.1	14.4	27.0	43.5
Male												
2002 .....	5.0	*	*	*	*	0.2	0.8	2.9	7.8	18.6	35.0	58.0
2001 .....	5.0	*	*	*	*	0.2	0.8	2.8	8.2	18.8	37.4	55.7
2000 .....	5.2	*	*	*	*	0.2	0.8	3.0	8.9	19.3	37.4	67.3
1999 .....	5.4	*	*	*	*	0.2	0.9	3.3	8.7	20.6	39.9	62.6

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of stomach (C16)</b>												
Non-Hispanic white												
Female												
2002 .....	3.2	*	*	*	*	0.2	0.5	1.2	3.2	7.5	16.8	33.3
2001 .....	3.4	*	*	*	*	*	0.6	1.3	3.1	9.0	18.1	33.5
2000 .....	3.5	*	*	*	*	0.2	0.5	1.4	3.6	8.6	18.7	37.9
1999 .....	3.5	*	*	*	*	0.2	0.6	1.4	3.7	9.1	18.6	35.9
Non-Hispanic black												
Both sexes												
2002 .....	5.8	*	*	*	*	0.6	1.6	5.5	13.2	31.5	57.7	100.1
2001 .....	5.8	*	*	*	*	0.5	2.2	5.5	13.1	31.0	59.1	89.3
2000 .....	6.1	*	*	*	*	0.7	2.1	5.8	14.1	32.3	64.3	102.7
1999 .....	6.3	*	*	*	*	0.7	2.3	6.0	16.1	31.5	65.1	103.1
Male												
2002 .....	6.9	*	*	*	*	0.8	1.9	7.9	19.5	47.3	80.9	136.7
2001 .....	6.9	*	*	*	*	*	2.4	8.0	20.0	49.2	87.9	106.3
2000 .....	7.2	*	*	*	*	0.8	3.0	8.1	20.9	47.0	91.0	138.6
1999 .....	7.6	*	*	*	*	*	2.8	8.7	23.3	46.4	101.6	149.8
Female												
2002 .....	4.9	*	*	*	*	*	1.4	3.4	8.1	20.3	44.6	86.1
2001 .....	4.7	*	*	*	*	*	2.0	3.3	7.6	18.1	43.0	83.0
2000 .....	5.2	*	*	*	*	*	1.4	3.9	8.6	22.0	49.3	89.4
1999 .....	5.1	*	*	*	*	*	1.9	3.7	10.2	21.0	44.6	85.7
<b>Malignant neoplasms of colon, rectum and anus (C18-C21)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	19.7	*	*	*	0.1	0.7	3.2	11.4	31.7	73.3	137.4	241.4
2001 .....	20.0	*	*	*	0.1	0.8	3.2	11.6	31.5	75.3	139.4	255.7
2000 .....	20.4	*	*	*	0.1	0.7	3.0	11.2	32.7	76.7	146.0	270.4
1999 .....	20.5	*	*	*	0.1	0.7	3.0	11.1	33.5	77.4	146.7	271.7
Male												
2002 .....	20.1	*	*	*	0.2	0.8	3.6	13.0	39.4	91.3	167.4	275.1
2001 .....	20.2	*	*	*	0.1	0.8	3.5	13.1	38.3	93.7	169.5	298.3
2000 .....	20.6	*	*	*	0.1	0.8	3.4	12.8	39.6	95.4	179.5	315.6
1999 .....	20.6	*	*	*	0.1	0.8	3.0	12.5	40.5	95.1	181.9	324.0
Female												
2002 .....	19.2	*	*	*	*	0.7	2.8	9.8	24.5	58.3	117.5	226.8
2001 .....	19.8	*	*	*	0.1	0.7	2.9	10.2	25.3	60.1	119.6	237.8
2000 .....	20.2	*	*	*	*	0.6	2.7	9.8	26.4	61.2	124.2	251.9
1999 .....	20.3	*	*	*	0.1	0.6	2.9	9.7	27.0	62.9	123.8	250.6
Hispanic												
Both sexes												
2002 .....	6.1	*	*	*	*	0.4	2.1	7.7	23.0	51.2	95.6	165.7
2001 .....	6.1	*	*	*	*	0.3	2.3	7.2	22.5	52.1	96.5	193.8
2000 .....	6.0	*	*	*	*	0.5	2.3	8.4	23.1	49.2	98.2	181.1
1999 .....	6.0	*	*	*	*	0.5	1.7	8.3	21.0	56.1	101.0	176.8
Male												
2002 .....	6.4	*	*	*	*	*	2.0	9.3	31.1	67.1	121.1	174.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms of colon, rectum and anus (C18-C21)</b>												
<b>Hispanic</b>												
<b>Male</b>												
2001 .....	6.2	*	*	*	*	*	2.7	7.4	28.2	66.5	116.7	257.1
2000 .....	6.2	*	*	*	*	*	2.0	9.0	28.2	63.5	131.3	235.8
1999 .....	6.1	*	*	*	*	*	1.5	9.1	24.1	72.7	134.0	211.4
<b>Female</b>												
2002 .....	5.7	*	*	*	*	*	2.1	6.1	15.7	38.6	78.1	161.6
2001 .....	5.9	*	*	*	*	*	1.9	7.1	17.4	40.8	82.8	162.9
2000 .....	5.8	*	*	*	*	*	2.7	7.8	18.5	38.0	75.9	154.3
1999 .....	5.9	*	*	*	*	*	2.0	7.5	18.3	43.0	79.1	160.1
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	21.7	*	*	*	0.1	0.8	3.3	11.8	32.3	74.6	139.1	244.0
2001 .....	22.0	*	*	*	0.1	0.9	3.3	12.0	32.2	76.7	141.1	257.4
2000 .....	22.4	*	*	*	0.1	0.7	3.1	11.5	33.3	78.2	147.8	273.1
1999 .....	22.4	*	*	*	0.1	0.7	3.1	11.3	34.3	78.5	148.1	274.3
<b>Male</b>												
2002 .....	22.3	*	*	*	0.2	0.9	3.8	13.4	39.9	92.7	169.3	279.2
2001 .....	22.3	*	*	*	0.1	0.9	3.6	13.6	38.9	95.2	171.8	299.6
2000 .....	22.8	*	*	*	0.1	0.8	3.6	13.1	40.3	97.2	181.3	318.6
1999 .....	22.7	*	*	*	0.1	0.9	3.2	12.8	41.6	96.2	183.2	327.5
<b>Female</b>												
2002 .....	21.2	*	*	*	*	0.7	2.9	10.2	25.2	59.6	119.1	228.8
2001 .....	21.7	*	*	*	*	0.8	3.0	10.5	25.9	61.3	121.0	239.9
2000 .....	22.1	*	*	*	*	0.6	2.7	9.9	26.9	62.6	126.1	254.7
1999 .....	22.2	*	*	*	*	0.6	3.0	9.9	27.6	63.9	125.3	253.0
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	23.2	*	*	*	0.1	0.8	3.0	10.9	30.6	73.0	136.8	243.2
2001 .....	23.5	*	*	*	0.1	0.8	3.1	11.1	30.6	74.7	138.5	256.9
2000 .....	24.0	*	*	*	0.1	0.6	2.8	10.6	31.7	76.5	145.7	271.9
1999 .....	24.0	*	*	*	0.1	0.7	2.9	10.6	32.8	76.7	145.9	273.4
<b>Male</b>												
2002 .....	23.8	*	*	*	0.2	0.9	3.4	12.5	38.0	91.1	165.8	277.6
2001 .....	23.9	*	*	*	*	0.9	3.3	12.8	37.4	93.2	169.3	298.2
2000 .....	24.2	*	*	*	*	0.8	3.2	12.2	38.2	95.1	178.4	315.8
1999 .....	24.3	*	*	*	*	0.8	3.1	12.1	39.7	94.8	180.6	328.2
<b>Female</b>												
2002 .....	22.7	*	*	*	*	0.7	2.7	9.4	23.7	57.6	117.3	228.3
2001 .....	23.1	*	*	*	*	0.7	2.8	9.5	24.2	59.0	118.1	239.8
2000 .....	23.8	*	*	*	*	0.5	2.4	9.0	25.5	60.9	124.1	254.1
1999 .....	23.7	*	*	*	*	0.6	2.7	9.0	26.2	61.5	123.2	251.5
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	18.9	*	*	*	*	0.7	5.1	19.0	50.5	100.6	185.9	278.3
2001 .....	19.2	*	*	*	*	1.1	5.0	19.5	50.6	105.9	189.7	289.0
2000 .....	19.4	*	*	*	*	1.0	5.2	20.0	51.1	106.3	192.5	312.9
1999 .....	19.3	*	*	*	*	1.1	4.9	18.1	53.7	107.2	194.5	308.8

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms of colon, rectum and anus (C18-C21)</b>												
<b>Non-Hispanic black</b>												
Male												
2002 .....	19.3	*	*	*	*	0.9	6.0	21.8	62.7	123.0	240.9	332.6
2001 .....	19.0	*	*	*	*	1.2	5.8	21.4	59.3	130.7	229.9	352.3
2000 .....	19.8	*	*	*	*	1.1	6.1	22.7	63.7	133.7	244.4	381.6
1999 .....	19.2	*	*	*	*	1.2	4.8	20.6	67.0	127.8	245.8	351.8
Female												
2002 .....	18.4	*	*	*	*	*	4.3	16.7	40.7	84.7	154.9	257.5
2001 .....	19.4	*	*	*	*	1.0	4.3	17.9	43.6	88.4	167.1	265.6
2000 .....	19.1	*	*	*	*	1.0	4.4	17.6	40.9	87.1	163.2	287.4
1999 .....	19.5	*	*	*	*	0.9	5.0	15.9	43.1	92.7	165.8	292.7
<b>Malignant neoplasms of liver and intrahepatic bile ducts (C22)</b>												
<b>All origins <sup>3</sup></b>												
Both sexes												
2002 .....	4.9	*	0.2	*	0.1	0.3	0.9	4.9	9.6	19.6	29.3	35.0
2001 .....	4.7	*	*	*	0.1	0.2	0.9	4.8	9.3	18.8	29.0	34.1
2000 .....	4.6	*	0.1	0.0	0.1	0.2	1.0	4.5	9.3	18.7	28.8	32.7
1999 .....	4.4	*	0.2	*	0.1	0.2	1.0	4.1	8.8	19.0	28.0	30.6
Male												
2002 .....	6.4	*	*	*	*	0.3	1.4	8.2	14.5	28.5	41.6	46.2
2001 .....	6.1	*	*	*	*	0.2	1.3	7.7	13.9	26.8	41.0	50.3
2000 .....	5.9	*	*	*	0.1	0.3	1.3	7.2	13.8	27.0	42.1	44.7
1999 .....	5.6	*	0.3	*	0.1	0.2	1.4	6.5	12.9	27.5	39.0	40.1
Female												
2002 .....	3.4	*	*	*	*	0.2	0.5	1.8	5.1	12.2	21.1	30.2
2001 .....	3.4	*	*	*	*	0.2	0.4	1.9	5.0	12.1	21.1	27.2
2000 .....	3.3	*	*	*	*	0.1	0.6	1.9	5.1	11.8	20.1	27.8
1999 .....	3.3	*	*	*	*	0.2	0.6	1.8	5.1	11.9	20.9	26.8
<b>Hispanic</b>												
Both sexes												
2002 .....	3.5	*	*	*	*	*	0.8	5.9	15.7	34.1	44.2	59.9
2001 .....	3.4	*	*	*	*	*	0.8	6.7	14.1	28.8	52.1	52.5
2000 .....	3.3	*	*	*	*	*	1.2	5.6	13.7	31.1	51.8	58.4
1999 .....	3.2	*	*	*	*	*	0.9	5.7	13.1	31.3	49.8	53.3
Male												
2002 .....	4.5	*	*	*	*	*	1.1	9.8	24.1	48.9	59.9	75.3
2001 .....	4.3	*	*	*	*	*	1.3	11.1	21.5	41.1	68.0	64.8
2000 .....	4.2	*	*	*	*	*	1.7	9.1	20.5	43.9	74.7	72.6
1999 .....	3.9	*	*	*	*	*	1.1	9.6	20.7	42.8	63.1	66.9
Female												
2002 .....	2.5	*	*	*	*	*	*	2.1	8.2	22.4	33.4	52.2
2001 .....	2.5	*	*	*	*	*	*	2.3	7.6	19.1	41.4	46.5
2000 .....	2.4	*	*	*	*	*	*	2.2	7.6	21.1	36.3	51.4
1999 .....	2.4	*	*	*	*	*	*	1.9	6.3	22.2	40.9	46.8
<b>Non-Hispanic <sup>4</sup></b>												
Both sexes												
2002 .....	5.1	*	0.2	*	0.1	0.3	0.9	4.8	9.1	18.6	28.4	33.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms of liver and intrahepatic bile ducts (C22)</b>												
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2001 .....	4.9	*	*	*	0.1	0.2	0.9	4.6	8.9	18.1	27.9	33.2
2000 .....	4.8	*	0.2	*	0.1	0.2	0.9	4.3	8.9	17.9	27.7	31.7
1999 .....	4.6	*	0.2	*	0.1	0.2	1.0	3.9	8.4	18.2	27.1	29.7
<b>Male</b>												
2002 .....	6.6	*	*	*	*	0.4	1.4	8.0	13.7	27.1	40.5	44.8
2001 .....	6.3	*	*	*	*	0.2	1.3	7.4	13.2	25.8	39.7	49.7
2000 .....	6.2	*	*	*	*	0.3	1.3	6.9	13.3	25.9	40.5	43.4
1999 .....	5.8	*	*	*	0.1	0.3	1.5	6.2	12.2	26.5	37.9	38.7
<b>Female</b>												
2002 .....	3.6	*	*	*	*	0.2	0.5	1.7	4.8	11.4	20.5	29.2
2001 .....	3.5	*	*	*	*	0.2	0.4	1.9	4.8	11.7	20.2	26.4
2000 .....	3.4	*	*	*	*	0.1	0.6	1.9	4.9	11.2	19.4	27.0
1999 .....	3.4	*	*	*	*	0.2	0.5	1.8	4.9	11.3	20.0	26.1
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	5.0	*	*	*	*	0.2	0.7	3.9	7.7	17.0	27.4	32.6
2001 .....	4.8	*	*	*	0.1	0.1	0.6	3.7	7.6	16.6	26.5	32.4
2000 .....	4.6	*	*	*	*	0.1	0.6	3.4	7.5	16.2	26.5	30.8
1999 .....	4.5	*	0.2	*	0.1	0.1	0.7	3.1	7.1	16.8	26.1	28.9
<b>Male</b>												
2002 .....	6.4	*	*	*	*	0.2	1.1	6.4	11.5	24.6	38.7	43.6
2001 .....	6.1	*	*	*	*	*	0.9	5.9	11.2	24.2	38.0	48.9
2000 .....	5.9	*	*	*	*	0.2	0.8	5.3	11.0	23.6	39.2	41.6
1999 .....	5.7	*	*	*	*	*	1.1	4.8	10.2	24.6	36.7	39.0
<b>Female</b>												
2002 .....	3.6	*	*	*	*	0.2	0.4	1.4	4.2	10.4	19.9	27.8
2001 .....	3.4	*	*	*	*	*	0.3	1.5	4.1	10.2	18.9	25.5
2000 .....	3.4	*	*	*	*	*	0.4	1.6	4.3	10.1	18.2	26.5
1999 .....	3.4	*	*	*	*	*	0.4	1.6	4.2	10.2	19.1	24.8
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	4.7	*	*	*	*	0.5	1.5	8.9	15.0	23.3	30.9	38.1
2001 .....	4.6	*	*	*	*	*	1.7	8.9	15.1	21.9	32.3	35.3
2000 .....	4.5	*	*	*	*	0.4	1.8	8.8	14.5	23.7	29.1	31.8
1999 .....	4.3	*	*	*	*	0.4	1.7	7.4	13.4	24.6	29.7	34.9
<b>Male</b>												
2002 .....	6.6	*	*	*	*	*	2.3	15.4	24.4	38.2	46.3	41.6
2001 .....	6.1	*	*	*	*	*	2.7	15.1	24.2	30.3	42.6	42.7
2000 .....	6.3	*	*	*	*	*	2.5	15.0	24.1	37.3	40.4	48.1
1999 .....	5.7	*	*	*	*	*	2.5	13.1	20.3	36.8	41.6	30.9
<b>Female</b>												
2002 .....	3.0	*	*	*	*	*	0.8	3.3	7.4	12.8	22.2	36.8
2001 .....	3.3	*	*	*	*	*	0.8	3.6	7.8	16.0	26.4	32.5
2000 .....	2.9	*	*	*	*	*	1.2	3.4	6.7	14.1	22.8	25.7
1999 .....	3.1	*	*	*	*	*	0.9	2.5	7.9	15.9	23.1	36.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of pancreas (C25)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.8	75.0	94.5
2001 .....	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.5	74.1	100.1
2000 .....	10.4	*	*	*	*	0.1	1.2	6.3	19.4	44.1	76.5	96.4
1999 .....	10.4	*	*	*	*	0.1	1.3	6.4	19.8	45.0	73.1	99.7
<b>Male</b>												
2002 .....	10.5	*	*	*	*	0.1	1.6	7.8	23.6	51.4	84.0	100.9
2001 .....	10.3	*	*	*	*	0.1	1.4	7.6	23.6	50.8	84.0	105.1
2000 .....	10.3	*	*	*	*	0.2	1.4	8.0	23.6	50.4	85.9	102.9
1999 .....	10.4	*	*	*	*	0.2	1.6	8.3	23.9	52.4	81.8	107.5
<b>Female</b>												
2002 .....	10.5	*	*	*	*	*	1.0	4.4	15.1	39.3	69.1	91.7
2001 .....	10.6	*	*	*	*	0.1	1.1	4.7	15.2	39.4	67.6	98.1
2000 .....	10.5	*	*	*	*	0.1	1.0	4.6	15.5	38.8	70.4	93.8
1999 .....	10.5	*	*	*	*	0.1	1.0	4.7	16.0	39.0	67.4	96.6
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	3.5	*	*	*	*	*	0.9	4.0	13.8	34.7	55.5	79.3
2001 .....	3.6	*	*	*	*	*	0.8	3.7	14.2	36.9	61.3	83.8
2000 .....	3.5	*	*	*	*	*	0.7	4.1	13.7	36.9	61.8	82.9
1999 .....	3.3	*	*	*	*	*	0.9	4.6	13.3	32.6	56.7	84.2
<b>Male</b>												
2002 .....	3.3	*	*	*	*	*	1.2	5.3	15.7	40.9	55.8	60.2
2001 .....	3.4	*	*	*	*	*	0.9	4.3	16.2	42.3	68.9	62.9
2000 .....	3.4	*	*	*	*	*	0.8	5.2	16.6	45.2	62.9	80.6
1999 .....	3.2	*	*	*	*	*	1.2	5.7	15.2	40.0	56.3	86.3
<b>Female</b>												
2002 .....	3.7	*	*	*	*	*	*	2.8	12.1	29.8	55.4	88.7
2001 .....	3.9	*	*	*	*	*	0.8	3.2	12.5	32.6	56.2	94.0
2000 .....	3.6	*	*	*	*	*	*	3.0	11.1	30.4	61.1	84.1
1999 .....	3.5	*	*	*	*	*	*	3.5	11.6	26.8	56.9	83.2
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	11.6	*	*	*	*	0.1	1.4	6.3	19.6	45.4	75.8	94.9
2001 .....	11.5	*	*	*	*	0.1	1.3	6.4	19.6	44.9	74.6	100.6
2000 .....	11.4	*	*	*	*	0.1	1.3	6.5	19.8	44.4	77.0	96.6
1999 .....	11.4	*	*	*	*	0.1	1.4	6.6	20.2	45.6	73.5	100.1
<b>Male</b>												
2002 .....	11.7	*	*	*	*	0.1	1.7	8.0	24.1	52.1	85.3	102.4
2001 .....	11.4	*	*	*	*	0.2	1.5	7.9	24.2	51.2	84.6	106.4
2000 .....	11.3	*	*	*	*	0.2	1.5	8.3	24.1	50.7	86.7	103.5
1999 .....	11.4	*	*	*	*	0.2	1.7	8.4	24.4	52.9	82.7	108.2
<b>Female</b>												
2002 .....	11.5	*	*	*	*	*	1.1	4.6	15.4	39.8	69.5	91.7
2001 .....	11.5	*	*	*	*	0.1	1.2	4.9	15.4	39.8	68.0	98.2
2000 .....	11.4	*	*	*	*	*	1.0	4.7	15.8	39.2	70.6	93.8
1999 .....	11.4	*	*	*	*	*	1.0	4.8	16.4	39.6	67.6	96.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of pancreas (C25)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	12.4	*	*	*	*	0.1	1.4	6.0	19.0	44.3	75.1	94.6
2001 .....	12.3	*	*	*	*	0.1	1.3	6.1	19.3	44.1	73.7	99.4
2000 .....	12.2	*	*	*	*	0.2	1.2	6.3	19.2	43.4	75.7	94.7
1999 .....	12.1	*	*	*	*	0.1	1.3	6.3	19.5	44.7	72.1	98.9
<b>Male</b>												
2002 .....	12.6	*	*	*	*	*	1.7	7.7	23.5	51.2	85.4	102.9
2001 .....	12.4	*	*	*	*	0.2	1.5	7.7	24.0	50.3	84.5	108.0
2000 .....	12.2	*	*	*	*	0.2	1.5	8.0	23.4	50.0	86.1	103.0
1999 .....	12.1	*	*	*	*	0.2	1.6	8.0	23.4	52.1	81.6	108.1
<b>Female</b>												
2002 .....	12.2	*	*	*	*	*	1.1	4.3	14.7	38.3	68.1	91.0
2001 .....	12.3	*	*	*	*	*	1.2	4.6	14.9	38.8	66.6	95.8
2000 .....	12.2	*	*	*	*	*	1.0	4.5	15.3	37.9	68.9	91.4
1999 .....	12.1	*	*	*	*	*	1.0	4.7	15.7	38.5	65.9	95.2
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	9.6	*	*	*	*	*	1.8	8.9	28.3	61.7	90.9	108.3
2001 .....	9.4	*	*	*	*	*	1.6	9.2	25.9	59.0	93.3	122.0
2000 .....	9.7	*	*	*	*	*	1.7	9.2	27.9	60.0	98.9	121.2
1999 .....	9.7	*	*	*	*	*	2.0	9.9	28.8	60.4	95.0	119.0
<b>Male</b>												
2002 .....	9.3	*	*	*	*	*	2.2	11.6	34.9	69.5	97.3	102.8
2001 .....	8.9	*	*	*	*	*	1.8	11.4	30.8	68.2	100.5	100.5
2000 .....	9.3	*	*	*	*	*	2.1	12.1	34.6	66.9	108.0	110.4
1999 .....	9.7	*	*	*	*	*	2.4	13.6	37.4	68.7	107.6	115.3
<b>Female</b>												
2002 .....	9.8	*	*	*	*	*	1.5	6.5	23.0	56.2	87.2	110.4
2001 .....	9.9	*	*	*	*	*	1.4	7.3	21.9	52.4	89.3	130.0
2000 .....	10.0	*	*	*	*	*	1.5	6.8	22.5	55.1	93.8	125.2
1999 .....	9.7	*	*	*	*	*	1.7	6.7	21.9	54.6	87.9	120.4
<b>Malignant neoplasm of larynx (C32)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	1.3	*	*	*	*	*	0.2	1.1	3.4	5.9	7.6	5.5
2001 .....	1.3	*	*	*	*	*	0.2	1.1	3.6	6.4	7.2	6.7
2000 .....	1.4	*	*	*	*	*	0.1	1.1	3.6	6.6	8.1	6.5
1999 .....	1.4	*	*	*	*	*	0.2	1.1	3.8	6.7	7.3	7.0
<b>Male</b>												
2002 .....	2.1	*	*	*	*	*	0.3	1.8	5.8	10.0	14.9	13.2
2001 .....	2.2	*	*	*	*	*	0.3	1.8	6.1	11.1	14.2	18.3
2000 .....	2.2	*	*	*	*	*	0.2	1.8	6.2	11.4	15.7	17.4
1999 .....	2.2	*	*	*	*	*	0.3	1.9	6.4	11.4	14.1	17.8
<b>Female</b>												
2002 .....	0.5	*	*	*	*	*	*	0.4	1.2	2.5	2.8	2.2
2001 .....	0.5	*	*	*	*	*	0.1	0.4	1.3	2.5	2.6	1.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of larynx (C32)</b>												
<b>All origins <sup>3</sup></b>												
Female												
2002 .....	0.6	*	*	*	*	*	*	0.4	1.2	2.7	3.1	2.1
1999 .....	0.6	*	*	*	*	*	*	0.3	1.4	2.9	2.8	2.6
<b>Hispanic</b>												
Both sexes												
2002 .....	0.4	*	*	*	*	*	*	*	2.2	4.1	5.5	*
2001 .....	0.4	*	*	*	*	*	*	0.6	2.1	5.3	5.2	*
2000 .....	0.4	*	*	*	*	*	*	*	1.7	4.2	6.9	*
1999 .....	0.5	*	*	*	*	*	*	0.8	1.8	5.3	8.3	*
Male												
2002 .....	0.7	*	*	*	*	*	*	*	4.3	8.0	11.1	*
2001 .....	0.7	*	*	*	*	*	*	*	3.9	10.6	10.7	*
2000 .....	0.7	*	*	*	*	*	*	*	3.5	7.8	14.8	*
1999 .....	0.9	*	*	*	*	*	*	1.6	3.5	10.5	16.2	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
Both sexes												
2002 .....	1.4	*	*	*	*	*	0.2	1.1	3.5	6.0	7.7	5.4
2001 .....	1.5	*	*	*	*	*	0.2	1.1	3.7	6.4	7.3	6.7
2000 .....	1.5	*	*	*	*	*	0.1	1.2	3.7	6.7	8.1	6.5
1999 .....	1.5	*	*	*	*	*	0.2	1.1	3.9	6.8	7.2	6.8
Male												
2002 .....	2.3	*	*	*	*	*	0.3	1.9	5.9	10.1	15.0	13.2
2001 .....	2.4	*	*	*	*	*	0.3	1.9	6.1	11.1	14.3	18.4
2000 .....	2.4	*	*	*	*	*	0.2	1.9	6.4	11.5	15.7	17.3
1999 .....	2.4	*	*	*	*	*	0.3	1.9	6.6	11.4	14.0	17.3
Female												
2002 .....	0.6	*	*	*	*	*	*	0.4	1.3	2.6	2.9	2.1
2001 .....	0.6	*	*	*	*	*	0.1	0.4	1.4	2.6	2.7	1.8
2000 .....	0.6	*	*	*	*	*	*	0.5	1.2	2.8	3.1	2.1
1999 .....	0.6	*	*	*	*	*	*	0.3	1.4	3.0	2.8	2.6
<b>Non-Hispanic white</b>												
Both sexes												
2002 .....	1.4	*	*	*	*	*	0.1	0.9	3.1	5.8	7.3	5.5
2001 .....	1.5	*	*	*	*	*	0.2	0.9	3.3	6.0	6.9	6.5
2000 .....	1.5	*	*	*	*	*	0.1	1.0	3.2	6.2	7.7	6.5
1999 .....	1.5	*	*	*	*	*	0.1	0.9	3.5	6.4	7.1	6.5
Male												
2002 .....	2.3	*	*	*	*	*	0.2	1.5	5.1	9.5	14.2	13.3
2001 .....	2.3	*	*	*	*	*	0.3	1.5	5.5	10.2	13.3	18.3
2000 .....	2.4	*	*	*	*	*	0.2	1.5	5.4	10.5	15.0	17.4
1999 .....	2.3	*	*	*	*	*	0.2	1.6	5.8	10.7	13.6	16.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of larynx (C32)</b>												
<b>Non-Hispanic white</b>												
<b>Female</b>												
2002 .....	0.6	*	*	*	*	*	*	0.3	1.3	2.6	2.7	2.2
2001 .....	0.6	*	*	*	*	*	*	0.3	1.2	2.4	2.7	1.6
2000 .....	0.6	*	*	*	*	*	*	0.4	1.1	2.6	3.0	2.1
1999 .....	0.7	*	*	*	*	*	*	0.3	1.3	2.9	2.9	2.7
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	1.9	*	*	*	*	*	0.5	2.7	7.7	10.6	13.9	*
2001 .....	1.9	*	*	*	*	*	0.4	2.9	7.9	11.5	12.0	8.4
2000 .....	2.0	*	*	*	*	*	*	2.9	9.2	12.9	14.3	7.0
1999 .....	1.9	*	*	*	*	*	0.6	3.0	8.7	11.7	10.0	8.4
<b>Male</b>												
2002 .....	3.1	*	*	*	*	*	0.8	4.6	15.1	19.8	29.6	*
2001 .....	3.1	*	*	*	*	*	*	5.2	14.2	21.4	28.2	*
2000 .....	3.4	*	*	*	*	*	*	5.2	17.5	24.6	29.0	*
1999 .....	3.2	*	*	*	*	*	1.0	5.3	16.3	22.1	22.5	25.0
<b>Female</b>												
2002 .....	0.7	*	*	*	*	*	*	1.1	1.8	4.1	5.0	*
2001 .....	0.7	*	*	*	*	*	*	*	2.9	4.6	*	*
2000 .....	0.8	*	*	*	*	*	*	1.0	2.4	4.7	6.1	*
1999 .....	0.7	*	*	*	*	*	*	0.9	2.7	4.3	*	*
<b>Malignant neoplasms of trachea, bronchus and lung (C33-C34)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	54.7	*	*	*	0.1	0.4	6.0	30.3	115.3	275.0	377.6	297.2
2001 .....	54.8	*	*	*	0.1	0.4	6.2	30.7	117.7	279.7	371.4	302.7
2000 .....	55.3	*	*	*	0.1	0.5	6.1	31.6	122.4	284.2	370.8	302.1
1999 .....	54.5	*	*	*	0.1	0.4	6.0	31.3	123.4	281.5	362.1	297.0
<b>Male</b>												
2002 .....	63.7	*	*	*	*	0.4	6.2	36.6	144.0	355.9	527.9	482.2
2001 .....	64.7	*	*	*	*	0.4	6.7	37.4	147.1	370.0	529.5	512.4
2000 .....	65.5	*	*	*	*	0.5	6.9	38.5	154.0	377.9	532.2	521.2
1999 .....	65.4	*	*	*	*	0.5	6.6	38.2	158.0	377.2	527.6	529.2
<b>Female</b>												
2002 .....	46.0	*	*	*	*	0.4	5.8	24.3	88.8	207.7	277.8	217.0
2001 .....	45.3	*	*	*	*	0.4	5.7	24.3	90.5	204.9	267.5	215.0
2000 .....	45.4	*	*	*	*	0.5	5.3	25.0	93.3	206.9	265.6	212.8
1999 .....	44.1	*	*	*	*	0.4	5.5	24.6	91.5	203.1	254.5	203.5
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	10.4	*	*	*	*	0.4	1.5	9.9	40.9	110.4	183.7	189.0
2001 .....	10.3	*	*	*	*	*	1.9	10.7	41.0	108.3	182.3	200.7
2000 .....	10.3	*	*	*	*	*	1.8	10.9	39.3	110.6	198.7	213.7
1999 .....	10.4	*	*	*	*	*	2.0	10.4	44.2	112.9	187.7	227.3
<b>Male</b>												
2002 .....	13.1	*	*	*	*	*	1.5	12.6	60.7	161.7	299.1	307.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Hispanic												
Male												
2001 .....	13.1	*	*	*	*	*	2.4	14.2	59.5	163.7	293.8	335.2
2000 .....	13.3	*	*	*	*	*	2.1	14.8	58.6	167.3	327.5	368.8
1999 .....	13.6	*	*	*	*	*	2.3	14.2	68.1	175.3	316.0	362.3
Female												
2002 .....	7.6	*	*	*	*	*	1.6	7.3	23.2	69.5	104.6	130.1
2001 .....	7.4	*	*	*	*	*	1.4	7.3	24.6	64.6	107.0	135.0
2000 .....	7.2	*	*	*	*	*	1.4	7.1	22.2	66.0	112.3	137.5
1999 .....	7.1	*	*	*	*	*	1.7	6.7	23.1	64.0	102.3	162.2
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	61.4	*	*	*	*	0.4	6.7	32.3	121.0	285.7	386.2	301.0
2001 .....	61.3	*	*	*	*	0.4	6.8	32.5	123.3	290.4	379.4	305.8
2000 .....	61.5	*	*	*	*	0.5	6.6	33.4	128.4	294.2	377.4	304.5
1999 .....	60.5	*	*	*	*	0.5	6.5	33.0	128.9	290.7	368.2	298.8
Male												
2002 .....	71.8	*	*	*	*	0.5	6.9	38.9	150.3	368.1	538.0	488.9
2001 .....	72.6	*	*	*	*	0.4	7.3	39.5	153.2	382.4	539.5	518.4
2000 .....	73.2	*	*	*	*	0.6	7.5	40.5	160.6	389.7	539.9	526.2
1999 .....	72.7	*	*	*	*	0.5	7.1	40.1	164.0	387.8	534.9	534.6
Female												
2002 .....	51.6	*	*	*	*	0.4	6.4	25.9	93.9	216.9	285.5	220.0
2001 .....	50.5	*	*	*	*	0.5	6.2	25.8	95.6	213.9	274.3	217.4
2000 .....	50.4	*	*	*	*	0.5	5.8	26.5	98.6	215.4	271.6	214.8
1999 .....	48.8	*	*	*	*	0.4	6.0	26.1	96.5	211.0	260.0	204.4
Non-Hispanic white												
Both sexes												
2002 .....	67.7	*	*	*	*	0.4	6.5	31.2	120.6	292.0	392.2	304.8
2001 .....	67.3	*	*	*	*	0.5	6.6	31.5	124.0	295.2	383.0	308.6
2000 .....	67.4	*	*	*	*	0.5	6.2	32.0	129.1	298.0	381.7	306.8
1999 .....	65.9	*	*	*	*	0.5	6.2	31.5	128.9	293.5	370.3	300.5
Male												
2002 .....	77.8	*	*	*	*	0.5	6.5	36.5	145.7	369.5	538.3	493.3
2001 .....	78.4	*	*	*	*	0.4	6.9	37.0	150.3	381.1	536.5	520.0
2000 .....	78.9	*	*	*	*	0.6	6.7	37.7	157.7	387.3	537.7	527.3
1999 .....	78.1	*	*	*	*	0.5	6.6	37.1	160.1	384.4	531.1	536.9
Female												
2002 .....	57.9	*	*	*	*	0.3	6.5	26.0	96.8	225.9	294.3	223.4
2001 .....	56.6	*	*	*	*	0.5	6.4	26.0	99.2	222.3	281.3	220.6
2000 .....	56.4	*	*	*	*	0.5	5.7	26.4	102.2	222.9	279.2	218.0
1999 .....	54.3	*	*	*	*	0.5	5.8	26.1	99.5	217.5	265.0	206.2
Non-Hispanic black												
Both sexes												
2002 .....	44.7	*	*	*	*	0.4	9.0	48.1	155.6	299.5	385.6	274.9
2001 .....	44.6	*	*	*	*	*	9.2	47.5	149.4	311.4	390.9	281.6
2000 .....	45.4	*	*	*	*	0.6	10.4	51.5	155.0	320.2	379.4	291.0
1999 .....	45.6	*	*	*	*	0.4	9.6	51.5	161.1	317.7	390.9	293.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms of trachea, bronchus and lung (C33-C34)</b>												
<b>Non-Hispanic black</b>												
Male												
2002 .....	58.1	*	*	*	*	*	11.0	66.3	228.0	451.4	633.9	490.1
2001 .....	59.2	*	*	*	*	*	11.4	67.5	218.3	484.4	668.0	550.9
2000 .....	60.3	*	*	*	*	*	13.8	72.4	227.1	495.4	649.5	567.1
1999 .....	60.9	*	*	*	*	*	11.7	73.7	241.9	491.0	665.1	580.0
Female												
2002 .....	32.6	*	*	*	*	*	7.3	32.5	97.5	191.8	245.1	192.7
2001 .....	31.3	*	*	*	*	*	7.3	30.4	94.0	189.4	235.3	181.8
2000 .....	31.9	*	*	*	*	*	7.4	33.6	97.0	196.9	227.3	188.4
1999 .....	31.8	*	*	*	*	*	7.9	32.5	96.4	195.5	237.4	186.1
<b>Malignant melanoma of skin (C43)</b>												
<b>All origins <sup>3</sup></b>												
Both sexes												
2002 .....	2.6	*	*	*	0.1	0.5	1.3	2.8	5.1	8.8	13.7	17.9
2001 .....	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.2	13.3	17.8
2000 .....	2.6	*	*	*	0.1	0.5	1.4	3.2	5.4	8.9	13.1	18.9
1999 .....	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.0	13.4	16.4
Male												
2002 .....	3.3	*	*	*	0.1	0.6	1.6	3.8	7.2	13.0	21.2	29.2
2001 .....	3.4	*	*	*	0.1	0.5	1.7	3.9	7.3	13.9	21.3	29.9
2000 .....	3.3	*	*	*	0.1	0.5	1.6	4.2	7.4	12.9	20.8	31.8
1999 .....	3.3	*	*	*	0.1	0.5	1.8	4.0	7.5	13.2	20.9	27.5
Female												
2002 .....	1.9	*	*	*	0.1	0.4	1.1	1.9	3.2	5.4	8.8	12.9
2001 .....	1.9	*	*	*	*	0.4	1.1	2.3	3.3	5.3	8.1	12.8
2000 .....	2.0	*	*	*	0.1	0.4	1.2	2.2	3.5	5.7	8.2	13.7
1999 .....	1.9	*	*	*	*	0.4	1.0	2.2	3.3	5.5	8.5	11.9
<b>Hispanic</b>												
Both sexes												
2002 .....	0.4	*	*	*	*	*	0.4	0.7	1.2	*	4.8	*
2001 .....	0.4	*	*	*	*	*	0.4	0.7	1.4	2.5	5.0	*
2000 .....	0.4	*	*	*	*	*	0.5	0.9	1.6	2.4	*	*
1999 .....	0.4	*	*	*	*	*	*	0.7	1.8	2.6	4.2	*
Male												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.5	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.5	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
Both sexes												
2002 .....	2.9	*	*	*	0.1	0.6	1.5	3.0	5.4	9.3	14.1	18.2

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant melanoma of skin (C43)</b>												
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2001 .....	3.0	*	*	*	0.1	0.5	1.5	3.3	5.6	9.7	13.7	18.3
2000 .....	3.0	*	*	*	0.1	0.5	1.5	3.4	5.7	9.3	13.5	19.3
1999 .....	2.9	*	*	*	0.1	0.6	1.5	3.3	5.6	9.3	13.7	16.7
<b>Male</b>												
2002 .....	3.8	*	*	*	*	0.7	1.8	4.1	7.6	13.7	22.0	30.2
2001 .....	3.9	*	*	*	0.1	0.6	1.9	4.2	7.8	14.6	22.0	31.0
2000 .....	3.7	*	*	*	0.1	0.6	1.7	4.4	7.8	13.4	21.3	32.6
1999 .....	3.7	*	*	*	0.1	0.6	2.0	4.2	7.8	13.7	21.5	28.2
<b>Female</b>												
2002 .....	2.1	*	*	*	*	0.4	1.2	2.0	3.3	5.7	8.9	12.9
2001 .....	2.1	*	*	*	*	0.5	1.2	2.4	3.5	5.6	8.3	12.9
2000 .....	2.2	*	*	*	*	0.4	1.3	2.4	3.7	5.9	8.4	13.9
1999 .....	2.1	*	*	*	*	0.5	1.1	2.3	3.5	5.7	8.7	12.0
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	3.6	*	*	*	0.1	0.7	1.8	3.6	6.2	10.5	15.6	19.6
2001 .....	3.7	*	*	*	0.2	0.7	1.9	4.0	6.5	10.8	15.0	19.9
2000 .....	3.6	*	*	*	0.2	0.7	1.8	4.0	6.5	10.4	14.7	20.8
1999 .....	3.5	*	*	*	0.1	0.7	1.9	3.9	6.4	10.5	14.9	17.9
<b>Male</b>												
2002 .....	4.7	*	*	*	*	0.9	2.2	4.8	8.6	15.3	24.1	32.9
2001 .....	4.8	*	*	*	0.2	0.8	2.3	5.0	9.0	16.3	24.1	34.1
2000 .....	4.6	*	*	*	0.2	0.8	2.0	5.2	8.9	15.0	23.2	35.7
1999 .....	4.5	*	*	*	*	0.8	2.4	5.0	8.9	15.4	23.3	30.5
<b>Female</b>												
2002 .....	2.6	*	*	*	*	0.6	1.4	2.4	3.9	6.5	9.8	13.8
2001 .....	2.6	*	*	*	*	0.6	1.5	2.9	4.1	6.2	9.0	13.9
2000 .....	2.7	*	*	*	*	0.6	1.6	2.8	4.2	6.6	9.1	14.7
1999 .....	2.5	*	*	*	*	0.6	1.4	2.8	4.1	6.4	9.4	12.9
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	0.3	*	*	*	*	*	*	*	0.8	1.5	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	2.2	2.6	*
2000 .....	0.4	*	*	*	*	*	*	*	*	1.9	3.6	6.7
1999 .....	0.3	*	*	*	*	*	*	*	*	1.4	2.8	*
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	2.1	*	*
2000 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of breast (C50)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	14.5	*	*	*	*	1.1	6.1	16.1	29.5	46.5	76.5	134.6
2001 .....	14.7	*	*	*	*	1.2	6.3	16.8	30.2	47.6	76.8	134.2
2000 .....	15.0	*	*	*	*	1.2	6.3	16.9	31.2	49.1	78.9	147.4
1999 .....	14.9	*	*	*	*	1.1	6.1	16.9	31.1	49.5	78.9	144.4
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001 .....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.3	3.5
2000 .....	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.2
1999 .....	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.0	4.5
<b>Female</b>												
2002 .....	28.3	*	*	*	*	2.1	12.0	31.4	56.2	84.4	125.9	191.5
2001 .....	28.6	*	*	*	*	2.4	12.4	32.8	57.5	85.8	125.8	188.9
2000 .....	29.2	*	*	*	*	2.3	12.4	33.0	59.3	88.3	128.9	205.7
1999 .....	28.9	*	*	*	*	2.2	12.0	32.9	59.2	88.9	128.9	200.8
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	4.5	*	*	*	*	0.7	3.8	11.0	17.6	27.4	43.7	71.0
2001 .....	4.8	*	*	*	*	0.8	4.5	12.0	22.0	28.0	39.5	68.8
2000 .....	4.8	*	*	*	*	0.8	4.3	12.1	21.0	31.3	45.6	71.7
1999 .....	4.6	*	*	*	*	0.7	4.0	12.2	19.6	28.0	43.5	86.3
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	9.2	*	*	*	*	1.6	7.8	21.6	33.5	48.7	73.1	105.3
2001 .....	9.7	*	*	*	*	1.7	9.2	23.8	41.5	49.3	64.7	102.4
2000 .....	9.7	*	*	*	*	1.8	8.7	23.9	39.1	54.9	74.9	105.8
1999 .....	9.3	*	*	*	*	1.6	8.3	23.9	36.6	49.1	71.8	127.9
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	16.0	*	*	*	*	1.1	6.4	16.6	30.3	47.7	78.0	136.8
2001 .....	16.1	*	*	*	*	1.2	6.5	17.2	30.7	48.8	78.4	136.4
2000 .....	16.5	*	*	*	*	1.2	6.5	17.3	31.9	50.1	80.1	149.7
1999 .....	16.3	*	*	*	*	1.2	6.3	17.3	31.9	50.6	80.2	146.2
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.3	3.5
2001 .....	0.3	*	*	*	*	*	*	0.3	0.6	1.5	2.3	3.7
2000 .....	0.3	*	*	*	*	*	*	0.3	0.7	1.4	2.2	4.2
1999 .....	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.1	4.7
<b>Female</b>												
2002 .....	31.0	*	*	*	*	2.2	12.6	32.3	57.9	86.7	128.2	194.4
2001 .....	31.1	*	*	*	*	2.5	12.8	33.6	58.6	88.1	128.3	191.6
2000 .....	31.8	*	*	*	*	2.4	12.8	33.8	60.7	90.3	130.8	208.6
1999 .....	31.4	*	*	*	*	2.3	12.4	33.6	60.7	91.2	130.9	202.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of breast (C50)</b>												
Non-Hispanic white												
Both sexes												
2002 .....	16.9	*	*	*	*	0.9	5.6	15.2	29.4	47.4	78.0	137.7
2001 .....	17.0	*	*	*	*	1.0	5.8	16.0	29.6	48.2	78.7	137.1
2000 .....	17.4	*	*	*	*	1.1	5.8	16.1	30.8	50.3	80.6	149.6
1999 .....	17.1	*	*	*	*	1.0	5.6	16.0	30.6	50.2	80.3	147.8
Male												
2002 .....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001 .....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.2	3.5
2000 .....	0.4	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.0
1999 .....	0.3	*	*	*	*	*	*	0.2	0.6	1.1	2.0	4.5
Female												
2002 .....	32.9	*	*	*	*	1.9	11.1	30.0	56.7	87.0	128.9	195.8
2001 .....	32.9	*	*	*	*	2.0	11.5	31.6	57.0	88.0	129.4	192.6
2000 .....	33.8	*	*	*	*	2.1	11.6	31.7	59.2	91.4	132.2	208.3
1999 .....	33.3	*	*	*	*	2.0	11.1	31.5	58.9	91.2	131.5	204.9
Non-Hispanic black												
Both sexes												
2002 .....	15.4	*	*	*	*	2.4	12.0	27.7	43.7	61.1	95.3	154.0
2001 .....	15.5	*	*	*	*	2.7	11.5	27.9	45.6	64.6	93.0	153.9
2000 .....	15.3	*	*	*	*	2.3	11.5	28.4	46.4	59.4	91.7	178.7
1999 .....	15.5	*	*	*	*	2.4	12.0	29.2	47.1	64.5	95.6	149.1
Male												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	29.1	*	*	*	*	4.5	22.5	51.0	77.8	102.7	147.1	210.7
2001 .....	29.2	*	*	*	*	5.1	21.5	51.3	81.6	108.4	143.0	208.3
2000 .....	28.9	*	*	*	*	4.3	21.5	52.5	82.5	100.1	141.5	241.6
1999 .....	29.2	*	*	*	*	4.5	22.3	53.7	83.7	108.2	147.2	201.7
<b>Malignant neoplasm of cervix uteri (C53)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.4	*	*	*	*	0.5	1.5	2.2	2.7	3.4	4.1	6.2
2001 .....	1.4	*	*	*	*	0.5	1.6	2.4	2.9	3.2	4.6	7.2
2000 .....	1.5	*	*	*	*	0.6	1.5	2.4	3.1	3.7	4.8	7.3
1999 .....	1.5	*	*	*	*	0.5	1.6	2.5	3.2	3.8	4.6	7.9
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	2.7	*	*	*	*	1.1	3.0	4.4	5.2	6.3	6.9	8.9
2001 .....	2.8	*	*	*	*	1.1	3.1	4.8	5.5	5.8	7.7	10.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of cervix uteri (C53)</b>												
<b>All origins <sup>3</sup></b>												
Female												
2002 .....	2.9	*	*	*	*	1.2	3.1	4.8	5.9	6.8	8.0	10.3
1999 .....	3.0	*	*	*	*	1.1	3.1	4.9	6.2	6.9	7.7	11.1
<b>Hispanic</b>												
Both sexes												
2002 .....	1.0	*	*	*	*	0.4	1.5	2.4	3.6	4.7	6.7	*
2001 .....	1.1	*	*	*	*	0.5	1.9	3.0	4.1	3.4	7.0	*
2000 .....	1.2	*	*	*	*	0.5	2.1	2.8	4.6	5.6	8.5	13.9
1999 .....	1.1	*	*	*	*	0.5	1.8	3.3	3.9	4.9	6.5	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	2.1	*	*	*	*	0.9	3.1	4.7	6.8	8.4	11.2	*
2001 .....	2.3	*	*	*	*	1.1	3.9	6.0	7.7	6.1	11.8	*
2000 .....	2.5	*	*	*	*	1.1	4.4	5.5	8.6	10.0	14.2	20.8
1999 .....	2.3	*	*	*	*	1.1	3.6	6.6	7.4	8.8	10.8	*
<b>Non-Hispanic <sup>4</sup></b>												
Both sexes												
2002 .....	1.4	*	*	*	*	0.5	1.5	2.2	2.6	3.4	4.0	6.1
2001 .....	1.5	*	*	*	*	0.5	1.5	2.4	2.8	3.2	4.5	7.0
2000 .....	1.5	*	*	*	*	0.6	1.5	2.4	3.0	3.6	4.7	7.1
1999 .....	1.6	*	*	*	*	0.5	1.5	2.4	3.2	3.7	4.6	7.9
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	2.8	*	*	*	*	1.1	2.9	4.3	5.1	6.2	6.7	8.7
2001 .....	2.9	*	*	*	*	1.0	3.0	4.6	5.3	5.8	7.5	10.0
2000 .....	3.0	*	*	*	*	1.2	2.9	4.7	5.7	6.6	7.7	9.9
1999 .....	3.0	*	*	*	*	1.1	3.0	4.7	6.1	6.7	7.5	11.0
<b>Non-Hispanic white</b>												
Both sexes												
2002 .....	1.3	*	*	*	*	0.5	1.3	1.9	2.3	2.7	3.4	4.9
2001 .....	1.4	*	*	*	*	0.5	1.4	2.1	2.4	2.6	3.8	6.1
2000 .....	1.4	*	*	*	*	0.5	1.3	2.1	2.5	2.9	4.1	6.0
1999 .....	1.4	*	*	*	*	0.6	1.4	2.3	2.7	3.0	3.9	6.7
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of cervix uteri (C53)</b>												
Non-Hispanic white												
Female												
2002 .....	2.5	*	*	*	*	1.0	2.7	3.8	4.5	5.0	5.7	7.0
2001 .....	2.7	*	*	*	*	1.0	2.8	4.2	4.7	4.9	6.3	8.6
2000 .....	2.7	*	*	*	*	1.1	2.5	4.2	4.8	5.3	6.8	8.4
1999 .....	2.8	*	*	*	*	1.1	2.8	4.5	5.2	5.5	6.5	9.3
Non-Hispanic black												
Both sexes												
2002 .....	2.3	*	*	*	*	0.8	2.6	4.2	5.1	9.2	10.7	17.5
2001 .....	2.2	*	*	*	*	0.8	2.5	3.8	5.3	7.9	12.0	19.0
2000 .....	2.5	*	*	*	*	1.0	2.6	4.3	6.4	9.8	12.0	19.7
1999 .....	2.4	*	*	*	*	0.7	2.5	3.6	7.2	10.8	11.3	21.7
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	4.3	*	*	*	*	1.6	4.9	7.8	9.2	15.8	16.7	24.2
2001 .....	4.2	*	*	*	*	1.6	4.6	7.1	9.6	13.5	18.7	26.1
2000 .....	4.7	*	*	*	*	1.9	5.0	8.0	11.5	16.6	18.8	27.0
1999 .....	4.5	*	*	*	*	1.2	4.8	6.6	13.0	18.3	17.7	29.8
<b>Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	2.4	*	*	*	*	0.1	0.3	1.4	4.5	10.0	15.4	24.6
2001 .....	2.4	*	*	*	*	0.1	0.3	1.4	4.4	9.7	16.4	24.7
2000 .....	2.3	*	*	*	*	0.1	0.3	1.3	4.6	9.7	15.8	24.7
1999 .....	2.3	*	*	*	*	0.1	0.3	1.3	4.3	9.8	16.4	24.0
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	4.7	*	*	*	*	0.1	0.6	2.8	8.7	18.4	25.6	35.2
2001 .....	4.7	*	*	*	*	0.2	0.6	2.8	8.4	17.8	27.2	35.0
2000 .....	4.6	*	*	*	*	0.1	0.6	2.6	8.9	17.7	26.2	34.8
1999 .....	4.5	*	*	*	*	0.1	0.6	2.5	8.2	17.8	27.1	33.7
Hispanic												
Both sexes												
2002 .....	0.9	*	*	*	*	*	0.3	1.1	3.2	9.0	11.0	20.5
2001 .....	0.9	*	*	*	*	*	*	1.4	3.8	7.5	11.7	18.8
2000 .....	0.8	*	*	*	*	*	*	1.1	3.6	7.0	10.7	19.9
1999 .....	0.8	*	*	*	*	*	*	1.1	3.7	8.3	10.6	15.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Hispanic												
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	1.8	*	*	*	*	*	0.7	2.1	6.2	16.1	18.5	30.7
2001 .....	1.8	*	*	*	*	*	*	2.8	7.3	13.4	19.6	27.9
2000 .....	1.6	*	*	*	*	*	*	2.2	6.8	12.4	17.8	29.7
1999 .....	1.6	*	*	*	*	*	*	2.2	7.0	14.8	17.7	22.9
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	2.6	*	*	*	*	*	0.3	1.4	4.6	10.1	15.5	24.6
2001 .....	2.6	*	*	*	*	0.1	0.3	1.4	4.4	9.9	16.6	24.8
2000 .....	2.6	*	*	*	*	0.1	0.3	1.4	4.7	9.8	16.0	24.9
1999 .....	2.5	*	*	*	*	0.1	0.3	1.3	4.3	9.9	16.6	24.3
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	5.1	*	*	*	*	*	0.6	2.8	8.8	18.5	25.8	35.3
2001 .....	5.1	*	*	*	*	0.2	0.6	2.8	8.5	18.1	27.5	35.1
2000 .....	5.0	*	*	*	*	0.1	0.6	2.7	9.0	17.9	26.5	34.9
1999 .....	4.9	*	*	*	*	0.1	0.6	2.5	8.3	18.0	27.4	34.1
Non-Hispanic white												
Both sexes												
2002 .....	2.6	*	*	*	*	*	0.3	1.3	4.1	8.9	14.6	23.8
2001 .....	2.7	*	*	*	*	*	0.3	1.3	4.0	8.9	15.9	24.5
2000 .....	2.6	*	*	*	*	*	0.3	1.3	4.3	8.6	15.0	24.5
1999 .....	2.6	*	*	*	*	*	0.3	1.3	4.0	8.9	16.0	23.6
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	5.1	*	*	*	*	*	0.6	2.7	8.0	16.5	24.4	34.1
2001 .....	5.2	*	*	*	*	*	0.6	2.7	7.8	16.4	26.5	34.7
2000 .....	5.1	*	*	*	*	*	0.5	2.7	8.4	15.9	24.8	34.3
1999 .....	5.1	*	*	*	*	*	0.6	2.5	7.7	16.3	26.4	33.1
Non-Hispanic black												
Both sexes												
2002 .....	3.1	*	*	*	*	*	0.4	2.0	8.6	22.7	28.4	39.6
2001 .....	2.9	*	*	*	*	*	0.4	2.1	7.7	21.6	28.6	32.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Non-Hispanic black												
Both sexes												
2000 .....	3.0	*	*	*	*	*	0.4	1.7	8.4	22.9	31.1	33.7
1999 .....	2.8	*	*	*	*	*	0.5	1.4	8.1	21.5	27.5	35.9
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	5.8	*	*	*	*	*	0.7	3.6	15.5	38.7	44.4	54.8
2001 .....	5.5	*	*	*	*	*	0.7	4.0	13.9	36.8	44.6	44.9
2000 .....	5.7	*	*	*	*	*	0.7	3.1	15.3	39.1	48.6	46.2
1999 .....	5.4	*	*	*	*	*	0.9	2.7	14.7	36.7	42.9	49.3
Malignant neoplasm of ovary (C56)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	5.1	*	*	*	0.1	0.2	1.0	4.3	10.3	19.7	33.4	38.1
2001 .....	5.1	*	*	*	0.1	0.3	1.0	4.1	10.3	20.0	33.2	39.6
2000 .....	5.0	*	*	*	0.1	0.3	1.0	4.2	10.1	19.6	33.2	40.9
1999 .....	4.9	*	*	*	0.1	0.2	1.1	4.2	10.2	19.6	31.7	37.6
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	10.0	*	*	*	0.2	0.5	2.1	8.4	19.9	36.1	55.6	54.6
2001 .....	9.9	*	*	*	0.1	0.5	2.1	8.1	19.9	36.6	55.0	56.1
2000 .....	9.8	*	*	*	0.1	0.5	2.1	8.3	19.4	35.7	54.8	57.6
1999 .....	9.6	*	*	*	0.1	0.4	2.2	8.3	19.6	35.7	52.3	52.8
Hispanic												
Both sexes												
2002 .....	1.6	*	*	*	*	*	0.8	3.2	5.6	12.3	24.2	23.8
2001 .....	1.6	*	*	*	*	*	0.8	2.7	7.1	14.4	20.0	30.6
2000 .....	1.6	*	*	*	*	*	0.6	3.0	6.5	14.3	24.9	32.5
1999 .....	1.4	*	*	*	*	*	0.6	2.7	6.1	12.2	20.8	23.9
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	3.3	*	*	*	*	*	1.7	6.3	10.7	22.1	40.8	35.6
2001 .....	3.4	*	*	*	*	*	1.6	5.3	13.4	25.8	33.5	45.6
2000 .....	3.4	*	*	*	*	*	1.2	5.9	12.2	25.5	41.6	48.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
Hispanic												
Female												
1999 .....	2.9	*	*	*	*	*	1.2	5.3	11.5	21.7	34.7	35.4
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	5.6	*	*	*	0.1	0.3	1.1	4.4	10.7	20.2	33.8	38.7
2001 .....	5.6	*	*	*	0.1	0.3	1.1	4.3	10.5	20.3	33.7	39.8
2000 .....	5.5	*	*	*	*	0.3	1.1	4.3	10.3	19.9	33.4	41.2
1999 .....	5.3	*	*	*	0.1	0.2	1.2	4.4	10.5	20.0	32.0	38.1
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	11.0	*	*	*	0.2	0.5	2.1	8.6	20.6	37.0	56.3	55.3
2001 .....	10.9	*	*	*	0.1	0.5	2.1	8.4	20.3	37.3	55.8	56.4
2000 .....	10.7	*	*	*	*	0.5	2.2	8.4	19.9	36.3	55.2	57.8
1999 .....	10.4	*	*	*	0.2	0.4	2.3	8.6	20.1	36.4	52.7	53.3
Non-Hispanic white												
Both sexes												
2002 .....	6.3	*	*	*	0.1	0.3	1.1	4.6	11.1	20.9	34.8	39.6
2001 .....	6.2	*	*	*	*	0.3	1.1	4.5	10.8	21.0	34.8	40.6
2000 .....	6.2	*	*	*	*	0.3	1.1	4.5	10.8	20.5	34.5	42.2
1999 .....	6.0	*	*	*	0.1	0.2	1.2	4.5	10.9	20.5	32.7	39.2
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	12.4	*	*	*	0.2	0.5	2.2	9.1	21.7	38.7	58.0	56.7
2001 .....	12.2	*	*	*	*	0.5	2.2	9.0	21.0	38.8	57.8	57.5
2000 .....	12.0	*	*	*	*	0.6	2.3	8.9	21.0	37.7	57.2	59.2
1999 .....	11.7	*	*	*	0.2	0.4	2.4	9.0	21.2	37.7	54.1	54.9
Non-Hispanic black												
Both sexes												
2002 .....	3.2	*	*	*	*	*	1.1	3.5	9.2	17.4	29.3	35.4
2001 .....	3.2	*	*	*	*	*	1.2	3.1	10.2	18.5	27.2	36.5
2000 .....	3.0	*	*	*	*	*	1.1	3.4	8.6	18.2	27.1	34.7
1999 .....	3.1	*	*	*	*	*	0.9	3.6	9.1	18.7	28.7	29.1
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	6.1	*	*	*	*	*	2.0	6.5	16.6	29.8	45.9	48.9

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of ovary (C56)</b>												
Non-Hispanic black												
Female												
2001 .....	6.1	*	*	*	*	*	2.2	5.7	18.4	31.5	42.5	50.0
2000 .....	5.8	*	*	*	*	*	2.0	6.3	15.6	31.1	42.4	47.5
1999 .....	5.8	*	*	*	*	*	1.6	6.7	16.4	31.9	44.8	40.0
<b>Malignant neoplasm of prostate (C61)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	10.6	*	*	*	*	*	0.1	1.0	7.6	34.5	100.1	195.0
2001 .....	10.8	*	*	*	*	*	0.1	1.1	7.5	35.7	103.2	200.9
2000 .....	11.0	*	*	*	*	*	0.1	1.0	7.9	37.0	105.7	209.3
1999 .....	11.4	*	*	*	*	*	0.1	1.0	8.2	39.7	109.9	207.5
Male												
2002 .....	21.5	*	*	*	*	*	0.1	1.9	15.8	75.9	251.0	644.5
2001 .....	22.0	*	*	*	*	*	0.1	2.1	15.7	78.9	260.2	681.2
2000 .....	22.5	*	*	*	*	*	0.1	2.0	16.6	82.0	267.9	723.1
1999 .....	23.2	*	*	*	*	*	0.1	2.1	17.1	88.2	279.1	722.4
Female												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic												
Both sexes												
2002 .....	3.1	*	*	*	*	*	*	0.8	5.4	24.7	78.7	170.7
2001 .....	3.3	*	*	*	*	*	*	0.8	5.8	29.2	87.8	170.1
2000 .....	2.9	*	*	*	*	*	*	0.8	4.6	25.7	79.2	165.9
1999 .....	3.1	*	*	*	*	*	*	0.7	5.6	28.5	80.4	176.1
Male												
2002 .....	6.0	*	*	*	*	*	*	1.6	11.3	55.8	193.7	515.4
2001 .....	6.4	*	*	*	*	*	*	1.6	12.4	66.3	217.8	518.0
2000 .....	5.7	*	*	*	*	*	*	1.7	9.7	58.5	197.2	503.9
1999 .....	6.0	*	*	*	*	*	*	1.5	12.0	64.8	201.3	541.3
Female												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	11.7	*	*	*	*	*	0.1	1.0	7.7	35.0	101.0	195.5
2001 .....	11.9	*	*	*	*	*	0.1	1.1	7.6	36.1	103.6	201.6
2000 .....	12.2	*	*	*	*	*	0.1	1.0	8.2	37.6	106.6	210.4
1999 .....	12.5	*	*	*	*	*	0.1	1.0	8.3	40.3	110.9	208.2
Male												
2002 .....	24.0	*	*	*	*	*	0.2	2.0	16.1	76.9	253.3	648.7
2001 .....	24.3	*	*	*	*	*	0.1	2.2	15.9	79.5	261.4	686.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of prostate (C61)</b>												
Non-Hispanic <sup>4</sup>												
Male												
2000 .....	25.0	*	*	*	*	*	0.1	2.0	17.0	83.2	270.2	730.7
1999 .....	25.6	*	*	*	*	*	0.2	2.1	17.4	89.4	281.8	728.2
Female												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white												
Both sexes												
2002 .....	11.9	*	*	*	*	*	0.1	0.8	6.5	30.9	93.1	186.6
2001 .....	12.0	*	*	*	*	*	*	0.9	6.4	31.9	95.1	189.5
2000 .....	12.3	*	*	*	*	*	*	0.8	6.9	32.8	97.6	199.3
1999 .....	12.6	*	*	*	*	*	0.1	0.9	6.9	35.6	102.6	195.4
Male												
2002 .....	24.3	*	*	*	*	*	0.2	1.5	13.4	67.1	232.1	618.5
2001 .....	24.6	*	*	*	*	*	*	1.8	13.3	69.6	238.4	645.0
2000 .....	25.2	*	*	*	*	*	*	1.7	14.2	71.7	246.1	693.6
1999 .....	25.9	*	*	*	*	*	0.1	1.8	14.3	78.2	259.4	685.4
Female												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black												
Both sexes												
2002 .....	14.1	*	*	*	*	*	*	2.6	19.5	82.1	212.3	333.3
2001 .....	14.6	*	*	*	*	*	*	2.9	19.5	84.2	221.6	365.8
2000 .....	15.1	*	*	*	*	*	*	2.4	20.8	91.6	230.8	359.6
1999 .....	15.3	*	*	*	*	*	*	2.7	22.0	94.7	227.5	368.6
Male												
2002 .....	29.6	*	*	*	*	*	*	5.7	43.8	197.9	587.6	1,205.5
2001 .....	30.8	*	*	*	*	*	*	6.2	43.7	203.5	616.3	1,353.6
2000 .....	31.7	*	*	*	*	*	*	5.3	46.6	221.7	640.8	1,328.0
1999 .....	32.2	*	*	*	*	*	*	5.8	49.5	229.1	633.8	1,354.9
Female												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Malignant neoplasms of kidney and renal pelvis (C64-C65)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	4.2	*	*	0.1	0.0	0.1	0.7	3.3	8.8	17.7	26.3	32.5
2001 .....	4.2	*	0.1	0.1	0.1	0.1	0.8	3.3	9.2	17.6	26.3	32.3
2000 .....	4.2	*	*	0.1	0.1	0.1	0.7	3.3	9.0	17.1	26.6	33.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms of kidney and renal pelvis (C64-C65)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
1999 .....	4.0	*	*	0.1	*	0.1	0.8	3.1	8.9	16.7	24.6	31.4
<b>Male</b>												
2002 .....	5.4	*	*	*	*	0.2	1.0	4.8	13.0	25.1	37.9	50.1
2001 .....	5.4	*	*	*	*	0.1	1.1	4.9	13.4	25.2	37.4	52.9
2000 .....	5.3	*	*	*	*	0.1	1.0	4.8	13.1	24.2	38.2	55.1
1999 .....	5.0	*	*	*	*	0.2	1.1	4.5	12.9	23.6	35.7	49.4
<b>Female</b>												
2002 .....	3.1	*	*	*	*	*	0.4	1.8	4.9	11.6	18.7	24.9
2001 .....	3.1	*	*	0.1	*	0.1	0.5	1.9	5.3	11.3	19.0	23.7
2000 .....	3.1	*	*	*	*	*	0.5	1.9	5.3	11.3	18.9	25.1
1999 .....	3.0	*	*	*	*	*	0.5	1.8	5.1	11.0	17.4	24.1
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	1.7	*	*	*	*	*	0.6	2.8	8.1	12.9	24.4	26.1
2001 .....	1.7	*	*	*	*	*	0.7	3.3	6.9	14.4	23.3	31.9
2000 .....	1.6	*	*	*	*	*	0.6	2.8	7.9	12.3	24.7	25.9
1999 .....	1.6	*	*	*	*	*	0.6	2.8	7.9	13.9	22.5	28.8
<b>Male</b>												
2002 .....	2.1	*	*	*	*	*	0.9	3.9	11.3	18.5	39.0	40.2
2001 .....	2.2	*	*	*	*	*	1.2	4.7	9.9	21.4	35.3	57.1
2000 .....	2.0	*	*	*	*	*	1.0	3.9	11.4	17.9	38.4	40.3
1999 .....	2.0	*	*	*	*	*	0.9	4.3	11.8	21.1	29.7	*
<b>Female</b>												
2002 .....	1.2	*	*	*	*	*	*	1.7	5.2	8.4	14.3	19.1
2001 .....	1.2	*	*	*	*	*	*	1.9	4.2	8.8	15.1	19.5
2000 .....	1.1	*	*	*	*	*	*	1.8	4.7	7.8	15.5	*
1999 .....	1.2	*	*	*	*	*	*	1.4	4.4	8.3	17.7	23.9
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	4.6	*	*	0.1	*	0.1	0.7	3.3	8.8	18.0	26.4	32.7
2001 .....	4.6	*	*	0.1	0.1	0.1	0.8	3.3	9.4	17.8	26.4	32.3
2000 .....	4.5	*	*	0.1	*	0.1	0.8	3.3	9.1	17.3	26.6	33.9
1999 .....	4.3	*	*	0.1	*	0.1	0.8	3.2	8.9	16.8	24.6	31.4
<b>Male</b>												
2002 .....	5.9	*	*	*	*	0.2	1.0	4.9	13.0	25.5	37.7	50.4
2001 .....	5.9	*	*	*	*	0.1	1.0	4.9	13.7	25.4	37.5	52.6
2000 .....	5.7	*	*	*	*	0.1	1.0	4.8	13.2	24.4	38.1	55.4
1999 .....	5.4	*	*	*	*	0.2	1.1	4.6	13.0	23.7	35.9	49.8
<b>Female</b>												
2002 .....	3.4	*	*	*	*	*	0.4	1.8	4.9	11.8	18.8	25.1
2001 .....	3.4	*	*	0.1	*	0.1	0.6	1.8	5.4	11.4	19.1	23.8
2000 .....	3.4	*	*	*	*	*	0.5	1.9	5.3	11.4	19.1	25.2
1999 .....	3.2	*	*	*	*	*	0.5	1.8	5.2	11.2	17.3	24.1
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	5.1	*	*	0.1	*	0.1	0.7	3.4	9.1	18.7	26.8	33.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasms of kidney and renal pelvis (C64-C65)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	5.1	*	*	*	*	0.1	0.7	3.4	9.6	18.4	26.8	32.8
2001 .....	5.0	*	*	*	*	0.1	0.7	3.4	9.3	17.7	26.9	34.6
2000 .....	4.7	*	*	0.1	*	0.1	0.7	3.1	9.2	17.2	24.9	31.5
1999 .....	4.7	*	*	0.1	*	0.1	0.7	3.1	9.2	17.2	24.9	31.5
<b>Male</b>												
2002 .....	6.5	*	*	*	*	*	1.0	5.0	13.3	26.3	38.4	52.4
2001 .....	6.5	*	*	*	*	*	1.0	4.9	13.8	26.2	37.7	54.3
2000 .....	6.3	*	*	*	*	*	1.0	4.9	13.5	24.6	38.5	56.9
1999 .....	6.0	*	*	*	*	*	1.0	4.5	13.3	24.0	36.5	50.2
<b>Female</b>												
2002 .....	3.8	*	*	*	*	*	0.4	1.8	5.1	12.2	19.0	25.3
2001 .....	3.8	*	*	*	*	*	0.5	1.8	5.5	11.8	19.5	23.8
2000 .....	3.8	*	*	*	*	*	0.5	1.9	5.4	11.8	19.3	25.7
1999 .....	3.5	*	*	*	*	*	0.5	1.8	5.3	11.5	17.3	24.1
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	2.9	*	*	*	*	*	0.8	3.6	8.8	15.8	26.5	28.1
2001 .....	3.1	*	*	*	*	*	1.3	3.8	9.7	15.3	26.0	30.0
2000 .....	2.9	*	*	*	*	*	0.9	3.5	9.0	16.8	25.9	29.6
1999 .....	3.0	*	*	*	*	0.4	1.1	3.9	8.7	16.4	23.9	31.7
<b>Male</b>												
2002 .....	3.6	*	*	*	*	*	1.2	5.0	14.4	22.1	36.6	39.4
2001 .....	4.0	*	*	*	*	*	1.5	5.5	15.3	22.7	40.8	46.2
2000 .....	3.7	*	*	*	*	*	1.0	5.0	13.4	26.8	38.3	49.3
1999 .....	3.7	*	*	*	*	*	1.6	6.0	13.3	23.9	32.5	49.9
<b>Female</b>												
2002 .....	2.3	*	*	*	*	*	*	2.5	4.4	11.3	20.7	23.8
2001 .....	2.3	*	*	*	*	*	1.1	2.2	5.3	10.0	17.7	24.0
2000 .....	2.2	*	*	*	*	*	0.8	2.2	5.4	9.7	18.9	22.2
1999 .....	2.2	*	*	*	*	*	*	2.1	5.0	11.0	19.1	24.9
<b>Malignant neoplasm of bladder (C67)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	4.4	*	*	*	*	*	0.3	1.3	4.9	15.9	37.0	66.6
2001 .....	4.3	*	*	*	*	*	0.3	1.2	5.0	15.4	35.7	68.9
2000 .....	4.3	*	*	*	*	*	0.2	1.2	5.0	15.5	35.7	69.6
1999 .....	4.3	*	*	*	*	*	0.2	1.3	5.2	15.9	35.3	68.5
<b>Male</b>												
2002 .....	6.0	*	*	*	*	*	0.3	1.8	7.7	25.2	63.0	128.9
2001 .....	6.0	*	*	*	*	*	0.3	1.8	7.9	25.2	62.5	134.8
2000 .....	5.9	*	*	*	*	*	0.3	1.7	7.9	25.2	62.6	139.4
1999 .....	5.9	*	*	*	*	*	0.3	2.0	8.2	25.7	62.5	134.8
<b>Female</b>												
2002 .....	2.8	*	*	*	*	*	0.2	0.7	2.4	8.2	19.7	39.6
2001 .....	2.7	*	*	*	*	*	0.3	0.6	2.3	7.2	18.1	41.4
2000 .....	2.7	*	*	*	*	*	0.2	0.6	2.4	7.5	18.1	41.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
All origins <sup>3</sup>												
Female												
1999 .....	2.7	*	*	*	*	*	0.2	0.6	2.5	7.8	17.7	41.8
Hispanic												
Both sexes												
2002 .....	1.0	*	*	*	*	*	*	0.7	2.9	7.7	18.3	54.9
2001 .....	0.9	*	*	*	*	*	*	0.6	2.2	7.9	18.8	46.3
2000 .....	0.9	*	*	*	*	*	*	*	2.9	6.6	17.6	47.1
1999 .....	0.9	*	*	*	*	*	*	0.7	3.1	8.5	16.5	38.6
Male												
2002 .....	1.3	*	*	*	*	*	*	*	4.2	12.8	27.9	100.4
2001 .....	1.2	*	*	*	*	*	*	*	4.3	11.8	30.0	87.6
2000 .....	1.2	*	*	*	*	*	*	*	4.6	10.6	33.9	84.6
1999 .....	1.3	*	*	*	*	*	*	*	5.6	15.2	27.1	77.6
Female												
2002 .....	0.7	*	*	*	*	*	*	*	*	3.6	11.8	32.3
2001 .....	0.6	*	*	*	*	*	*	*	*	4.9	11.2	26.1
2000 .....	0.5	*	*	*	*	*	*	*	*	3.5	6.6	28.7
1999 .....	0.5	*	*	*	*	*	*	*	*	*	9.4	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	4.9	*	*	*	*	*	0.3	1.3	5.1	16.4	37.8	66.9
2001 .....	4.8	*	*	*	*	*	0.3	1.3	5.2	15.8	36.4	69.6
2000 .....	4.7	*	*	*	*	*	0.3	1.2	5.2	16.1	36.3	70.3
1999 .....	4.7	*	*	*	*	*	0.2	1.3	5.4	16.3	36.0	69.3
Male												
2002 .....	6.8	*	*	*	*	*	0.4	1.9	7.9	26.0	64.7	129.8
2001 .....	6.7	*	*	*	*	*	0.4	1.9	8.1	26.0	63.8	136.4
2000 .....	6.6	*	*	*	*	*	0.3	1.8	8.1	26.1	63.6	141.3
1999 .....	6.6	*	*	*	*	*	0.3	2.0	8.4	26.3	63.8	136.6
Female												
2002 .....	3.1	*	*	*	*	*	0.2	0.8	2.4	8.5	20.0	39.8
2001 .....	3.0	*	*	*	*	*	0.3	0.6	2.5	7.4	18.3	41.9
2000 .....	3.0	*	*	*	*	*	0.2	0.6	2.5	7.8	18.5	41.6
1999 .....	3.0	*	*	*	*	*	0.2	0.6	2.6	8.1	17.9	42.3
Non-Hispanic white												
Both sexes												
2002 .....	5.6	*	*	*	*	*	0.3	1.4	5.3	17.1	39.1	68.0
2001 .....	5.5	*	*	*	*	*	0.3	1.3	5.4	16.6	37.4	71.5
2000 .....	5.4	*	*	*	*	*	0.3	1.3	5.5	16.8	37.1	72.8
1999 .....	5.4	*	*	*	*	*	0.2	1.3	5.5	16.8	37.1	70.7
Male												
2002 .....	7.9	*	*	*	*	*	0.4	2.0	8.5	27.4	67.2	135.2
2001 .....	7.8	*	*	*	*	*	0.4	2.0	8.6	27.6	66.6	143.0
2000 .....	7.7	*	*	*	*	*	0.3	2.0	8.6	27.5	65.9	148.0
1999 .....	7.6	*	*	*	*	*	0.3	2.0	8.7	27.6	66.2	143.0
Female												
2002 .....	3.4	*	*	*	*	*	0.2	0.8	2.3	8.4	20.2	39.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malignant neoplasm of bladder (C67)</b>												
<b>Non-Hispanic white</b>												
<b>Female</b>												
2001 .....	3.2	*	*	*	*	*	0.3	0.6	2.4	7.4	18.1	41.7
2000 .....	3.3	*	*	*	*	*	0.2	0.7	2.6	7.8	18.2	42.5
1999 .....	3.2	*	*	*	*	*	0.2	0.7	2.5	7.8	18.0	41.9
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	2.6	*	*	*	*	*	*	1.1	4.5	14.7	31.2	64.1
2001 .....	2.4	*	*	*	*	*	0.4	1.3	5.1	12.1	31.4	55.6
2000 .....	2.4	*	*	*	*	*	*	1.1	4.2	13.5	34.9	49.6
1999 .....	2.6	*	*	*	*	*	*	1.5	5.6	15.2	29.4	60.8
<b>Male</b>												
2002 .....	2.8	*	*	*	*	*	*	1.3	5.5	18.7	48.1	91.9
2001 .....	2.6	*	*	*	*	*	*	1.8	6.5	16.5	43.2	78.5
2000 .....	2.8	*	*	*	*	*	*	1.4	5.9	17.8	52.8	88.1
1999 .....	3.0	*	*	*	*	*	*	2.6	7.4	19.2	46.6	86.8
<b>Female</b>												
2002 .....	2.4	*	*	*	*	*	*	0.9	3.7	11.8	21.7	53.5
2001 .....	2.3	*	*	*	*	*	*	*	3.9	9.0	24.8	47.1
2000 .....	2.1	*	*	*	*	*	*	*	2.7	10.4	24.8	35.3
1999 .....	2.3	*	*	*	*	*	*	*	4.2	12.3	19.8	51.1
<b>Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	4.4	*	0.7	0.8	0.5	1.0	2.2	5.1	9.6	16.2	19.1	16.4
2001 .....	4.4	*	0.7	0.7	0.6	1.1	2.3	5.0	9.7	15.4	20.4	16.4
2000 .....	4.5	0.5	0.7	0.8	0.6	1.0	2.4	5.1	9.6	16.5	20.0	16.9
1999 .....	4.6	*	0.7	0.8	0.6	1.1	2.4	5.1	10.3	16.9	20.4	16.0
<b>Male</b>												
2002 .....	5.0	*	0.8	0.9	0.7	1.1	2.8	6.2	11.7	20.2	23.8	20.1
2001 .....	5.0	*	0.7	0.7	0.6	1.3	2.8	6.1	12.1	19.2	25.8	20.1
2000 .....	5.1	*	0.7	0.8	0.7	1.3	3.1	6.2	11.8	20.2	24.3	22.4
1999 .....	5.1	*	0.7	0.7	0.7	1.4	2.8	6.2	12.4	20.8	25.5	20.7
<b>Female</b>												
2002 .....	3.9	*	0.7	0.7	0.4	0.8	1.7	4.1	7.7	12.8	16.0	14.8
2001 .....	3.9	*	0.6	0.8	0.5	0.8	1.7	3.9	7.6	12.3	16.8	14.8
2000 .....	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.6	13.4	17.3	14.7
1999 .....	4.1	*	0.8	0.8	0.5	0.8	1.9	4.1	8.4	13.7	17.1	14.1
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	1.7	*	0.7	0.7	0.4	0.5	0.9	3.4	5.8	10.0	13.7	13.3
2001 .....	1.7	*	0.7	0.5	0.4	0.6	1.4	2.6	5.9	10.3	14.8	*
2000 .....	1.8	*	*	0.9	0.4	0.7	1.5	2.9	6.2	9.8	14.2	15.9
1999 .....	1.7	*	0.9	0.7	*	0.5	1.3	3.0	6.0	10.0	14.0	*
<b>Male</b>												
2002 .....	1.8	*	*	0.7	*	*	1.0	4.4	6.6	12.4	16.0	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Hispanic												
Male												
2001 .....	1.8	*	*	*	*	0.6	1.4	3.0	6.8	14.4	17.9	*
2000 .....	1.9	*	*	1.2	*	0.8	1.5	2.6	7.7	12.5	18.7	*
1999 .....	1.7	*	*	0.7	*	*	1.8	2.9	7.8	10.5	16.2	*
Female												
2002 .....	1.5	*	*	0.7	*	*	0.9	2.3	5.1	8.1	12.1	*
2001 .....	1.6	*	*	0.6	*	*	1.4	2.2	5.1	7.1	12.7	*
2000 .....	1.7	*	*	0.7	*	*	1.5	3.2	4.9	7.6	11.2	*
1999 .....	1.7	*	*	0.8	*	*	0.8	3.1	4.5	9.6	12.5	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	4.9	*	0.7	0.8	0.5	1.1	2.4	5.3	9.9	16.6	19.4	16.5
2001 .....	4.8	*	0.6	0.8	0.6	1.2	2.4	5.2	10.0	15.7	20.6	16.5
2000 .....	4.9	*	0.7	0.7	0.6	1.1	2.5	5.3	9.8	16.9	20.2	16.9
1999 .....	5.0	*	0.7	0.8	0.6	1.2	2.5	5.3	10.6	17.3	20.6	16.1
Male												
2002 .....	5.5	*	0.8	0.9	0.7	1.3	3.0	6.4	12.0	20.7	24.1	20.2
2001 .....	5.5	*	0.6	0.7	0.7	1.5	3.0	6.4	12.5	19.4	26.1	20.5
2000 .....	5.5	*	0.7	0.8	0.7	1.4	3.3	6.5	12.1	20.6	24.5	22.3
1999 .....	5.6	*	0.7	0.8	0.7	1.5	3.0	6.5	12.7	21.4	25.8	21.3
Female												
2002 .....	4.2	*	0.7	0.7	0.4	0.9	1.8	4.2	7.9	13.1	16.2	14.9
2001 .....	4.2	*	0.6	0.8	0.5	0.8	1.8	4.1	7.7	12.7	17.0	14.9
2000 .....	4.2	*	0.7	0.7	0.5	0.7	1.8	4.1	7.8	13.8	17.5	14.7
1999 .....	4.4	*	0.7	0.8	0.5	0.9	2.0	4.2	8.6	14.0	17.2	14.0
Non-Hispanic white												
Both sexes												
2002 .....	5.6	*	0.6	0.8	0.6	1.2	2.8	5.9	10.7	17.9	20.4	17.3
2001 .....	5.6	*	0.7	0.8	0.6	1.3	2.7	5.7	10.8	16.8	21.9	17.2
2000 .....	5.6	*	0.7	0.8	0.6	1.2	2.8	5.9	10.7	18.3	21.2	17.5
1999 .....	5.7	*	0.7	0.8	0.6	1.4	2.8	5.8	11.5	18.7	21.7	16.7
Male												
2002 .....	6.4	*	0.7	0.9	0.8	1.4	3.5	7.1	12.8	22.3	25.2	21.4
2001 .....	6.3	*	0.7	0.7	0.7	1.7	3.5	7.0	13.4	20.9	27.5	20.9
2000 .....	6.3	*	0.7	0.8	0.7	1.5	3.7	7.1	13.1	22.2	25.7	23.5
1999 .....	6.4	*	0.8	0.8	0.7	1.8	3.3	7.1	13.7	23.0	27.2	22.4
Female												
2002 .....	4.9	*	0.6	0.8	0.4	1.1	2.1	4.7	8.7	14.2	17.1	15.5
2001 .....	4.8	*	0.6	0.9	0.5	0.9	2.0	4.5	8.4	13.4	18.2	15.7
2000 .....	4.9	*	0.7	0.7	0.5	0.9	1.9	4.6	8.4	15.0	18.3	15.1
1999 .....	5.0	*	0.7	0.7	0.5	1.1	2.3	4.6	9.5	15.1	18.1	14.4
Non-Hispanic black												
Both sexes												
2002 .....	2.1	*	1.1	0.8	0.4	0.6	1.2	2.8	5.6	8.6	11.9	9.7
2001 .....	2.1	*	*	0.7	0.5	0.5	1.1	3.1	5.9	9.4	9.9	11.9
2000 .....	2.2	*	*	0.8	0.6	0.7	1.6	2.9	5.8	8.2	12.1	10.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Non-Hispanic black												
Both sexes												
1999 .....	2.2	*	*	0.8	0.7	0.7	1.4	3.3	6.1	8.2	11.7	10.0
Male												
2002 .....	2.3	*	*	1.1	*	0.9	1.3	2.7	8.1	10.2	16.1	*
2001 .....	2.2	*	*	0.8	*	*	1.3	3.7	7.9	10.4	12.0	*
2000 .....	2.4	*	*	0.9	0.8	1.1	2.0	3.7	7.2	9.9	13.3	*
1999 .....	2.4	*	*	0.7	0.9	0.9	1.8	4.5	7.4	10.2	12.5	*
Female												
2002 .....	1.9	*	*	*	*	*	1.1	2.8	3.7	7.5	9.5	10.5
2001 .....	1.9	*	*	*	*	*	1.0	2.5	4.2	8.7	8.8	9.8
2000 .....	1.9	*	*	0.7	*	*	1.2	2.2	4.7	7.1	11.5	10.5
1999 .....	1.9	*	*	0.9	*	*	1.0	2.2	5.1	6.8	11.2	9.3
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	19.5	0.7	0.8	0.9	1.7	2.3	3.9	9.9	28.2	73.9	143.6	197.5
2001 .....	19.8	0.7	1.0	0.9	1.8	2.5	4.1	10.4	29.5	74.6	145.2	199.9
2000 .....	20.1	1.0	0.9	0.9	1.8	2.3	4.1	10.7	30.0	76.3	149.2	204.6
1999 .....	20.2	0.8	0.9	0.9	1.9	2.4	4.2	10.8	31.8	77.8	148.0	199.4
Male												
2002 .....	21.4	*	0.8	1.0	2.0	2.7	4.6	11.9	35.0	95.2	188.0	269.1
2001 .....	21.5	*	1.0	1.0	2.1	2.9	5.1	12.7	36.3	94.7	187.0	275.3
2000 .....	21.6	1.1	1.0	1.1	2.3	2.7	4.8	13.0	36.1	96.8	192.6	281.8
1999 .....	21.7	*	1.0	1.0	2.2	2.8	5.1	13.2	38.8	97.3	189.0	279.0
Female												
2002 .....	17.7	*	0.7	0.8	1.4	1.9	3.2	7.9	21.9	56.2	114.1	166.5
2001 .....	18.1	*	1.0	0.7	1.4	2.1	3.2	8.1	23.1	57.9	117.7	168.4
2000 .....	18.6	*	0.9	0.7	1.4	1.9	3.4	8.5	24.4	59.5	120.9	173.1
1999 .....	18.7	*	0.8	0.7	1.5	2.0	3.4	8.5	25.3	61.9	121.3	167.3
Hispanic												
Both sexes												
2002 .....	7.3	*	0.9	1.3	2.1	1.9	3.9	7.7	22.1	53.7	100.8	132.5
2001 .....	7.4	*	1.3	1.2	2.2	2.2	3.9	8.2	23.9	52.6	103.3	134.4
2000 .....	7.5	*	1.0	1.1	2.0	1.9	3.9	8.9	25.0	58.8	103.3	143.3
1999 .....	7.5	*	1.0	1.2	2.0	2.1	3.9	10.1	24.2	58.5	101.0	131.9
Male												
2002 .....	7.6	*	*	1.3	2.7	2.3	4.4	8.9	26.2	63.3	119.0	180.7
2001 .....	8.3	*	1.4	1.3	2.7	2.7	5.0	9.7	28.9	67.5	142.2	180.9
2000 .....	8.0	*	*	1.4	2.3	2.4	4.3	11.1	30.7	71.3	126.9	197.5
1999 .....	7.6	*	*	1.3	2.1	2.2	4.2	12.0	29.4	68.5	117.8	135.9
Female												
2002 .....	7.0	*	*	1.3	1.5	1.4	3.3	6.5	18.5	46.1	88.3	108.6
2001 .....	6.5	*	*	1.1	1.5	1.7	2.6	6.7	19.5	40.8	77.1	111.7
2000 .....	6.9	*	*	0.8	1.5	1.4	3.4	6.8	19.8	48.9	87.5	116.7

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Hispanic												
Female												
1999 .....	7.4	*	*	1.0	1.9	2.1	3.4	8.3	19.6	50.6	89.9	130.0
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	21.3	0.8	0.7	0.8	1.6	2.4	3.9	10.1	28.6	75.1	145.3	199.8
2001 .....	21.6	0.6	0.9	0.8	1.7	2.5	4.1	10.5	29.8	75.8	146.8	202.1
2000 .....	21.8	1.0	0.9	0.9	1.8	2.4	4.1	10.8	30.3	77.3	150.9	206.5
1999 .....	21.9	0.7	0.9	0.8	1.8	2.4	4.3	10.8	32.3	78.8	149.5	201.4
Male												
2002 .....	23.6	*	0.8	1.0	1.8	2.8	4.7	12.1	35.6	97.0	190.9	272.8
2001 .....	23.5	*	1.0	1.0	1.9	2.9	5.0	12.9	36.8	96.3	188.6	278.8
2000 .....	23.7	*	1.0	1.0	2.2	2.7	4.8	13.1	36.4	98.1	195.2	285.0
1999 .....	23.7	*	0.9	0.9	2.2	2.9	5.2	13.3	39.5	98.7	191.5	284.3
Female												
2002 .....	19.2	*	0.7	0.7	1.4	2.0	3.1	8.0	22.2	56.8	115.1	168.3
2001 .....	19.7	*	0.9	0.7	1.4	2.2	3.3	8.2	23.3	58.9	119.3	170.2
2000 .....	20.1	*	0.9	0.7	1.3	2.0	3.4	8.6	24.7	60.1	122.1	174.8
1999 .....	20.1	*	0.8	0.7	1.4	2.0	3.4	8.5	25.6	62.5	122.3	168.2
Non-Hispanic white												
Both sexes												
2002 .....	23.7	*	0.8	0.8	1.7	2.3	3.8	9.7	28.4	76.7	149.5	206.5
2001 .....	23.8	*	1.1	0.8	1.7	2.4	4.0	10.3	29.5	77.2	150.5	208.2
2000 .....	24.2	1.3	1.0	0.9	1.7	2.3	4.0	10.5	30.3	79.0	155.1	210.7
1999 .....	24.1	*	1.0	0.8	1.8	2.4	4.2	10.6	32.0	80.4	153.0	206.4
Male												
2002 .....	26.2	*	0.9	1.0	1.9	2.7	4.5	11.8	35.4	99.3	196.8	283.5
2001 .....	25.9	*	1.0	0.8	2.0	2.8	4.7	12.8	36.4	97.9	193.6	288.7
2000 .....	26.2	*	1.1	1.0	2.1	2.6	4.8	12.7	36.3	100.1	201.0	293.8
1999 .....	26.1	*	1.0	0.9	2.2	2.7	5.0	13.1	39.2	100.7	196.1	291.8
Female												
2002 .....	21.2	*	0.7	0.7	1.4	1.9	3.1	7.6	21.7	57.5	117.8	173.2
2001 .....	21.8	*	1.1	0.7	1.4	2.0	3.2	7.9	23.0	59.5	122.0	174.6
2000 .....	22.3	*	0.9	0.8	1.2	1.9	3.2	8.4	24.6	61.2	125.0	177.2
1999 .....	22.2	*	0.9	0.6	1.3	2.0	3.3	8.3	25.2	63.4	124.7	172.4
Non-Hispanic black												
Both sexes												
2002 .....	14.1	*	*	0.8	1.5	3.0	4.8	13.9	36.4	77.0	123.5	153.7
2001 .....	14.5	*	*	1.0	1.8	3.5	5.8	13.5	37.1	79.0	127.6	157.3
2000 .....	14.3	*	*	0.7	2.3	3.0	5.4	15.0	36.5	76.2	122.3	177.1
1999 .....	14.5	*	*	0.9	2.4	3.0	5.6	14.1	39.5	77.5	128.2	162.3
Male												
2002 .....	15.1	*	*	1.0	1.5	3.8	6.2	16.5	45.2	99.4	155.3	203.5
2001 .....	15.6	*	*	1.3	2.0	4.0	7.6	16.3	46.6	100.7	161.4	206.7
2000 .....	15.2	*	*	0.8	2.8	3.7	5.7	18.2	45.3	97.8	157.4	226.6
1999 .....	15.6	*	*	1.2	2.6	3.9	7.0	17.5	48.4	96.9	162.9	221.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Non-Hispanic black												
Female												
2002 .....	13.1	*	*	*	1.5	2.3	3.5	11.6	29.4	61.2	105.4	134.6
2001 .....	13.5	*	*	*	1.7	3.1	4.2	11.1	29.5	63.7	108.7	139.0
2000 .....	13.5	*	*	*	1.8	2.5	5.2	12.2	29.4	61.0	102.6	158.7
1999 .....	13.6	*	*	0.7	2.1	2.2	4.3	11.1	32.3	63.8	108.7	140.4
Hodgkin's disease (C81)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.7
2001 .....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.2	2.2	2.2
2000 .....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.1	2.5
1999 .....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.5	2.1	2.4
Male												
2002 .....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.5	3.6
2001 .....	0.5	*	*	*	0.2	0.4	0.5	0.6	0.7	1.6	2.6	2.6
2000 .....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.5	3.7
1999 .....	0.6	*	*	*	0.3	0.4	0.5	0.5	1.0	1.9	2.5	3.8
Female												
2002 .....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	1.6	2.3
2001 .....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.8	2.0	2.0
2000 .....	0.4	*	*	*	0.2	0.3	0.3	0.3	0.6	1.0	1.8	2.0
1999 .....	0.4	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.8
Hispanic												
Both sexes												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	1.8	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	4.5	*
1999 .....	0.3	*	*	*	*	*	*	*	*	2.5	*	*
Male												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.8
2001 .....	0.5	*	*	*	0.2	0.5	0.4	0.5	0.5	1.1	2.2	2.2
2000 .....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.4
1999 .....	0.5	*	*	*	0.3	0.4	0.4	0.4	0.7	1.5	2.1	2.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Non-Hispanic <sup>4</sup>												
Male												
2002 .....	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.4	3.7
2001 .....	0.5	*	*	*	0.2	0.5	0.5	0.6	0.7	1.5	2.6	2.5
2000 .....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.3	3.7
1999 .....	0.6	*	*	*	0.3	0.5	0.5	0.5	1.0	1.9	2.3	3.7
Female												
2002 .....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.5	0.8	1.5	2.4
2001 .....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.5	0.8	2.0	2.0
2000 .....	0.4	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	1.7	2.0
1999 .....	0.5	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.7
Non-Hispanic white												
Both sexes												
2002 .....	0.6	*	*	*	0.2	0.5	0.4	0.5	0.6	1.4	2.0	2.9
2001 .....	0.5	*	*	*	0.2	0.5	0.4	0.5	0.6	1.2	2.3	2.3
2000 .....	0.5	*	*	*	0.2	0.5	0.4	0.4	0.6	1.3	2.1	2.6
1999 .....	0.6	*	*	*	0.3	0.4	0.4	0.5	0.8	1.5	2.2	2.3
Male												
2002 .....	0.6	*	*	*	0.2	0.5	0.4	0.5	0.8	1.9	2.6	4.0
2001 .....	0.6	*	*	*	0.3	0.5	0.5	0.6	0.7	1.6	2.7	2.6
2000 .....	0.6	*	*	*	0.2	0.5	0.4	0.5	0.7	1.6	2.5	3.9
1999 .....	0.7	*	*	*	0.3	0.5	0.5	0.5	1.1	2.0	2.5	3.8
Female												
2002 .....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	1.6	2.5
2001 .....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	2.0	2.3
2000 .....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.6	1.1	1.9	2.1
1999 .....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	1.2	2.0	1.7
Non-Hispanic black												
Both sexes												
2002 .....	0.3	*	*	*	*	0.4	0.4	0.4	0.8	*	*	*
2001 .....	0.4	*	*	*	*	0.8	0.4	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	0.4	*	*	*	*	*
1999 .....	0.4	*	*	*	*	0.5	*	*	*	*	*	*
Male												
2002 .....	0.5	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	0.8	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hodgkin's lymphoma (C82-C85)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	7.6	*	*	0.1	0.3	0.7	1.5	3.9	11.5	28.8	57.2	77.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
All origins <sup>3</sup>												
Both sexes												
2001 .....	7.8	*	*	0.1	0.4	0.8	1.7	4.4	12.2	29.3	58.6	79.6
2000 .....	8.1	*	*	0.1	0.5	0.7	1.7	4.6	13.1	30.4	61.0	80.2
1999 .....	8.2	*	*	0.1	0.4	0.7	1.8	4.8	13.3	31.7	61.8	77.7
Male												
2002 .....	8.1	*	*	0.2	0.4	0.9	2.0	5.0	14.4	36.0	70.7	99.7
2001 .....	8.4	*	*	0.1	0.5	0.9	2.2	5.8	15.0	36.5	71.3	106.6
2000 .....	8.6	*	*	0.1	0.6	0.9	2.1	6.0	15.9	37.4	74.5	105.9
1999 .....	8.6	*	*	0.1	0.5	0.9	2.3	6.2	16.2	38.3	74.7	104.4
Female												
2002 .....	7.1	*	*	*	0.2	0.5	1.1	3.0	8.8	22.8	48.2	67.6
2001 .....	7.3	*	*	*	0.3	0.6	1.1	3.1	9.7	23.3	50.2	68.3
2000 .....	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5	24.7	52.1	69.8
1999 .....	7.7	*	*	*	0.3	0.6	1.2	3.4	10.6	26.4	53.4	66.9
Hispanic												
Both sexes												
2002 .....	2.8	*	*	*	0.4	0.6	1.3	3.0	9.4	22.7	43.7	50.4
2001 .....	2.7	*	*	*	0.3	0.5	1.4	3.6	10.9	18.3	43.3	59.4
2000 .....	2.9	*	*	*	0.3	0.6	1.6	3.6	11.1	25.7	41.5	59.7
1999 .....	2.8	*	*	*	*	0.6	1.4	4.4	9.3	25.2	42.7	56.1
Male												
2002 .....	2.7	*	*	*	*	0.6	1.5	3.6	11.7	26.0	45.5	60.2
2001 .....	3.0	*	*	*	*	0.6	2.0	4.4	13.2	24.4	58.6	68.6
2000 .....	3.1	*	*	*	*	0.7	1.9	4.6	13.9	30.2	48.2	82.6
1999 .....	2.8	*	*	*	*	*	1.7	5.7	11.8	27.5	48.0	60.4
Female												
2002 .....	2.8	*	*	*	*	*	1.2	2.3	7.4	20.1	42.4	45.6
2001 .....	2.4	*	*	*	*	*	0.8	2.8	8.8	13.5	32.9	54.9
2000 .....	2.7	*	*	*	*	*	1.3	2.6	8.6	22.2	37.0	48.5
1999 .....	2.8	*	*	*	*	*	1.0	3.1	7.1	23.4	39.2	54.1
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	8.3	*	*	0.1	0.3	0.7	1.5	4.0	11.6	29.2	57.7	78.2
2001 .....	8.6	*	*	0.1	0.4	0.8	1.7	4.5	12.3	29.9	59.2	80.3
2000 .....	8.8	*	*	0.1	0.5	0.8	1.7	4.7	13.2	30.7	61.7	80.9
1999 .....	8.9	*	*	0.1	0.4	0.7	1.8	4.8	13.5	32.1	62.4	78.3
Male												
2002 .....	9.0	*	*	0.2	0.4	0.9	2.0	5.1	14.6	36.5	71.8	101.4
2001 .....	9.2	*	*	0.1	0.5	0.9	2.3	5.9	15.1	37.2	71.7	108.1
2000 .....	9.4	*	*	0.2	0.7	0.9	2.2	6.1	15.9	37.7	75.6	106.7
1999 .....	9.4	*	*	0.2	0.5	0.9	2.3	6.2	16.5	38.8	75.5	106.0
Female												
2002 .....	7.7	*	*	*	0.3	0.5	1.1	3.0	8.9	23.0	48.3	68.2
2001 .....	8.0	*	*	*	0.3	0.6	1.2	3.1	9.7	23.8	50.9	68.7
2000 .....	8.3	*	*	*	0.3	0.6	1.2	3.4	10.6	24.8	52.6	70.4
1999 .....	8.4	*	*	*	0.3	0.5	1.3	3.4	10.8	26.5	53.9	67.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Non-Hodgkin's lymphoma (C82-C85)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	9.5	*	*	0.2	0.3	0.7	1.5	4.0	12.0	30.6	60.9	83.1
2001 .....	9.7	*	*	0.1	0.3	0.7	1.7	4.5	12.7	31.4	62.3	84.1
2000 .....	10.1	*	*	0.1	0.4	0.7	1.7	4.8	13.8	32.3	65.1	85.1
1999 .....	10.1	*	*	0.1	0.4	0.7	1.7	4.9	14.1	33.9	65.6	82.0
<b>Male</b>												
2002 .....	10.2	*	*	0.2	0.4	0.8	1.9	5.1	15.0	38.3	75.6	107.6
2001 .....	10.4	*	*	*	0.4	0.9	2.1	6.0	15.5	38.8	75.2	113.1
2000 .....	10.6	*	*	0.2	0.6	0.8	2.2	6.1	16.5	39.5	79.6	113.0
1999 .....	10.6	*	*	0.2	0.5	0.8	2.2	6.3	17.0	40.8	78.9	110.0
<b>Female</b>												
2002 .....	8.8	*	*	*	0.2	0.5	1.1	2.9	9.2	24.1	51.0	72.6
2001 .....	9.2	*	*	*	0.3	0.5	1.2	3.1	10.1	25.2	53.7	72.0
2000 .....	9.5	*	*	*	0.3	0.6	1.1	3.4	11.3	26.3	55.6	73.8
1999 .....	9.6	*	*	*	0.2	0.5	1.2	3.5	11.4	28.1	56.9	70.8
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	4.0	*	*	*	0.4	1.2	2.0	4.8	10.8	21.6	30.1	31.2
2001 .....	4.1	*	*	*	0.6	1.2	2.3	5.1	10.9	21.3	30.6	40.0
2000 .....	3.9	*	*	*	0.6	1.1	2.1	5.0	10.8	19.7	27.1	36.6
1999 .....	4.1	*	*	*	0.6	1.1	2.4	4.8	11.2	20.5	32.1	40.4
<b>Male</b>												
2002 .....	4.5	*	*	*	*	1.6	2.9	5.7	13.9	28.3	35.8	43.8
2001 .....	4.7	*	*	*	0.8	1.3	3.4	6.7	13.7	28.2	37.2	55.4
2000 .....	4.3	*	*	*	*	1.2	2.6	7.0	14.8	25.3	31.8	43.4
1999 .....	4.7	*	*	*	*	1.6	3.3	6.6	14.8	26.2	41.0	59.4
<b>Female</b>												
2002 .....	3.6	*	*	*	*	0.8	1.2	4.0	8.3	16.9	26.9	26.3
2001 .....	3.7	*	*	*	*	1.1	1.4	3.7	8.6	16.5	27.0	34.2
2000 .....	3.5	*	*	*	*	0.9	1.7	3.3	7.5	15.7	24.5	34.0
1999 .....	3.6	*	*	*	*	*	1.7	3.2	8.2	16.5	27.1	33.3
<b>Leukemia (C91-C95)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	7.5	0.5	0.7	0.8	1.2	1.2	1.7	3.6	9.8	26.4	53.1	80.5
2001 .....	7.5	0.6	1.0	0.8	1.2	1.3	1.7	3.8	10.0	26.7	53.3	81.4
2000 .....	7.6	1.0	0.9	0.8	1.2	1.2	1.8	3.7	9.9	27.2	54.5	83.6
1999 .....	7.5	0.7	0.8	0.7	1.2	1.2	1.8	3.7	10.6	27.2	52.8	82.4
<b>Male</b>												
2002 .....	8.5	*	0.7	0.8	1.4	1.4	1.9	4.2	12.2	36.3	74.9	114.5
2001 .....	8.5	*	1.0	0.9	1.4	1.5	1.9	4.2	12.8	35.7	74.2	117.5
2000 .....	8.5	1.1	0.9	0.9	1.5	1.4	1.9	4.3	12.5	37.2	75.7	119.8
1999 .....	8.4	*	0.8	0.8	1.5	1.4	1.9	4.3	13.1	36.8	73.2	119.2
<b>Female</b>												
2002 .....	6.5	*	0.6	0.7	1.0	1.0	1.5	3.1	7.6	18.1	38.7	65.7
2001 .....	6.6	*	1.0	0.7	1.0	1.1	1.5	3.3	7.5	19.2	39.5	66.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
All origins <sup>3</sup>												
Female												
2000 .....	6.7	*	0.9	0.6	0.9	1.0	1.7	3.2	7.6	19.0	40.6	68.8
1999 .....	6.7	*	0.7	0.7	1.0	1.1	1.6	3.2	8.3	19.3	39.6	67.6
Hispanic												
Both sexes												
2002 .....	2.8	*	0.8	1.1	1.6	1.1	2.1	2.5	7.2	15.5	27.4	47.1
2001 .....	3.0	*	1.3	1.1	1.7	1.5	1.9	2.9	6.5	17.3	33.4	42.5
2000 .....	2.9	*	0.9	1.1	1.5	1.1	1.9	3.5	7.8	16.9	32.6	49.1
1999 .....	2.9	*	0.9	1.1	1.6	1.4	2.0	3.4	7.0	17.3	31.0	44.2
Male												
2002 .....	3.0	*	*	1.1	2.0	1.4	2.3	2.8	7.2	18.5	38.6	65.3
2001 .....	3.4	*	1.4	1.1	2.2	1.8	2.3	3.3	8.4	19.6	45.6	68.6
2000 .....	3.3	*	*	1.3	1.7	1.5	2.0	4.1	10.2	20.7	42.8	76.6
1999 .....	3.1	*	*	1.1	1.7	1.4	1.9	3.6	8.6	23.5	39.1	49.6
Female												
2002 .....	2.6	*	*	1.2	1.1	0.7	1.9	2.3	7.3	13.1	19.7	38.1
2001 .....	2.6	*	*	1.1	1.1	1.1	1.4	2.6	4.8	15.6	25.1	29.8
2000 .....	2.6	*	*	0.8	1.3	*	1.8	2.9	5.6	13.9	25.8	35.6
1999 .....	2.8	*	*	1.0	1.5	1.3	2.1	3.2	5.6	12.4	25.7	41.6
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	8.2	*	0.6	0.7	1.1	1.2	1.6	3.7	10.0	27.0	54.3	81.7
2001 .....	8.2	*	0.9	0.7	1.1	1.2	1.7	3.8	10.3	27.2	54.1	82.7
2000 .....	8.2	1.0	0.9	0.7	1.1	1.2	1.7	3.8	10.1	27.8	55.3	84.7
1999 .....	8.2	*	0.7	0.7	1.2	1.2	1.7	3.8	10.8	27.8	53.6	83.7
Male												
2002 .....	9.4	*	0.7	0.8	1.2	1.4	1.9	4.3	12.6	37.3	76.5	116.5
2001 .....	9.3	*	0.9	0.8	1.2	1.4	1.9	4.3	13.1	36.6	75.4	119.3
2000 .....	9.3	*	0.9	0.8	1.4	1.4	1.9	4.3	12.6	38.2	77.0	121.5
1999 .....	9.2	*	0.8	0.8	1.4	1.3	1.9	4.4	13.4	37.5	74.6	122.1
Female												
2002 .....	7.0	*	0.5	0.6	0.9	1.0	1.4	3.2	7.6	18.4	39.5	66.6
2001 .....	7.2	*	0.9	0.6	0.9	1.1	1.5	3.4	7.7	19.4	40.1	67.5
2000 .....	7.2	*	0.9	0.6	0.8	1.0	1.6	3.3	7.7	19.3	41.1	69.8
1999 .....	7.2	*	0.7	0.6	0.9	1.1	1.5	3.2	8.5	19.8	40.0	68.3
Non-Hispanic white												
Both sexes												
2002 .....	9.1	*	0.7	0.7	1.1	1.2	1.7	3.7	10.0	28.2	56.2	84.4
2001 .....	9.1	*	1.0	0.7	1.1	1.2	1.7	3.8	10.5	28.1	56.0	86.1
2000 .....	9.2	1.3	0.9	0.7	1.1	1.1	1.7	3.7	10.2	29.0	57.5	87.3
1999 .....	9.1	*	0.8	0.7	1.1	1.2	1.7	3.7	11.0	28.6	55.3	86.8
Male												
2002 .....	10.5	*	0.8	0.7	1.3	1.3	1.9	4.2	12.7	38.7	79.2	121.3
2001 .....	10.3	*	1.0	0.7	1.3	1.4	1.8	4.3	13.3	37.7	78.0	125.3
2000 .....	10.4	*	1.0	0.8	1.3	1.3	1.9	4.2	12.7	39.5	79.9	125.5
1999 .....	10.2	*	0.8	0.7	1.4	1.3	1.9	4.3	13.6	38.6	77.0	127.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Leukemia (C91-C95)</b>												
<b>Non-Hispanic white</b>												
Female												
2002 .....	7.8	*	0.5	0.6	0.9	1.0	1.5	3.2	7.5	19.3	40.7	68.4
2001 .....	8.0	*	1.1	0.6	0.9	1.0	1.6	3.3	7.8	19.9	41.4	69.8
2000 .....	8.1	*	0.9	0.6	0.8	0.9	1.5	3.2	7.8	20.1	42.7	72.0
1999 .....	8.0	*	0.9	0.6	0.8	1.1	1.5	3.2	8.5	20.3	41.1	70.8
<b>Non-Hispanic black</b>												
Both sexes												
2002 .....	4.8	*	*	0.6	0.9	1.3	1.6	4.4	11.5	22.6	42.9	65.3
2001 .....	4.9	*	*	0.9	1.0	1.5	2.1	4.3	10.1	25.0	42.1	57.1
2000 .....	4.9	*	*	0.6	1.4	1.7	2.1	5.0	10.8	23.1	38.8	66.1
1999 .....	4.9	*	*	0.8	1.5	1.3	1.9	4.4	10.8	25.2	41.6	55.9
Male												
2002 .....	5.3	*	*	0.7	1.0	1.5	1.8	5.2	14.1	32.2	58.0	87.5
2001 .....	5.4	*	*	1.2	1.1	1.8	2.6	4.6	13.4	34.4	56.1	75.1
2000 .....	5.5	*	*	0.7	1.9	2.0	2.0	5.6	14.0	32.3	55.8	98.6
1999 .....	5.4	*	*	1.0	1.6	1.5	2.1	5.7	13.7	33.4	55.7	78.4
Female												
2002 .....	4.4	*	*	*	0.8	1.1	1.4	3.6	9.4	15.8	34.3	56.9
2001 .....	4.4	*	*	*	1.0	1.1	1.7	4.0	7.4	18.4	34.2	50.5
2000 .....	4.4	*	*	*	0.9	1.4	2.2	4.4	8.3	16.5	29.2	54.1
1999 .....	4.4	*	*	0.6	1.3	1.1	1.7	3.3	8.5	19.4	33.8	47.5
<b>Multiple myeloma and immunoproliferative neoplasms (C88,C90)</b>												
<b>All origins <sup>3</sup></b>												
Both sexes												
2002 .....	4.0	*	*	*	*	*	0.3	1.8	6.3	17.4	31.2	36.7
2001 .....	3.9	*	*	*	*	*	0.3	1.7	6.6	17.4	30.9	36.3
2000 .....	3.9	*	*	*	*	*	0.3	1.9	6.3	17.3	31.5	37.8
1999 .....	3.9	*	*	*	*	0.1	0.3	1.8	7.1	17.2	31.0	36.6
Male												
2002 .....	4.2	*	*	*	*	*	0.3	2.2	7.6	21.0	39.7	50.8
2001 .....	4.1	*	*	*	*	*	0.4	2.1	7.8	20.8	38.6	48.0
2000 .....	4.0	*	*	*	*	*	0.4	2.1	6.9	20.4	39.6	51.8
1999 .....	4.1	*	*	*	*	*	0.4	2.1	8.5	20.1	38.3	51.3
Female												
2002 .....	3.7	*	*	*	*	*	0.3	1.5	5.1	14.3	25.5	30.6
2001 .....	3.8	*	*	*	*	*	0.3	1.3	5.5	14.5	25.8	31.3
2000 .....	3.8	*	*	*	*	*	0.3	1.6	5.7	14.7	26.3	32.2
1999 .....	3.8	*	*	*	*	*	0.3	1.5	5.9	14.8	26.2	30.6
<b>Hispanic</b>												
Both sexes												
2002 .....	1.5	*	*	*	*	*	*	1.8	5.0	14.3	26.7	33.3
2001 .....	1.4	*	*	*	*	*	*	1.3	5.8	14.9	24.7	29.4
2000 .....	1.4	*	*	*	*	*	*	1.4	5.3	15.0	24.5	31.2
1999 .....	1.4	*	*	*	*	*	*	1.7	6.8	13.4	23.1	28.1
Male												
2002 .....	1.6	*	*	*	*	*	*	2.1	6.6	17.6	31.6	53.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Hispanic												
Male												
2001 .....	1.5	*	*	*	*	*	*	1.4	6.0	20.4	35.8	38.1
2000 .....	1.3	*	*	*	*	*	*	1.5	5.5	18.4	30.0	*
1999 .....	1.3	*	*	*	*	*	*	1.9	7.8	14.7	24.5	*
Female												
2002 .....	1.4	*	*	*	*	*	*	1.5	3.5	11.7	23.3	23.2
2001 .....	1.3	*	*	*	*	*	*	1.2	5.6	10.6	17.2	25.1
2000 .....	1.4	*	*	*	*	*	*	*	5.1	12.3	20.8	30.7
1999 .....	1.5	*	*	*	*	*	*	1.5	5.9	12.4	22.2	31.2
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	4.3	*	*	*	*	*	0.3	1.8	6.4	17.5	31.3	36.7
2001 .....	4.3	*	*	*	*	*	0.3	1.7	6.6	17.5	31.1	36.5
2000 .....	4.3	*	*	*	*	*	0.3	1.9	6.4	17.4	31.8	38.0
1999 .....	4.3	*	*	*	*	*	0.4	1.8	7.1	17.4	31.2	36.8
Male												
2002 .....	4.6	*	*	*	*	*	0.4	2.2	7.6	21.2	40.0	50.6
2001 .....	4.5	*	*	*	*	*	0.4	2.2	7.9	20.8	38.6	48.4
2000 .....	4.4	*	*	*	*	*	0.4	2.2	7.0	20.5	39.9	52.5
1999 .....	4.5	*	*	*	*	*	0.4	2.1	8.5	20.4	38.9	52.2
Female												
2002 .....	4.0	*	*	*	*	*	0.3	1.5	5.2	14.5	25.5	30.8
2001 .....	4.1	*	*	*	*	*	0.3	1.4	5.5	14.8	26.1	31.5
2000 .....	4.2	*	*	*	*	*	0.3	1.6	5.7	14.8	26.5	32.1
1999 .....	4.1	*	*	*	*	*	0.3	1.5	5.8	15.0	26.3	30.6
Non-Hispanic white												
Both sexes												
2002 .....	4.5	*	*	*	*	*	0.3	1.5	5.6	16.4	30.3	35.7
2001 .....	4.4	*	*	*	*	*	0.3	1.5	5.7	16.3	29.7	35.1
2000 .....	4.3	*	*	*	*	*	0.3	1.6	5.6	16.2	30.2	35.2
1999 .....	4.3	*	*	*	*	*	0.3	1.5	6.1	16.2	29.6	35.0
Male												
2002 .....	4.8	*	*	*	*	*	0.3	1.9	6.9	20.2	39.2	50.1
2001 .....	4.6	*	*	*	*	*	0.3	1.9	6.9	19.5	37.4	47.3
2000 .....	4.6	*	*	*	*	*	0.4	1.8	6.3	19.2	38.5	50.9
1999 .....	4.6	*	*	*	*	*	0.3	1.9	7.5	19.2	37.4	50.6
Female												
2002 .....	4.1	*	*	*	*	*	0.2	1.1	4.5	13.1	24.4	29.5
2001 .....	4.1	*	*	*	*	*	0.2	1.1	4.7	13.6	24.6	30.0
2000 .....	4.1	*	*	*	*	*	0.2	1.4	4.9	13.6	24.8	28.9
1999 .....	4.1	*	*	*	*	*	0.2	1.2	4.7	13.7	24.6	28.7
Non-Hispanic black												
Both sexes												
2002 .....	4.8	*	*	*	*	*	0.7	4.3	13.4	31.8	49.2	55.7
2001 .....	5.0	*	*	*	*	*	0.9	3.7	15.3	31.8	53.2	59.3
2000 .....	5.2	*	*	*	*	*	0.8	4.5	14.1	32.5	55.6	72.8
1999 .....	5.1	*	*	*	*	*	0.9	4.3	16.8	30.7	53.6	63.4

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Non-Hispanic black												
Male												
2002 .....	4.8	*	*	*	*	*	0.8	4.8	15.9	37.6	60.1	71.1
2001 .....	5.0	*	*	*	*	*	1.1	4.3	18.5	37.2	66.0	72.8
2000 .....	5.0	*	*	*	*	*	*	5.0	15.4	39.1	68.5	81.0
1999 .....	5.0	*	*	*	*	*	1.1	4.4	19.0	35.6	65.0	80.8
Female												
2002 .....	4.9	*	*	*	*	*	*	3.8	11.3	27.7	43.1	49.7
2001 .....	5.0	*	*	*	*	*	0.8	3.1	12.7	28.0	46.0	54.3
2000 .....	5.4	*	*	*	*	*	0.9	4.1	13.1	27.9	48.3	69.8
1999 .....	5.3	*	*	*	*	*	0.7	4.3	15.1	27.2	47.3	56.9
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001 .....	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
2000 .....	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
1999 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Male												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001 .....	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
2000 .....	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
1999 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Male												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Male												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
Non-Hispanic black												
Female												
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	21.8	0.6	0.8	0.7	1.3	1.8	4.5	13.7	36.6	80.8	146.7	230.1
2001 .....	22.0	*	0.8	0.7	1.3	1.8	4.6	14.1	37.3	82.0	148.7	234.5
2000 .....	22.4	0.6	0.8	0.6	1.3	1.8	4.6	14.1	38.3	85.5	152.6	239.8
1999 .....	23.7	*	0.8	0.7	1.3	1.9	5.1	15.0	41.5	91.4	160.8	250.4
Male												
2002 .....	22.7	*	0.8	0.7	1.6	2.3	5.0	15.9	43.2	98.1	180.0	290.8
2001 .....	22.6	*	0.9	0.7	1.7	2.2	4.9	16.4	43.8	98.5	181.1	295.2
2000 .....	23.0	*	1.0	0.7	1.6	2.1	5.1	16.0	44.9	103.2	187.8	300.7
1999 .....	24.5	*	0.8	0.8	1.7	2.4	5.8	17.3	49.1	110.4	196.1	318.9
Female												
2002 .....	21.0	*	0.8	0.7	0.9	1.4	4.1	11.6	30.4	66.3	124.6	203.8
2001 .....	21.4	*	0.6	0.6	0.9	1.5	4.2	11.7	31.3	68.4	127.4	209.1
2000 .....	21.9	*	0.7	0.6	1.0	1.4	4.2	12.2	32.2	70.8	129.6	215.0
1999 .....	23.1	*	0.7	0.7	1.0	1.4	4.5	12.8	34.5	75.8	137.9	222.8
Hispanic												
Both sexes												
2002 .....	7.8	*	*	0.8	1.6	1.5	3.0	9.5	25.4	60.3	112.8	164.1
2001 .....	7.7	*	*	0.6	1.3	1.2	3.4	9.8	26.0	60.7	110.9	193.8
2000 .....	7.8	*	0.8	0.7	1.2	1.5	2.7	9.2	26.6	64.1	124.6	192.4
1999 .....	8.1	*	*	0.7	1.3	1.6	3.5	10.6	28.8	64.8	122.3	202.1
Male												
2002 .....	7.8	*	*	0.8	1.9	1.7	3.1	10.4	29.6	69.2	138.7	204.1
2001 .....	7.7	*	*	0.6	1.8	1.5	3.4	10.7	30.2	67.7	140.9	213.3
2000 .....	7.7	*	*	0.7	1.4	1.9	2.9	9.5	29.2	73.6	151.9	231.8
1999 .....	8.0	*	*	0.7	1.8	1.9	3.5	11.1	33.3	77.1	144.9	235.1
Female												
2002 .....	7.7	*	*	0.7	1.2	1.1	2.9	8.6	21.6	53.2	95.0	144.2
2001 .....	7.7	*	*	0.6	0.7	0.8	3.5	8.9	22.4	55.3	90.7	184.3
2000 .....	7.9	*	*	0.6	1.1	1.1	2.5	9.0	24.3	56.6	106.3	173.1
1999 .....	8.2	*	*	0.7	0.8	1.2	3.6	10.1	24.7	55.1	107.2	186.1
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	24.0	*	0.9	0.7	1.2	1.9	4.8	14.1	37.3	81.9	148.0	232.1
2001 .....	24.1	*	0.8	0.7	1.3	2.0	4.7	14.4	38.1	83.1	150.1	235.3
2000 .....	24.4	0.7	0.8	0.6	1.3	1.8	4.8	14.5	39.1	86.5	153.3	241.0
1999 .....	25.8	*	0.8	0.7	1.3	1.9	5.3	15.3	42.3	92.7	161.9	251.6

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
Non-Hispanic <sup>4</sup>												
Male												
2002 .....	25.0	*	0.9	0.7	1.6	2.4	5.3	16.4	44.1	99.7	181.5	294.0
2001 .....	24.9	*	1.0	0.8	1.7	2.4	5.1	16.9	44.7	100.0	182.7	297.9
2000 .....	25.2	*	0.9	0.6	1.6	2.2	5.3	16.5	45.9	104.7	188.7	303.0
1999 .....	26.8	*	1.0	0.8	1.6	2.5	6.1	17.8	50.1	112.0	197.6	321.9
Female												
2002 .....	22.9	*	0.8	0.7	0.9	1.4	4.3	11.8	31.0	67.1	125.8	205.4
2001 .....	23.3	*	0.6	0.6	0.9	1.6	4.3	12.0	31.9	69.1	128.7	209.3
2000 .....	23.7	*	0.8	0.6	1.0	1.4	4.4	12.5	32.8	71.5	130.2	215.9
1999 .....	25.0	*	0.7	0.6	1.1	1.4	4.6	13.0	35.1	76.9	138.8	223.5
Non-Hispanic white												
Both sexes												
2002 .....	26.0	*	0.8	0.7	1.2	1.9	4.6	13.5	36.5	81.8	149.1	235.2
2001 .....	26.0	*	0.9	0.7	1.3	2.0	4.5	13.6	36.8	82.5	150.8	238.9
2000 .....	26.5	*	0.8	0.7	1.3	1.9	4.5	13.6	38.2	86.3	153.9	244.9
1999 .....	27.9	*	0.9	0.7	1.3	2.0	5.0	14.6	40.9	92.7	162.3	253.6
Male												
2002 .....	27.2	*	0.8	0.7	1.5	2.4	5.0	15.7	42.9	100.0	183.2	300.2
2001 .....	26.8	*	1.1	0.7	1.7	2.3	5.0	15.9	42.8	99.3	183.7	302.9
2000 .....	27.2	*	1.0	0.7	1.6	2.3	5.0	15.6	44.1	104.6	189.1	309.5
1999 .....	28.8	*	1.1	0.8	1.5	2.6	5.8	16.6	48.3	111.5	198.9	326.9
Female												
2002 .....	24.9	*	0.7	0.7	0.9	1.4	4.3	11.3	30.5	66.3	126.3	207.0
2001 .....	25.3	*	0.7	0.6	1.0	1.6	4.0	11.4	31.3	68.3	129.1	212.3
2000 .....	25.8	*	0.6	0.6	1.0	1.4	4.1	11.7	32.6	71.0	130.8	218.9
1999 .....	27.0	*	0.8	0.6	1.1	1.3	4.2	12.6	33.9	77.0	138.3	224.3
Non-Hispanic black												
Both sexes												
2002 .....	18.7	*	1.1	0.8	1.3	2.4	6.5	20.5	51.8	97.9	159.6	223.2
2001 .....	18.9	*	*	0.7	1.3	2.4	6.1	22.2	54.0	101.8	158.4	218.5
2000 .....	18.9	*	0.9	0.6	1.5	1.7	7.7	22.6	52.8	101.2	163.6	211.1
1999 .....	20.2	*	*	0.8	1.5	2.0	7.6	23.6	60.6	105.9	172.7	244.8
Male												
2002 .....	19.4	*	*	0.9	1.8	2.7	7.6	24.5	64.7	119.0	197.8	265.8
2001 .....	20.0	*	*	0.8	1.8	2.8	6.3	27.1	69.0	128.4	195.0	287.6
2000 .....	20.0	*	*	0.7	2.0	1.9	8.7	26.9	69.1	123.8	212.0	253.6
1999 .....	21.3	*	*	0.7	2.0	2.4	8.5	29.7	77.5	134.5	208.6	293.6
Female												
2002 .....	18.0	*	*	0.8	0.8	2.1	5.5	17.1	41.4	83.0	137.9	206.9
2001 .....	17.8	*	*	0.6	0.9	2.0	5.8	18.0	42.0	83.0	137.8	192.9
2000 .....	17.8	*	*	*	1.0	1.5	6.8	18.9	39.6	85.2	136.3	195.4
1999 .....	19.2	*	*	0.9	1.0	1.7	6.9	18.3	47.0	85.8	152.7	226.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	4.6	1.7	0.4	0.2	0.2	0.4	0.8	1.8	4.5	12.8	36.1	78.7
2001 .....	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.8	5.0	14.1	37.0	82.7
2000 .....	4.8	1.6	0.3	0.2	0.2	0.4	0.8	1.8	5.0	13.8	37.6	84.9
1999 .....	4.7	1.5	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.6	36.6	82.9
Male												
2002 .....	4.6	1.5	0.4	0.2	0.2	0.4	0.9	1.9	5.3	15.8	45.9	101.1
2001 .....	4.8	1.3	0.4	0.2	0.3	0.4	0.9	2.0	5.5	17.3	48.2	107.6
2000 .....	4.8	1.6	0.4	0.2	0.3	0.4	0.8	2.1	5.6	17.0	47.5	108.9
1999 .....	4.8	1.5	0.4	0.2	0.3	0.5	0.9	2.0	5.8	18.4	46.1	109.2
Female												
2002 .....	4.6	2.0	0.4	0.2	0.2	0.4	0.8	1.7	3.7	10.4	29.6	68.9
2001 .....	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.6	4.4	11.3	29.7	72.3
2000 .....	4.9	1.5	0.3	0.2	0.2	0.4	0.8	1.6	4.5	11.2	31.1	75.2
1999 .....	4.7	1.5	0.4	0.3	0.2	0.4	0.6	1.6	3.9	11.5	30.4	72.3
Hispanic												
Both sexes												
2002 .....	1.5	*	*	0.4	*	*	0.6	1.5	3.0	8.9	24.7	48.8
2001 .....	1.5	*	*	0.3	*	0.3	0.7	1.5	3.6	8.8	23.1	57.5
2000 .....	1.5	*	*	*	*	0.4	0.7	1.4	3.1	9.1	24.3	70.3
1999 .....	1.5	*	*	*	*	0.4	0.6	1.3	3.7	10.5	23.1	51.9
Male												
2002 .....	1.4	*	*	*	*	*	*	1.4	3.7	10.3	31.6	56.9
2001 .....	1.5	*	*	*	*	*	0.8	1.8	4.0	10.0	30.0	70.5
2000 .....	1.3	*	*	*	*	*	*	*	3.1	9.7	26.6	86.7
1999 .....	1.3	*	*	*	*	*	0.8	1.8	3.9	10.3	24.5	49.6
Female												
2002 .....	1.5	*	*	*	*	*	*	1.5	2.4	7.8	20.0	44.8
2001 .....	1.5	*	*	*	*	*	*	1.2	3.3	7.9	18.4	51.2
2000 .....	1.7	*	*	*	*	*	0.9	1.6	3.1	8.6	22.8	62.3
1999 .....	1.6	*	*	*	*	*	*	*	3.5	10.7	22.2	53.0
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	5.1	1.6	0.4	0.2	0.2	0.5	0.8	1.8	4.6	13.1	36.6	79.8
2001 .....	5.3	1.4	0.4	0.2	0.3	0.4	0.8	1.9	5.0	14.4	37.5	83.5
2000 .....	5.3	1.7	0.4	0.2	0.3	0.4	0.8	1.9	5.2	14.1	38.1	85.2
1999 .....	5.2	1.6	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.8	37.0	83.8
Male												
2002 .....	5.1	1.3	0.4	0.2	0.3	0.5	0.9	1.9	5.4	16.2	46.4	102.9
2001 .....	5.3	1.5	0.4	0.2	0.3	0.4	0.9	2.1	5.6	17.8	48.9	109.0
2000 .....	5.3	1.8	0.5	0.2	0.3	0.4	0.9	2.2	5.8	17.5	48.3	109.4
1999 .....	5.3	1.6	0.5	0.2	0.3	0.5	0.9	2.0	5.9	18.8	46.9	111.4
Female												
2002 .....	5.1	2.0	0.4	0.2	0.2	0.4	0.8	1.7	3.8	10.5	30.0	69.8
2001 .....	5.2	1.4	0.4	0.3	0.3	0.4	0.8	1.7	4.5	11.5	30.1	72.9
2000 .....	5.3	1.7	*	0.3	0.2	0.4	0.8	1.6	4.6	11.4	31.4	75.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
Non-Hispanic <sup>4</sup>												
Female												
1999 .....	5.1	1.7	0.4	0.3	0.2	0.4	0.7	1.7	3.9	11.5	30.6	72.8
Non-Hispanic white												
Both sexes												
2002 .....	5.7	1.8	0.4	0.2	0.2	0.4	0.8	1.8	4.6	13.4	37.5	81.8
2001 .....	5.9	1.3	0.4	0.2	0.3	0.4	0.8	1.9	5.0	14.7	38.6	84.8
2000 .....	5.9	1.8	0.4	0.2	0.3	0.4	0.8	1.8	5.2	14.2	39.1	87.6
1999 .....	5.7	1.5	0.4	0.3	0.2	0.4	0.7	1.8	4.8	15.0	37.8	86.3
Male												
2002 .....	5.8	*	*	0.2	0.2	0.5	1.0	1.9	5.4	16.7	47.6	106.0
2001 .....	6.0	*	*	0.2	0.2	0.5	0.9	2.1	5.7	18.3	50.1	111.4
2000 .....	5.9	1.8	0.5	0.2	0.3	0.4	0.8	2.1	5.8	17.4	49.9	113.1
1999 .....	5.9	*	*	0.2	0.2	0.4	0.9	2.0	5.9	19.2	48.1	116.8
Female												
2002 .....	5.6	2.1	*	*	0.2	0.3	0.7	1.6	3.9	10.7	30.7	71.4
2001 .....	5.8	*	*	0.3	0.3	0.4	0.7	1.7	4.5	11.6	30.9	73.7
2000 .....	5.9	1.8	*	0.3	0.2	0.4	0.8	1.5	4.7	11.5	32.1	77.3
1999 .....	5.6	*	*	0.3	0.2	0.4	0.6	1.5	3.7	11.5	31.1	74.1
Non-Hispanic black												
Both sexes												
2002 .....	3.1	*	*	*	*	0.6	1.2	2.5	5.6	12.8	31.8	64.4
2001 .....	3.2	*	*	0.4	0.4	*	1.1	2.1	6.0	14.5	31.9	72.7
2000 .....	3.2	*	*	*	*	0.6	1.1	2.7	5.6	15.1	31.3	64.2
1999 .....	3.3	*	*	*	0.4	0.6	1.2	2.6	6.4	15.9	33.1	62.7
Male												
2002 .....	2.9	*	*	*	*	*	1.0	2.2	7.3	14.3	39.3	81.0
2001 .....	3.1	*	*	*	*	*	1.1	1.9	6.6	17.6	42.3	87.8
2000 .....	3.1	*	*	*	*	*	1.2	2.8	7.0	19.5	36.4	76.3
1999 .....	3.1	*	*	*	*	0.8	1.2	2.4	7.3	19.2	38.8	64.2
Female												
2002 .....	3.4	*	*	*	*	0.8	1.4	2.7	4.3	11.7	27.5	58.1
2001 .....	3.3	*	*	*	*	*	1.0	2.2	5.4	12.2	26.1	67.2
2000 .....	3.3	*	*	*	*	0.8	1.0	2.6	4.5	12.1	28.5	59.7
1999 .....	3.5	*	*	*	*	*	1.2	2.8	5.7	13.6	29.9	62.2
Anemias (D50-D64)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.6	*	0.1	0.1	0.3	0.5	0.4	0.6	1.0	2.8	9.1	40.3
2001 .....	1.6	*	0.1	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.0
2000 .....	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1	2.8	9.5	42.4
1999 .....	1.6	*	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	9.5	43.8
Male												
2002 .....	1.3	*	*	0.1	0.3	0.5	0.4	0.6	1.0	2.9	8.9	38.3
2001 .....	1.3	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	9.9	40.0
2000 .....	1.2	*	*	0.1	0.2	0.3	0.3	0.6	1.1	2.8	9.5	40.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Anemias (D50-D64)</b>												
<b>All origins <sup>3</sup></b>												
Male												
1999 .....	1.3	*	*	0.1	0.3	0.4	0.5	0.6	0.9	2.7	10.1	43.4
Female												
2002 .....	1.9	*	*	0.1	0.2	0.4	0.4	0.7	1.0	2.6	9.2	41.2
2001 .....	2.0	*	*	0.1	0.3	0.3	0.5	0.6	1.0	2.5	9.6	42.9
2000 .....	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0	2.8	9.4	43.3
1999 .....	1.9	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.2	44.0
<b>Hispanic</b>												
Both sexes												
2002 .....	0.5	*	*	*	*	*	*	*	*	3.1	7.3	34.4
2001 .....	0.5	*	*	*	*	*	*	*	*	2.2	7.8	28.8
2000 .....	0.4	*	*	*	*	*	*	*	*	2.2	7.1	17.3
1999 .....	0.5	*	*	*	*	*	*	*	*	2.6	7.3	35.1
Male												
2002 .....	0.5	*	*	*	*	*	*	*	*	*	*	53.5
2001 .....	0.4	*	*	*	*	*	*	*	*	*	9.4	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.5	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.5	*	*	*	*	*	*	*	*	3.0	7.3	24.9
2001 .....	0.6	*	*	*	*	*	*	*	*	*	6.6	32.6
2000 .....	0.5	*	*	*	*	*	*	*	*	*	6.9	*
1999 .....	0.5	*	*	*	*	*	*	*	*	*	*	32.2
<b>Non-Hispanic <sup>4</sup></b>												
Both sexes												
2002 .....	1.8	*	*	0.1	0.3	0.5	0.5	0.7	1.0	2.7	9.2	40.5
2001 .....	1.8	*	*	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.5
2000 .....	1.8	*	0.3	0.1	0.2	0.4	0.5	0.6	1.1	2.8	9.5	43.2
1999 .....	1.8	*	0.2	0.1	0.4	0.4	0.5	0.7	1.0	2.6	9.6	44.0
Male												
2002 .....	1.4	*	*	0.1	0.4	0.6	0.4	0.6	1.0	2.9	9.0	37.6
2001 .....	1.4	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	10.0	40.8
2000 .....	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	2.8	9.6	41.0
1999 .....	1.4	*	*	0.1	0.4	0.5	0.5	0.7	1.0	2.6	10.1	43.3
Female												
2002 .....	2.1	*	*	0.2	0.2	0.4	0.5	0.7	1.0	2.6	9.3	41.8
2001 .....	2.2	*	*	0.2	0.3	0.4	0.6	0.6	1.0	2.6	9.7	43.1
2000 .....	2.2	*	*	*	0.3	0.4	0.5	0.6	1.0	2.8	9.5	44.1
1999 .....	2.1	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.3	44.2
<b>Non-Hispanic white</b>												
Both sexes												
2002 .....	1.6	*	*	*	0.1	0.1	0.1	0.3	0.7	2.4	8.7	39.9
2001 .....	1.7	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.4	42.1
2000 .....	1.7	*	*	*	*	*	0.1	0.3	0.8	2.5	9.3	42.9
1999 .....	1.6	*	*	*	0.1	*	0.1	0.3	0.6	2.4	9.4	43.2
Male												
2002 .....	1.2	*	*	*	*	0.2	*	0.3	0.7	2.5	8.5	36.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Anemias (D50-D64)</b>												
Non-Hispanic white												
Male												
2001 .....	1.3	*	*	*	*	*	*	0.3	0.7	2.6	9.8	41.2
2000 .....	1.2	*	*	*	*	*	0.1	0.3	0.9	2.7	9.3	39.9
1999 .....	1.2	*	*	*	*	*	*	0.3	0.5	2.4	9.9	43.4
Female												
2002 .....	2.1	*	*	*	*	*	*	0.3	0.7	2.4	8.8	41.4
2001 .....	2.1	*	*	*	*	*	0.2	0.3	0.8	2.4	9.2	42.4
2000 .....	2.1	*	*	*	*	*	*	0.3	0.7	2.4	9.4	44.1
1999 .....	2.0	*	*	*	*	*	*	0.3	0.7	2.4	9.1	43.1
Non-Hispanic black												
Both sexes												
2002 .....	2.9	*	*	0.4	1.3	2.5	2.5	3.1	3.7	6.1	16.2	52.9
2001 .....	2.8	*	*	0.4	1.2	2.3	2.8	3.1	3.0	5.5	16.1	49.3
2000 .....	2.7	*	*	0.5	1.1	2.4	2.5	3.1	3.5	5.5	13.4	51.8
1999 .....	2.9	*	0.9	0.5	1.5	2.7	2.6	3.2	4.3	5.0	13.4	53.0
Male												
2002 .....	2.7	*	*	*	1.7	2.9	2.5	2.6	4.0	7.5	17.0	58.0
2001 .....	2.6	*	*	*	1.4	3.0	2.5	3.6	3.2	6.8	15.3	42.7
2000 .....	2.4	*	*	*	1.0	2.2	2.1	3.4	3.5	4.6	13.9	56.4
1999 .....	2.7	*	*	*	1.8	2.9	2.8	3.4	5.1	5.6	12.8	44.0
Female												
2002 .....	3.0	*	*	*	1.0	2.0	2.6	3.5	3.4	5.1	15.8	51.0
2001 .....	2.9	*	*	*	1.1	1.6	3.0	2.7	2.9	4.6	16.5	51.8
2000 .....	3.1	*	*	*	1.1	2.5	2.8	2.9	3.5	6.1	13.2	50.2
1999 .....	3.1	*	*	*	1.2	2.4	2.4	3.0	3.6	4.7	13.7	56.4
<b>Diabetes mellitus (E10-E14)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	25.4	*	*	0.1	0.4	1.6	4.8	13.7	37.7	91.4	182.8	320.6
2001 .....	25.1	*	*	0.1	0.4	1.5	4.3	13.6	37.8	91.4	181.4	321.8
2000 .....	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7
1999 .....	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2
Male												
2002 .....	24.2	*	*	0.1	0.5	1.9	5.9	16.4	44.5	101.4	202.6	348.5
2001 .....	23.5	*	*	0.1	0.4	1.7	5.3	16.4	43.1	101.7	198.9	343.1
2000 .....	22.9	*	*	*	0.5	1.9	5.1	15.3	42.4	99.1	199.6	345.1
1999 .....	22.8	*	*	*	0.4	1.6	5.1	15.6	42.6	99.8	198.2	345.7
Female												
2002 .....	26.5	*	*	*	0.4	1.3	3.7	11.1	31.4	83.1	169.7	308.4
2001 .....	26.6	*	*	*	0.3	1.3	3.4	11.0	32.9	82.8	169.9	312.9
2000 .....	26.3	*	*	*	0.4	1.2	3.4	11.1	33.6	83.7	166.3	309.4
1999 .....	26.2	*	*	*	0.4	1.3	3.5	10.4	34.3	85.2	164.9	305.8
Hispanic												
Both sexes												
2002 .....	15.3	*	*	*	*	0.8	3.5	14.0	52.1	134.7	285.6	437.9
2001 .....	15.3	*	*	*	*	0.7	3.1	14.9	53.2	141.2	279.2	489.6

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Diabetes mellitus (E10-E14)</b>												
<b>Hispanic</b>												
<b>Both sexes</b>												
2000 .....	15.1	*	*	*	*	1.0	3.0	14.0	55.5	147.9	278.5	465.1
1999 .....	15.3	*	*	*	*	0.7	3.2	13.9	56.6	149.7	285.2	493.9
<b>Male</b>												
2002 .....	13.9	*	*	*	*	0.9	4.1	16.7	58.9	143.7	306.9	425.0
2001 .....	13.6	*	*	*	*	0.8	3.9	18.5	58.7	151.4	285.3	422.8
2000 .....	13.8	*	*	*	*	1.4	3.6	15.5	64.7	160.8	299.4	449.4
1999 .....	13.4	*	*	*	*	0.6	3.3	17.2	62.5	155.8	293.0	468.0
<b>Female</b>												
2002 .....	16.7	*	*	*	*	0.7	2.9	11.4	46.0	127.6	270.9	444.3
2001 .....	17.1	*	*	*	*	0.7	2.2	11.4	48.3	133.2	275.0	522.2
2000 .....	16.5	*	*	*	*	2.4	12.6	47.4	137.7	264.5	472.8	
1999 .....	17.3	*	*	*	*	0.8	3.1	10.7	51.4	144.9	280.0	506.4
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	26.9	*	*	0.1	0.5	1.8	5.0	13.6	36.4	88.2	177.2	315.0
2001 .....	26.4	*	*	0.1	0.4	1.7	4.5	13.4	36.5	87.7	176.4	314.8
2000 .....	25.9	*	*	0.1	0.5	1.7	4.4	13.0	36.4	86.8	174.8	313.4
1999 .....	25.7	*	*	0.1	0.4	1.6	4.4	12.8	36.8	88.1	173.1	310.0
<b>Male</b>												
2002 .....	25.8	*	*	0.1	0.5	2.2	6.2	16.3	43.1	98.2	196.6	344.0
2001 .....	25.0	*	*	0.1	0.4	1.9	5.5	16.1	41.8	98.0	194.3	338.9
2000 .....	24.2	*	*	*	0.5	2.0	5.3	15.2	40.7	95.0	194.6	339.7
1999 .....	24.1	*	*	*	0.4	1.8	5.3	15.3	41.0	96.3	193.8	339.5
<b>Female</b>												
2002 .....	27.9	*	*	*	0.4	1.4	3.8	11.1	30.1	79.8	164.3	302.5
2001 .....	27.8	*	*	*	0.3	1.4	3.6	10.9	31.6	79.1	164.6	304.7
2000 .....	27.6	*	*	*	0.4	1.4	3.5	11.0	32.4	80.1	161.8	302.8
1999 .....	27.3	*	*	*	0.4	1.4	3.6	10.4	32.8	81.4	159.6	298.2
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	26.4	*	*	*	0.4	1.5	4.3	11.3	30.6	76.8	163.3	296.4
2001 .....	26.0	*	*	*	0.3	1.3	3.9	11.3	30.7	77.0	162.7	297.4
2000 .....	25.4	*	*	*	0.4	1.4	3.9	10.7	30.9	76.1	159.9	295.5
1999 .....	25.0	*	*	*	0.4	1.4	3.8	10.5	30.1	76.8	159.1	290.2
<b>Male</b>												
2002 .....	26.0	*	*	*	0.4	1.9	5.5	13.8	37.5	88.9	187.4	338.4
2001 .....	25.1	*	*	*	0.3	1.5	4.8	13.7	36.4	89.8	183.5	332.1
2000 .....	24.3	*	*	*	0.5	1.7	4.8	13.0	35.4	86.6	184.5	330.4
1999 .....	24.0	*	*	*	0.4	1.5	4.6	13.0	35.0	87.7	184.6	326.0
<b>Female</b>												
2002 .....	26.8	*	*	*	0.4	1.2	3.2	8.8	24.1	66.5	147.0	278.3
2001 .....	26.9	*	*	*	0.2	1.2	2.9	8.8	25.3	66.0	148.9	283.0
2000 .....	26.5	*	*	*	0.3	1.1	3.1	8.6	26.6	67.3	143.8	281.4
1999 .....	26.0	*	*	*	0.3	1.2	3.0	8.1	25.4	67.7	142.4	275.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Diabetes mellitus (E10-E14)</b>												
Non-Hispanic black												
Both sexes												
2002 .....	34.8	*	*	*	1.0	3.8	9.9	30.9	85.9	191.5	340.2	544.7
2001 .....	34.1	*	*	*	1.0	3.8	9.1	30.1	86.5	190.8	337.9	533.8
2000 .....	34.0	*	*	*	1.0	3.6	8.2	31.0	83.9	190.6	354.8	530.1
1999 .....	34.1	*	*	*	0.8	3.1	9.1	31.0	92.0	194.0	336.1	530.9
Male												
2002 .....	30.0	*	*	*	1.1	4.6	12.1	35.8	96.6	193.3	332.6	423.4
2001 .....	29.4	*	*	*	0.9	5.0	10.5	35.0	93.8	189.5	345.1	452.7
2000 .....	28.3	*	*	*	1.0	4.0	9.7	34.2	89.7	186.9	339.4	456.8
1999 .....	28.6	*	*	*	*	3.4	10.7	35.3	95.4	189.7	317.7	506.3
Female												
2002 .....	39.2	*	*	*	0.8	3.0	8.0	26.8	77.3	190.3	344.5	591.1
2001 .....	38.4	*	*	*	1.0	2.8	7.8	25.9	80.6	191.7	333.9	563.8
2000 .....	39.1	*	*	*	1.0	3.3	7.0	28.2	79.1	193.3	363.5	557.3
1999 .....	39.0	*	*	*	0.9	2.7	7.7	27.2	89.2	197.0	346.5	540.1
<b>Nutritional deficiencies (E40-E64)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.1	8.9	41.5
2001 .....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.7	42.9
2000 .....	1.5	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.0
1999 .....	1.5	*	*	*	*	*	0.1	0.3	0.7	2.3	10.2	53.9
Male												
2002 .....	0.9	*	*	*	*	*	0.1	0.2	0.7	2.2	8.5	37.0
2001 .....	0.9	*	*	*	*	*	0.1	0.3	0.7	2.1	8.5	35.2
2000 .....	1.0	*	*	*	*	*	*	0.2	0.8	2.7	9.7	43.7
1999 .....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	10.3	48.4
Female												
2002 .....	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	9.1	43.5
2001 .....	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	8.9	46.1
2000 .....	2.0	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.6
1999 .....	2.0	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	56.2
Hispanic												
Both sexes												
2002 .....	0.4	*	*	*	*	*	*	*	*	1.8	8.2	29.4
2001 .....	0.4	*	*	*	*	*	*	*	*	*	8.1	38.8
2000 .....	0.4	*	*	*	*	*	*	*	*	2.2	8.5	39.8
1999 .....	0.4	*	*	*	*	*	*	*	*	*	6.5	51.2
Male												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	8.9	43.8
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	51.8
Female												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	8.4	29.0
2001 .....	0.5	*	*	*	*	*	*	*	*	*	7.6	36.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
Hispanic												
Female												
2000 .....	0.5	*	*	*	*	*	*	*	*	*	8.6	41.5
1999 .....	0.5	*	*	*	*	*	*	*	*	*	*	51.0
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.5	*	*	*	*	*	0.1	0.3	0.7	2.1	8.9	41.8
2001 .....	1.4	*	*	*	*	*	0.1	0.3	0.6	2.1	8.8	43.0
2000 .....	1.7	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.3
1999 .....	1.7	*	*	*	*	*	0.1	0.3	0.7	2.4	10.3	53.9
Male												
2002 .....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.2	8.5	37.1
2001 .....	1.0	*	*	*	*	*	*	0.3	0.7	2.2	8.5	34.8
2000 .....	1.1	*	*	*	*	*	*	0.3	0.9	2.7	9.6	43.9
1999 .....	1.2	*	*	*	*	*	0.1	0.3	0.9	2.7	10.4	48.1
Female												
2002 .....	1.8	*	*	*	*	*	0.1	0.3	0.6	2.0	9.1	43.9
2001 .....	1.9	*	*	*	*	*	*	0.3	0.6	2.0	8.9	46.4
2000 .....	2.2	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.8
1999 .....	2.2	*	*	*	*	*	0.2	0.3	0.6	2.1	10.3	56.2
Non-Hispanic white												
Both sexes												
2002 .....	1.5	*	*	*	*	*	0.1	0.3	0.6	1.9	8.5	40.6
2001 .....	1.5	*	*	*	*	*	0.1	0.2	0.6	2.0	8.2	42.2
2000 .....	1.8	*	*	*	*	*	0.1	0.2	0.6	2.3	10.0	49.0
1999 .....	1.8	*	*	*	*	*	0.1	0.3	0.6	2.0	9.6	52.8
Male												
2002 .....	1.1	*	*	*	*	*	*	0.2	0.6	2.0	8.0	36.5
2001 .....	1.0	*	*	*	*	*	*	0.2	0.5	1.9	7.9	33.3
2000 .....	1.2	*	*	*	*	*	*	0.2	0.7	2.3	9.0	43.0
1999 .....	1.2	*	*	*	*	*	*	0.2	0.7	2.2	9.4	46.2
Female												
2002 .....	2.0	*	*	*	*	*	*	0.3	0.6	1.8	8.8	42.3
2001 .....	2.0	*	*	*	*	*	*	0.3	0.6	2.0	8.3	45.9
2000 .....	2.3	*	*	*	*	*	0.2	0.3	0.6	2.3	10.6	51.5
1999 .....	2.3	*	*	*	*	*	0.2	0.3	0.5	1.9	9.8	55.4
Non-Hispanic black												
Both sexes												
2002 .....	1.4	*	*	*	*	*	*	0.4	1.8	4.3	14.7	60.2
2001 .....	1.3	*	*	*	*	*	*	0.8	1.2	3.7	16.1	55.2
2000 .....	1.5	*	*	*	*	*	*	0.6	1.7	4.8	16.3	67.4
1999 .....	1.7	*	*	*	*	*	*	0.7	2.0	5.6	19.6	70.8
Male												
2002 .....	1.1	*	*	*	*	*	*	*	2.9	4.5	16.4	45.9
2001 .....	1.2	*	*	*	*	*	*	1.2	1.9	5.6	16.5	58.9
2000 .....	1.3	*	*	*	*	*	*	*	2.8	6.6	17.9	64.6
1999 .....	1.5	*	*	*	*	*	*	*	2.9	8.3	23.1	76.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Nutritional deficiencies (E40-E64)</b>												
<b>Non-Hispanic black</b>												
<b>Female</b>												
2002 .....	1.6	*	*	*	*	*	*	*	*	4.1	13.8	65.6
2001 .....	1.4	*	*	*	*	*	*	*	*	2.4	15.8	53.9
2000 .....	1.8	*	*	*	*	*	*	*	*	3.6	15.5	68.5
1999 .....	1.8	*	*	*	*	*	*	*	*	3.7	17.7	68.8
<b>Malnutrition (E40-E46)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	8.4	38.4
2001 .....	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	8.2	40.2
2000 .....	1.4	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.3
1999 .....	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	9.5	50.0
<b>Male</b>												
2002 .....	0.9	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.1
2001 .....	0.8	*	*	*	*	*	*	0.2	0.6	1.9	8.0	33.6
2000 .....	0.9	*	*	*	*	*	*	0.2	0.7	2.5	8.9	40.7
1999 .....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.5	9.4	44.4
<b>Female</b>												
2002 .....	1.6	*	*	*	*	*	0.1	0.2	0.5	1.8	8.6	39.8
2001 .....	1.6	*	*	*	*	*	0.1	0.3	0.5	1.9	8.2	43.0
2000 .....	1.8	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	48.6
1999 .....	1.8	*	*	*	*	*	0.1	0.3	0.5	2.0	9.5	52.3
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	0.4	*	*	*	*	*	*	*	*	1.8	8.2	28.3
2001 .....	0.4	*	*	*	*	*	*	*	*	*	7.9	38.1
2000 .....	0.4	*	*	*	*	*	*	*	*	2.0	7.7	38.5
1999 .....	0.4	*	*	*	*	*	*	*	*	*	6.2	47.7
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	8.9	41.9
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	45.3
<b>Female</b>												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	8.4	28.2
2001 .....	0.4	*	*	*	*	*	*	*	*	*	7.3	36.3
2000 .....	0.5	*	*	*	*	*	*	*	*	*	7.6	41.5
1999 .....	0.5	*	*	*	*	*	*	*	*	*	*	48.9
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	1.3	*	*	*	*	*	0.1	0.2	0.6	1.9	8.3	38.7
2001 .....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.2	40.2
2000 .....	1.5	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.5
1999 .....	1.6	*	*	*	*	*	0.1	0.3	0.7	2.2	9.6	50.0
<b>Male</b>												
2002 .....	1.0	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Malnutrition (E40-E46)</b>												
<b>Non-Hispanic <sup>4</sup></b>												
Male												
2001 .....	0.9	*	*	*	*	*	*	0.3	0.6	2.0	8.0	33.2
2000 .....	1.0	*	*	*	*	*	*	0.2	0.8	2.5	8.8	40.9
1999 .....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	9.5	44.3
Female												
2002 .....	1.7	*	*	*	*	*	0.1	0.3	0.5	1.8	8.6	40.1
2001 .....	1.7	*	*	*	*	*	*	0.3	0.6	1.9	8.3	43.2
2000 .....	2.0	*	*	*	*	*	0.2	0.3	0.6	2.2	10.3	48.7
1999 .....	2.0	*	*	*	*	*	0.2	0.3	0.5	2.0	9.7	52.2
<b>Non-Hispanic white</b>												
Both sexes												
2002 .....	1.4	*	*	*	*	*	0.1	0.2	0.5	1.8	8.0	37.4
2001 .....	1.4	*	*	*	*	*	*	0.2	0.5	1.8	7.6	39.4
2000 .....	1.6	*	*	*	*	*	0.1	0.2	0.6	2.1	9.3	45.3
1999 .....	1.6	*	*	*	*	*	0.1	0.2	0.5	1.9	8.9	48.8
Male												
2002 .....	1.0	*	*	*	*	*	*	0.2	0.5	1.9	7.5	34.5
2001 .....	0.9	*	*	*	*	*	*	0.2	0.5	1.7	7.4	31.7
2000 .....	1.1	*	*	*	*	*	*	0.2	0.6	2.1	8.3	39.8
1999 .....	1.1	*	*	*	*	*	*	0.2	0.6	2.0	8.6	42.0
Female												
2002 .....	1.8	*	*	*	*	*	*	0.3	0.5	1.7	8.3	38.6
2001 .....	1.9	*	*	*	*	*	*	0.3	0.6	1.9	7.7	42.7
2000 .....	2.2	*	*	*	*	*	0.1	0.2	0.6	2.0	10.0	47.4
1999 .....	2.2	*	*	*	*	*	0.2	0.3	0.4	1.8	9.2	51.6
<b>Non-Hispanic black</b>												
Both sexes												
2002 .....	1.3	*	*	*	*	*	*	*	1.7	3.6	14.2	57.2
2001 .....	1.2	*	*	*	*	*	*	0.6	1.1	3.4	15.0	52.1
2000 .....	1.4	*	*	*	*	*	*	0.6	1.6	4.8	15.1	64.2
1999 .....	1.6	*	*	*	*	*	*	0.6	1.8	5.5	18.1	66.0
Male												
2002 .....	1.0	*	*	*	*	*	*	*	2.7	4.0	15.2	44.9
2001 .....	1.1	*	*	*	*	*	*	1.0	*	5.0	15.9	56.6
2000 .....	1.2	*	*	*	*	*	*	*	2.5	6.4	17.3	62.2
1999 .....	1.4	*	*	*	*	*	*	*	2.6	8.0	20.6	74.9
Female												
2002 .....	1.5	*	*	*	*	*	*	*	*	3.4	13.6	61.9
2001 .....	1.3	*	*	*	*	*	*	*	*	2.3	14.5	50.5
2000 .....	1.6	*	*	*	*	*	*	*	*	3.6	13.9	65.0
1999 .....	1.7	*	*	*	*	*	*	*	*	3.7	16.6	62.6
<b>Other nutritional deficiencies (E50-E64)</b>												
<b>All origins <sup>3</sup></b>												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	3.1
2001 .....	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
All origins <sup>3</sup>												
Both sexes												
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.7
1999 .....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.6	1.9
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.9	3.9
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.6
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.1
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.0
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.9
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	3.2
2001 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.6	2.7
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.8
1999 .....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.6	1.9
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.9	3.8
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.7
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.2
2000 .....	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.1
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.7	4.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Other nutritional deficiencies (E50-E64)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.2
2001 .....	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.8
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.6	3.8
1999 .....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
<b>Male</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.0
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.8	3.1
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.8	4.2
<b>Female</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	0.5	3.7
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.3
2000 .....	0.2	*	*	*	*	*	*	*	*	0.3	0.6	4.0
1999 .....	0.2	*	*	*	*	*	*	*	*	*	0.6	3.9
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Meningitis (G00,G03)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	0.2	1.8	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.7	1.2
2001 .....	0.3	1.9	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.8	0.9
2000 .....	0.3	2.4	0.2	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999 .....	0.3	3.1	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.5
<b>Male</b>												
2002 .....	0.3	2.0	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001 .....	0.3	1.9	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000 .....	0.3	2.5	0.3	*	*	0.1	0.2	0.3	0.5	0.7	1.3	*
1999 .....	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.0	1.7
<b>Female</b>												
2002 .....	0.2	1.7	*	*	*	*	*	0.2	0.3	0.5	0.6	1.1
2001 .....	0.2	2.0	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Meningitis (G00,G03)</b>												
<b>All origins <sup>3</sup></b>												
<b>Female</b>												
2000 .....	0.2	2.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.0
1999 .....	0.3	3.0	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	0.3	2.1	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.6	1.2
2001 .....	0.3	2.2	0.2	0.1	0.1	0.1	0.2	0.3	0.5	0.5	0.7	0.9
2000 .....	0.3	2.7	0.3	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999 .....	0.3	3.2	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.4
<b>Male</b>												
2002 .....	0.3	2.1	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001 .....	0.3	2.1	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000 .....	0.3	2.7	*	*	*	*	0.2	0.3	0.5	0.8	1.2	*
1999 .....	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.1	*
<b>Female</b>												
2002 .....	0.2	2.0	*	*	*	*	*	0.2	0.3	0.5	0.5	1.2
2001 .....	0.2	2.3	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.7
2000 .....	0.3	2.7	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.1
1999 .....	0.3	3.3	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	0.2	1.9	*	0.1	*	*	0.1	0.2	0.2	0.5	0.5	1.2
2001 .....	0.2	1.6	*	*	0.1	0.1	0.1	0.2	0.4	0.5	0.7	0.7
2000 .....	0.2	2.1	*	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.1
1999 .....	0.3	2.2	0.3	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.3
<b>Male</b>												
2002 .....	0.2	1.9	*	*	*	*	0.1	0.2	0.3	0.7	0.7	*
2001 .....	0.3	*	*	*	*	*	*	0.3	0.4	0.5	0.8	*
2000 .....	0.3	2.3	*	*	*	*	0.1	0.3	0.4	0.7	1.1	*
1999 .....	0.3	2.8	*	*	*	*	0.1	0.3	0.4	0.6	0.9	*

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Meningitis (G00,G03)</b>												
<b>Non-Hispanic white</b>												
<b>Female</b>												
2002 .....	0.2	1.8	*	*	*	*	*	0.2	0.2	0.4	0.4	1.3
2001 .....	0.2	1.9	*	*	*	*	*	0.2	0.3	0.4	0.6	*
2000 .....	0.2	1.8	*	*	*	*	*	0.2	0.2	0.3	0.8	1.0
1999 .....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8	1.2
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	0.5	3.1	*	*	*	*	*	0.9	0.9	1.3	*	*
2001 .....	0.5	4.7	*	*	*	*	0.5	0.7	1.0	*	*	*
2000 .....	0.6	5.9	*	*	*	*	0.5	0.6	0.9	1.7	*	*
1999 .....	0.6	7.0	*	*	*	*	0.4	0.9	*	1.3	2.7	*
<b>Male</b>												
2002 .....	0.6	*	*	*	*	*	*	1.3	*	*	*	*
2001 .....	0.6	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.6	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.7	*	*	*	*	*	*	1.5	*	*	*	*
<b>Female</b>												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.6	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.5	9.6	*	*	*	*	*	*	*	*	*	*
<b>Parkinson's disease (G20-G21)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	5.9	*	*	*	*	*	*	0.1	1.2	12.2	63.9	135.2
2001 .....	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.6	134.2
2000 .....	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9
1999 .....	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4
<b>Male</b>												
2002 .....	6.8	*	*	*	*	*	*	0.2	1.5	17.9	94.7	220.1
2001 .....	6.7	*	*	*	*	*	*	0.2	1.8	16.8	97.0	224.4
2000 .....	6.4	*	*	*	*	*	*	0.2	1.5	16.1	92.8	222.3
1999 .....	6.0	*	*	*	*	*	*	0.2	1.4	15.7	89.5	205.6
<b>Female</b>												
2002 .....	5.0	*	*	*	*	*	*	*	0.9	7.4	43.5	98.4
2001 .....	4.9	*	*	*	*	*	*	*	0.7	7.5	43.3	96.4
2000 .....	4.8	*	*	*	*	*	*	0.1	0.8	7.6	41.7	95.0
1999 .....	4.5	*	*	*	*	*	*	*	0.7	7.1	37.8	91.7
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	1.3	*	*	*	*	*	*	*	1.1	8.6	37.2	76.5
2001 .....	1.3	*	*	*	*	*	*	*	*	8.6	41.5	86.3
2000 .....	1.2	*	*	*	*	*	*	*	*	7.8	37.7	86.3
1999 .....	1.1	*	*	*	*	*	*	*	*	8.1	38.3	77.2
<b>Male</b>												
2002 .....	1.4	*	*	*	*	*	*	*	*	13.6	50.1	113.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
Hispanic												
Male												
2001 .....	1.4	*	*	*	*	*	*	*	*	11.2	63.0	127.6
2000 .....	1.3	*	*	*	*	*	*	*	*	9.7	55.1	143.1
1999 .....	1.3	*	*	*	*	*	*	*	*	11.4	58.9	105.7
Female												
2002 .....	1.1	*	*	*	*	*	*	*	*	4.7	28.4	58.0
2001 .....	1.2	*	*	*	*	*	*	*	*	6.6	26.9	66.1
2000 .....	1.1	*	*	*	*	*	*	*	*	6.3	26.1	58.4
1999 .....	1.0	*	*	*	*	*	*	*	*	5.5	24.6	63.4
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	6.6	*	*	*	*	*	*	0.1	1.2	12.4	65.1	137.4
2001 .....	6.5	*	*	*	*	*	*	0.1	1.2	11.9	65.5	135.6
2000 .....	6.2	*	*	*	*	*	*	0.1	1.1	11.7	62.8	133.3
1999 .....	5.8	*	*	*	*	*	*	0.1	1.1	11.1	58.9	125.8
Male												
2002 .....	7.6	*	*	*	*	*	*	0.2	1.5	18.2	96.7	224.4
2001 .....	7.5	*	*	*	*	*	*	0.2	1.8	17.1	98.3	227.8
2000 .....	7.1	*	*	*	*	*	*	0.2	1.5	16.5	94.2	225.2
1999 .....	6.7	*	*	*	*	*	*	0.2	1.5	15.9	90.6	209.4
Female												
2002 .....	5.6	*	*	*	*	*	*	*	0.9	7.6	44.2	99.8
2001 .....	5.4	*	*	*	*	*	*	*	0.7	7.6	44.0	97.3
2000 .....	5.3	*	*	*	*	*	*	*	0.8	7.7	42.3	96.1
1999 .....	4.9	*	*	*	*	*	*	*	0.7	7.1	38.3	92.3
Non-Hispanic white												
Both sexes												
2002 .....	7.9	*	*	*	*	*	*	0.2	1.3	13.3	69.6	145.8
2001 .....	7.7	*	*	*	*	*	*	0.1	1.3	12.8	69.9	143.5
2000 .....	7.3	*	*	*	*	*	*	0.1	1.2	12.6	66.7	141.2
1999 .....	6.8	*	*	*	*	*	*	0.1	1.1	11.9	62.4	132.8
Male												
2002 .....	9.1	*	*	*	*	*	*	0.2	1.6	19.3	102.9	237.8
2001 .....	9.0	*	*	*	*	*	*	0.2	2.0	18.3	104.5	241.7
2000 .....	8.4	*	*	*	*	*	*	0.2	1.6	17.6	99.2	238.6
1999 .....	7.9	*	*	*	*	*	*	0.2	1.6	17.1	95.7	221.0
Female												
2002 .....	6.7	*	*	*	*	*	*	*	1.0	8.1	47.3	106.0
2001 .....	6.5	*	*	*	*	*	*	*	0.7	8.2	46.9	102.7
2000 .....	6.3	*	*	*	*	*	*	*	0.9	8.3	45.3	101.9
1999 .....	5.8	*	*	*	*	*	*	*	0.7	7.6	40.5	97.7
Non-Hispanic black												
Both sexes												
2002 .....	1.5	*	*	*	*	*	*	*	*	6.7	25.7	51.7
2001 .....	1.5	*	*	*	*	*	*	*	0.8	5.6	25.7	59.0
2000 .....	1.5	*	*	*	*	*	*	*	*	5.8	27.3	52.5
1999 .....	1.4	*	*	*	*	*	*	*	0.9	5.9	24.8	51.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
Non-Hispanic black												
Male												
2002 .....	1.8	*	*	*	*	*	*	*	*	11.1	40.4	87.5
2001 .....	1.7	*	*	*	*	*	*	*	*	8.8	38.1	99.3
2000 .....	1.8	*	*	*	*	*	*	*	*	9.0	47.5	88.1
1999 .....	1.6	*	*	*	*	*	*	*	*	8.7	38.5	87.9
Female												
2002 .....	1.3	*	*	*	*	*	*	*	*	3.6	17.4	38.0
2001 .....	1.4	*	*	*	*	*	*	*	*	3.3	18.7	44.1
2000 .....	1.2	*	*	*	*	*	*	*	*	3.5	16.0	39.2
1999 .....	1.3	*	*	*	*	*	*	*	*	3.9	17.2	38.2
Alzheimer's disease (G30)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	20.4	*	*	*	*	*	*	0.1	1.9	19.7	158.1	752.3
2001 .....	18.9	*	*	*	*	*	*	0.2	2.1	18.7	147.5	710.3
2000 .....	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7
1999 .....	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3
Male												
2002 .....	12.0	*	*	*	*	*	*	0.1	1.8	18.6	142.2	572.0
2001 .....	11.3	*	*	*	*	*	*	0.2	1.9	17.7	136.3	556.1
2000 .....	10.5	*	*	*	*	*	*	0.2	1.8	18.7	127.6	522.1
1999 .....	9.8	*	*	*	*	*	*	0.2	1.9	17.3	122.6	485.7
Female												
2002 .....	28.5	*	*	*	*	*	*	0.1	2.0	20.6	168.7	830.5
2001 .....	26.3	*	*	*	*	*	*	0.2	2.3	19.5	154.8	774.9
2000 .....	24.5	*	*	*	*	*	*	0.2	2.2	18.6	147.4	727.0
1999 .....	21.9	*	*	*	*	*	*	0.2	1.8	17.5	134.0	648.0
Hispanic												
Both sexes												
2002 .....	3.8	*	*	*	*	*	*	*	1.0	10.8	89.2	438.5
2001 .....	3.4	*	*	*	*	*	*	*	*	9.9	83.9	399.5
2000 .....	3.2	*	*	*	*	*	*	*	*	11.9	79.2	380.2
1999 .....	2.9	*	*	*	*	*	*	*	*	9.7	75.4	351.5
Male												
2002 .....	2.3	*	*	*	*	*	*	*	*	10.5	80.4	341.3
2001 .....	2.1	*	*	*	*	*	*	*	*	8.8	83.2	306.6
2000 .....	2.1	*	*	*	*	*	*	*	*	13.3	81.6	284.2
1999 .....	1.8	*	*	*	*	*	*	*	*	11.2	66.7	278.2
Female												
2002 .....	5.4	*	*	*	*	*	*	*	*	10.9	95.3	486.6
2001 .....	4.7	*	*	*	*	*	*	*	*	10.7	84.3	444.9
2000 .....	4.3	*	*	*	*	*	*	*	*	10.8	77.6	427.3
1999 .....	4.0	*	*	*	*	*	*	*	*	8.4	81.2	386.8
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	22.9	*	*	*	*	*	*	0.1	2.0	20.3	161.1	763.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2001 .....	21.2	*	*	*	*	*	*	0.2	2.2	19.2	150.2	720.5
2000 .....	19.6	*	*	*	*	*	*	0.2	2.1	19.0	142.0	676.7
1999 .....	17.7	*	*	*	*	*	*	0.2	2.0	17.8	131.5	608.9
Male												
2002 .....	13.6	*	*	*	*	*	*	0.2	1.9	19.1	145.0	581.4
2001 .....	12.7	*	*	*	*	*	*	0.2	2.0	18.3	138.6	565.2
2000 .....	11.7	*	*	*	*	*	*	0.2	1.8	19.0	129.6	531.0
1999 .....	10.9	*	*	*	*	*	*	0.2	2.0	17.7	124.8	492.8
Female												
2002 .....	31.9	*	*	*	*	*	*	0.1	2.1	21.2	171.8	841.6
2001 .....	29.3	*	*	*	*	*	*	0.2	2.4	20.0	157.8	785.1
2000 .....	27.2	*	*	*	*	*	*	0.2	2.3	19.1	150.0	735.6
1999 .....	24.2	*	*	*	*	*	*	0.2	1.9	18.0	135.9	655.4
Non-Hispanic white												
Both sexes												
2002 .....	26.9	*	*	*	*	*	*	0.2	2.1	21.3	168.2	793.9
2001 .....	24.8	*	*	*	*	*	*	0.2	2.4	20.3	155.6	749.2
2000 .....	22.9	*	*	*	*	*	*	0.2	2.2	19.9	147.4	705.1
1999 .....	20.7	*	*	*	*	*	*	0.2	2.1	18.6	136.7	636.6
Male												
2002 .....	15.8	*	*	*	*	*	*	0.2	2.0	19.9	150.5	605.4
2001 .....	14.7	*	*	*	*	*	*	0.2	2.1	19.1	143.0	587.2
2000 .....	13.6	*	*	*	*	*	*	0.2	2.0	19.8	134.6	551.7
1999 .....	12.7	*	*	*	*	*	*	0.2	2.2	18.3	129.8	515.1
Female												
2002 .....	37.6	*	*	*	*	*	*	0.2	2.3	22.5	180.1	875.3
2001 .....	34.4	*	*	*	*	*	*	0.2	2.7	21.3	163.9	816.6
2000 .....	31.9	*	*	*	*	*	*	0.2	2.5	20.0	155.8	767.0
1999 .....	28.4	*	*	*	*	*	*	0.2	2.0	18.8	141.2	685.0
Non-Hispanic black												
Both sexes												
2002 .....	9.1	*	*	*	*	*	*	*	1.2	16.8	124.2	548.7
2001 .....	8.7	*	*	*	*	*	*	*	1.1	15.5	126.9	513.2
2000 .....	7.7	*	*	*	*	*	*	*	1.3	16.1	111.6	450.9
1999 .....	6.7	*	*	*	*	*	*	*	1.2	15.5	99.1	374.7
Male												
2002 .....	5.5	*	*	*	*	*	*	*	*	17.9	122.5	414.6
2001 .....	5.3	*	*	*	*	*	*	*	*	16.3	123.6	420.4
2000 .....	4.6	*	*	*	*	*	*	*	*	16.9	98.7	378.1
1999 .....	4.2	*	*	*	*	*	*	*	*	16.8	90.7	320.9
Female												
2002 .....	12.5	*	*	*	*	*	*	*	*	16.1	125.2	599.9
2001 .....	11.8	*	*	*	*	*	*	*	*	15.0	128.7	547.5
2000 .....	10.6	*	*	*	*	*	*	*	*	15.6	118.8	478.0
1999 .....	9.0	*	*	*	*	*	*	*	*	14.5	103.8	394.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Major cardiovascular diseases (I00-I78)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	318.6	15.9	1.4	0.9	3.2	10.0	37.8	114.3	294.8	788.4	2,257.0	7,389.6
2001 .....	323.9	14.9	1.9	0.9	3.2	10.2	37.1	113.4	301.2	810.4	2,322.2	7,673.7
2000 .....	332.9	16.7	1.5	0.9	3.3	9.5	36.8	115.7	319.2	849.3	2,396.2	8,030.3
1999 .....	340.6	17.0	1.5	0.9	3.4	9.6	37.7	116.2	327.3	887.3	2,474.9	8,197.1
<b>Male</b>												
2002 .....	303.3	16.6	1.5	0.9	4.1	12.7	50.8	162.3	407.0	1,027.8	2,731.1	7,638.8
2001 .....	306.2	15.3	1.9	0.9	3.9	12.8	49.9	160.4	414.0	1,057.6	2,823.3	7,936.0
2000 .....	315.8	17.5	1.6	1.0	3.9	11.8	49.5	165.1	441.6	1,114.1	2,926.6	8,448.1
1999 .....	323.0	17.9	1.6	1.0	4.0	12.1	51.2	165.8	452.0	1,168.4	3,024.0	8,603.2
<b>Female</b>												
2002 .....	333.3	15.1	1.3	0.8	2.2	7.2	24.9	68.0	191.0	589.1	1,942.2	7,281.5
2001 .....	340.9	14.5	1.8	0.9	2.4	7.6	24.3	68.1	196.9	605.6	1,992.9	7,563.9
2000 .....	349.5	15.9	1.4	0.8	2.8	7.2	24.3	68.0	206.3	631.2	2,050.4	7,860.1
1999 .....	357.5	16.0	1.4	0.8	2.7	7.2	24.4	68.5	212.5	657.0	2,118.2	8,033.5
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	94.3	13.1	1.2	0.7	2.6	6.1	20.4	73.6	225.8	621.8	1,728.4	5,369.7
2001 .....	96.2	12.0	1.5	0.8	2.4	6.4	19.8	75.4	216.4	644.9	1,828.4	6,036.2
2000 .....	96.5	13.7	1.3	0.7	2.6	6.4	21.3	75.0	236.3	660.8	1,852.8	6,168.2
1999 .....	99.2	15.2	1.4	0.7	2.5	6.5	21.4	80.5	242.7	690.0	1,935.5	6,335.9
<b>Male</b>												
2002 .....	94.2	12.2	*	0.6	3.3	8.1	27.4	102.9	316.5	822.6	2,067.7	5,494.9
2001 .....	95.4	12.7	1.4	0.8	3.0	8.3	26.8	107.2	297.2	852.4	2,220.5	6,155.6
2000 .....	96.0	13.7	1.4	0.9	3.1	8.1	27.9	107.7	330.6	870.1	2,272.6	6,304.3
1999 .....	99.2	12.7	1.4	0.9	3.0	8.6	29.7	116.3	333.2	925.3	2,356.2	6,595.2
<b>Female</b>												
2002 .....	94.4	14.0	1.3	0.8	1.7	3.7	12.8	44.4	144.4	462.0	1,496.0	5,307.7
2001 .....	97.1	11.2	1.6	0.8	1.8	4.2	12.3	44.1	144.6	481.2	1,563.4	5,977.9
2000 .....	96.9	13.8	*	*	2.0	4.5	14.3	42.9	152.6	496.2	1,570.9	6,101.4
1999 .....	99.1	17.9	1.4	*	2.0	4.0	12.7	45.7	162.6	505.6	1,655.5	6,210.9
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	352.2	16.5	1.5	0.9	3.3	10.8	40.2	117.8	299.1	796.7	2,276.3	7,451.4
2001 .....	356.7	15.5	1.9	0.9	3.3	10.9	39.3	116.5	306.3	818.6	2,338.1	7,715.2
2000 .....	365.6	17.3	1.6	0.9	3.5	10.1	38.6	118.8	323.8	857.8	2,412.2	8,076.4
1999 .....	372.7	17.4	1.5	0.9	3.6	10.2	39.5	118.8	331.8	895.4	2,489.2	8,239.8
<b>Male</b>												
2002 .....	336.4	17.6	1.7	1.0	4.2	13.8	54.2	167.5	412.3	1,037.1	2,755.7	7,713.8
2001 .....	338.1	15.8	2.0	0.9	4.1	13.7	53.0	164.7	420.6	1,066.8	2,842.6	7,990.6
2000 .....	347.7	18.3	1.7	1.0	4.1	12.6	52.0	169.4	447.3	1,124.0	2,946.0	8,513.1
1999 .....	354.2	19.1	1.7	1.0	4.2	12.7	53.6	169.3	457.9	1,177.2	3,041.8	8,657.6
<b>Female</b>												
2002 .....	367.2	15.3	1.3	0.8	2.3	7.9	26.5	70.0	194.1	595.9	1,958.5	7,338.2
2001 .....	374.3	15.2	1.8	0.9	2.5	8.2	25.9	70.1	200.4	612.3	2,006.9	7,600.7
2000 .....	382.6	16.3	1.5	0.8	2.9	7.6	25.4	70.0	209.6	638.0	2,064.5	7,899.8
1999 .....	390.2	15.5	1.4	0.8	2.9	7.7	25.7	70.2	215.4	664.0	2,130.7	8,072.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Major cardiovascular diseases (I00-I78)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	381.9	13.4	1.4	0.8	2.7	8.6	34.9	102.9	272.0	760.6	2,252.2	7,546.1
2001 .....	385.8	12.1	1.5	0.8	2.7	8.9	33.3	101.3	278.5	779.4	2,313.0	7,811.4
2000 .....	395.5	14.6	1.4	0.8	2.8	8.1	32.8	103.9	295.3	820.2	2,384.9	8,179.0
1999 .....	402.7	14.2	1.3	0.8	3.0	8.3	33.3	104.1	302.5	856.8	2,462.9	8,342.9
<b>Male</b>												
2002 .....	362.2	14.8	1.6	0.9	3.5	11.1	48.7	151.1	381.0	1,001.2	2,744.3	7,845.8
2001 .....	363.0	13.0	1.6	0.8	3.3	11.4	46.7	148.1	389.7	1,027.1	2,827.4	8,116.3
2000 .....	373.6	15.5	1.6	0.9	3.3	10.5	46.1	153.2	415.4	1,088.2	2,930.7	8,650.5
1999 .....	379.8	16.1	1.5	0.8	3.4	10.6	46.9	152.7	424.4	1,139.8	3,030.2	8,776.1
<b>Female</b>												
2002 .....	400.8	12.0	1.1	0.7	1.9	6.0	21.1	55.4	168.8	555.5	1,922.5	7,416.6
2001 .....	407.6	11.0	1.5	0.8	2.0	6.4	19.9	55.3	173.3	569.4	1,971.8	7,684.6
2000 .....	416.5	13.6	1.1	0.6	2.2	5.7	19.5	55.3	182.0	594.6	2,026.2	7,988.9
1999 .....	424.7	12.1	1.1	0.7	2.5	6.0	19.8	56.2	187.7	620.0	2,091.4	8,170.1
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	285.1	28.3	2.1	1.4	6.0	24.3	78.2	242.6	578.0	1,243.1	2,851.9	7,136.9
2001 .....	289.7	30.1	3.7	1.3	6.8	23.5	80.7	243.2	587.2	1,293.5	2,915.8	7,366.6
2000 .....	294.2	28.8	2.5	1.5	7.0	22.3	79.1	246.4	609.6	1,333.1	2,995.1	7,576.5
1999 .....	298.5	30.2	2.6	1.4	7.0	21.6	83.7	246.8	623.7	1,382.8	3,044.9	7,624.4
<b>Male</b>												
2002 .....	277.2	29.1	2.0	1.5	7.3	30.1	97.5	319.3	775.5	1,576.9	3,318.5	6,982.7
2001 .....	281.2	29.7	3.8	1.2	8.3	28.9	100.1	319.1	772.1	1,639.6	3,461.6	7,413.8
2000 .....	284.3	30.0	2.3	1.4	7.8	26.5	97.6	324.0	807.5	1,658.1	3,527.8	7,755.5
1999 .....	291.1	31.5	2.7	1.6	8.7	26.6	105.2	329.7	833.1	1,738.6	3,542.5	7,952.2
<b>Female</b>												
2002 .....	292.3	27.4	2.2	1.4	4.7	18.9	61.2	176.8	419.5	1,006.4	2,588.0	7,195.9
2001 .....	297.4	30.6	3.6	1.4	5.3	18.6	63.5	178.1	438.5	1,049.3	2,609.4	7,349.1
2000 .....	303.1	27.5	2.9	1.5	6.3	18.6	62.8	179.8	450.3	1,104.6	2,695.3	7,510.0
1999 .....	305.1	28.9	2.4	1.3	5.3	17.2	64.7	176.0	455.9	1,131.8	2,766.3	7,501.9
<b>Diseases of heart (I00-I09,I11,I13,I20-I51)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	241.7	12.4	1.1	0.6	2.5	7.9	30.5	93.7	241.5	615.9	1,677.2	5,446.8
2001 .....	245.8	11.9	1.5	0.7	2.5	8.0	29.6	92.9	246.9	635.1	1,725.7	5,664.2
2000 .....	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2	665.6	1,780.3	5,926.1
1999 .....	259.9	13.8	1.2	0.7	2.8	7.6	30.2	95.7	269.9	701.7	1,849.9	6,063.0
<b>Male</b>												
2002 .....	240.7	12.9	1.1	0.7	3.3	10.5	43.1	138.4	343.4	827.1	2,110.1	5,823.5
2001 .....	242.5	11.8	1.5	0.7	3.2	10.3	41.7	136.6	349.8	851.3	2,177.3	6,040.5
2000 .....	249.8	13.3	1.4	0.8	3.2	9.6	41.4	140.2	371.7	898.3	2,248.1	6,430.0
1999 .....	257.0	13.9	1.3	0.8	3.4	9.8	43.0	142.0	383.0	950.6	2,336.4	6,561.6
<b>Female</b>												
2002 .....	242.7	11.8	1.0	0.6	1.7	5.2	18.0	50.6	147.2	440.1	1,389.7	5,283.3
2001 .....	249.0	12.0	1.4	0.7	1.8	5.6	17.6	50.7	151.8	455.9	1,428.9	5,506.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Diseases of heart (I00-I09,I11,I13,I20-I51)</b>												
<b>All origins <sup>3</sup></b>												
<b>Female</b>												
2002 .....	255.3	12.5	1.0	0.5	2.1	5.2	17.2	49.8	159.3	474.0	1,475.1	5,720.9
1999 .....	262.6	13.7	1.1	0.6	2.1	5.4	17.5	51.0	165.6	497.9	1,533.9	5,862.1
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	71.9	10.2	0.8	0.4	1.9	4.4	14.7	55.3	176.8	484.3	1,297.6	4,121.9
2001 .....	73.3	9.7	1.1	0.5	1.7	4.6	13.8	55.2	170.7	504.5	1,376.3	4,607.5
2000 .....	73.1	11.2	1.0	0.6	1.9	4.3	14.8	55.9	183.3	506.5	1,395.9	4,701.8
1999 .....	76.2	12.7	1.2	0.6	1.8	4.5	15.4	60.9	192.7	536.3	1,467.8	4,888.6
<b>Male</b>												
2002 .....	74.0	9.6	*	*	2.6	6.1	20.6	80.5	256.0	657.7	1,599.5	4,301.8
2001 .....	74.6	10.3	*	*	2.1	6.1	20.0	82.9	242.2	683.7	1,702.7	4,784.3
2000 .....	74.7	11.2	1.4	0.7	2.4	5.7	20.7	84.3	264.8	684.8	1,733.2	4,897.5
1999 .....	77.7	10.3	1.4	0.7	2.1	6.5	22.5	92.2	272.1	737.0	1,816.0	5,012.2
<b>Female</b>												
2002 .....	69.7	10.8	*	*	1.1	2.4	8.2	30.2	105.7	346.4	1,090.8	4,032.8
2001 .....	71.8	9.0	*	*	1.2	2.8	7.1	27.9	107.2	363.1	1,155.7	4,521.1
2000 .....	71.5	11.2	*	*	1.3	2.7	8.5	28.2	111.2	366.3	1,169.4	4,605.8
1999 .....	74.6	15.2	*	*	1.5	2.3	7.9	30.5	122.5	379.0	1,236.1	4,828.9
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	267.1	12.9	1.1	0.7	2.6	8.7	32.7	97.1	245.5	622.2	1,690.6	5,484.9
2001 .....	270.7	12.3	1.5	0.7	2.6	8.7	31.6	96.0	251.6	641.4	1,736.4	5,688.6
2000 .....	277.3	13.3	1.2	0.7	2.8	8.0	30.8	97.1	265.7	672.7	1,791.1	5,953.6
1999 .....	284.3	14.0	1.2	0.7	2.9	8.2	31.8	98.3	274.1	708.5	1,859.2	6,086.3
<b>Male</b>												
2002 .....	267.0	13.7	1.2	0.7	3.4	11.6	46.4	143.6	348.6	834.6	2,128.7	5,874.4
2001 .....	268.0	12.0	1.7	0.7	3.3	11.2	44.6	141.1	356.1	858.7	2,192.5	6,077.8
2000 .....	275.2	13.7	1.4	0.8	3.3	10.3	43.8	144.5	377.5	907.0	2,263.4	6,474.6
1999 .....	282.0	14.8	1.3	0.8	3.6	10.5	45.4	145.7	388.7	958.3	2,349.8	6,602.0
<b>Female</b>												
2002 .....	267.2	12.0	1.1	0.6	1.8	5.8	19.3	52.4	150.0	444.8	1,400.2	5,316.8
2001 .....	273.3	12.7	1.4	0.7	1.9	6.1	18.9	52.6	154.8	460.9	1,437.0	5,526.8
2000 .....	279.3	12.8	1.1	0.6	2.2	5.7	18.1	51.5	162.2	479.3	1,483.4	5,742.9
1999 .....	286.4	13.3	1.1	0.6	2.3	5.9	18.5	52.6	168.2	503.4	1,540.8	5,879.8
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	290.8	10.7	1.0	0.6	2.1	6.9	29.3	87.5	227.2	599.7	1,679.2	5,568.4
2001 .....	294.1	9.8	1.2	0.6	2.1	7.1	27.6	86.4	233.0	616.7	1,724.0	5,776.0
2000 .....	301.3	11.4	1.1	0.5	2.1	6.4	27.2	87.7	246.4	648.1	1,777.4	6,047.9
1999 .....	308.2	11.3	1.0	0.6	2.4	6.7	27.6	88.7	254.0	683.2	1,846.1	6,176.6
<b>Male</b>												
2002 .....	289.2	11.6	1.1	0.7	2.8	9.4	42.9	133.1	327.6	813.5	2,129.9	5,994.1
2001 .....	289.5	10.3	1.3	0.6	2.6	9.4	40.4	130.7	335.8	834.7	2,190.4	6,195.4
2000 .....	297.5	12.0	1.2	0.7	2.6	8.6	40.1	134.3	356.3	885.1	2,261.9	6,606.6
1999 .....	304.0	12.4	1.1	0.6	2.8	8.7	40.6	135.0	365.6	935.2	2,352.6	6,713.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Diseases of heart (I00-I09,I11,I13,I20-I51)</b>												
<b>Non-Hispanic white</b>												
Female												
2002 .....	292.3	9.6	0.8	0.5	1.4	4.4	15.8	42.6	132.0	417.4	1,377.2	5,384.5
2001 .....	298.4	9.2	1.1	0.6	1.5	4.7	14.8	42.7	136.0	431.8	1,414.7	5,601.6
2000 .....	304.9	10.8	0.9	0.4	1.7	4.2	14.3	41.9	142.9	448.5	1,458.9	5,822.7
1999 .....	312.3	10.2	0.8	0.6	1.9	4.6	14.5	43.1	149.0	472.3	1,514.5	5,962.4
<b>Non-Hispanic black</b>												
Both sexes												
2002 .....	212.2	21.9	1.7	1.0	5.1	19.2	59.4	184.7	450.2	936.3	2,067.0	5,151.0
2001 .....	215.5	22.9	3.1	1.1	5.6	18.7	61.3	183.6	454.5	977.9	2,123.4	5,307.7
2000 .....	218.6	20.5	2.0	1.2	5.9	17.9	58.7	185.4	473.2	1,014.8	2,177.9	5,424.6
1999 .....	223.8	24.8	2.1	1.2	6.0	17.5	63.6	189.1	488.5	1,058.0	2,223.2	5,514.7
Male												
2002 .....	212.9	23.0	*	1.0	6.3	25.0	77.6	250.9	615.0	1,207.5	2,470.0	5,161.2
2001 .....	215.7	20.7	3.4	1.0	7.1	23.6	78.9	247.6	610.5	1,259.3	2,586.9	5,495.4
2000 .....	217.6	20.1	2.0	1.2	6.9	22.1	75.6	252.5	639.6	1,281.7	2,615.1	5,677.2
1999 .....	224.9	24.6	2.3	1.4	7.7	22.1	83.8	259.8	668.4	1,355.1	2,631.6	5,877.1
Female												
2002 .....	211.6	20.8	1.9	1.0	3.9	14.1	43.3	128.0	317.9	744.0	1,839.1	5,147.0
2001 .....	215.4	25.2	2.9	1.1	4.0	14.2	45.7	128.7	328.9	779.4	1,863.2	5,238.2
2000 .....	219.5	20.8	2.0	1.1	4.9	14.1	43.7	127.8	339.3	827.1	1,931.9	5,330.8
1999 .....	222.8	25.0	1.9	1.0	4.4	13.3	45.8	128.7	344.3	848.5	1,994.5	5,379.3
<b>Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)</b>												
<b>All origins <sup>3</sup></b>												
Both sexes												
2002 .....	1.2	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.6	17.8
2001 .....	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.4	17.7
2000 .....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.5	16.8
1999 .....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	18.0
Male												
2002 .....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	6.7	11.8
2001 .....	0.7	*	*	*	*	*	0.2	0.5	1.1	2.8	6.6	12.8
2000 .....	0.8	*	*	*	*	*	0.3	0.6	1.4	2.9	6.6	12.2
1999 .....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.5	6.8	10.3
Female												
2002 .....	1.7	*	*	*	*	*	0.2	0.7	2.0	4.6	11.6	20.4
2001 .....	1.7	*	*	*	*	*	0.3	0.7	1.7	5.1	11.3	19.8
2000 .....	1.8	*	*	*	*	0.1	0.4	0.8	2.0	5.8	11.4	18.7
1999 .....	1.9	*	*	*	*	0.1	0.3	0.8	2.2	5.6	12.2	21.1
<b>Hispanic</b>												
Both sexes												
2002 .....	0.5	*	*	*	*	*	*	0.7	1.8	4.6	7.0	*
2001 .....	0.4	*	*	*	*	*	*	*	1.5	4.2	5.4	*
2000 .....	0.4	*	*	*	*	*	*	*	1.8	4.4	5.3	*
1999 .....	0.5	*	*	*	*	*	*	0.8	1.6	4.2	7.3	*

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Hispanic												
Male												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.8	*	*	*	*	*	*	*	2.1	5.9	10.7	*
2001 .....	0.6	*	*	*	*	*	*	*	2.0	5.2	*	*
2000 .....	0.6	*	*	*	*	*	*	*	*	5.6	6.9	*
1999 .....	0.8	*	*	*	*	*	*	*	*	5.9	10.4	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.4	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.7	18.1
2001 .....	1.3	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.6	18.0
2000 .....	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.7	17.1
1999 .....	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.1	18.3
Male												
2002 .....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	7.0	12.0
2001 .....	0.8	*	*	*	*	*	0.3	0.5	1.1	2.8	6.7	13.1
2000 .....	0.9	*	*	*	*	*	0.3	0.6	1.4	2.9	6.7	12.5
1999 .....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.6	6.9	10.7
Female												
2002 .....	1.8	*	*	*	*	*	0.2	0.7	2.0	4.5	11.6	20.8
2001 .....	1.8	*	*	*	*	*	0.3	0.7	1.7	5.1	11.5	20.0
2000 .....	1.9	*	*	*	*	*	0.4	0.8	2.0	5.8	11.6	19.0
1999 .....	2.0	*	*	*	*	0.1	0.3	0.8	2.3	5.5	12.2	21.3
Non-Hispanic white												
Both sexes												
2002 .....	1.5	*	*	*	*	*	0.2	0.5	1.5	3.9	10.2	19.3
2001 .....	1.5	*	*	*	*	*	0.2	0.5	1.3	4.0	10.2	19.0
2000 .....	1.6	*	*	*	*	*	0.3	0.6	1.7	4.6	10.3	18.0
1999 .....	1.6	*	*	*	*	0.1	0.2	0.6	1.7	4.8	10.6	19.2
Male												
2002 .....	0.9	*	*	*	*	*	0.2	0.4	1.1	3.0	7.3	12.9
2001 .....	0.9	*	*	*	*	*	0.2	0.4	1.1	2.7	7.0	13.7
2000 .....	0.9	*	*	*	*	*	0.3	0.5	1.5	2.9	7.2	13.0
1999 .....	0.9	*	*	*	*	*	0.2	0.5	1.1	3.8	7.2	11.6
Female												
2002 .....	2.1	*	*	*	*	*	0.2	0.6	1.9	4.6	12.2	22.0
2001 .....	2.0	*	*	*	*	*	0.2	0.5	1.5	5.1	12.2	21.2
2000 .....	2.1	*	*	*	*	*	0.3	0.7	1.9	6.0	12.4	20.0
1999 .....	2.2	*	*	*	*	*	0.2	0.7	2.2	5.7	12.9	22.3
Non-Hispanic black												
Both sexes												
2002 .....	0.7	*	*	*	*	*	0.4	1.0	2.4	2.9	4.4	*
2001 .....	0.8	*	*	*	*	*	0.7	1.4	2.1	4.2	4.3	6.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Non-Hispanic black												
Both sexes												
2000 .....	0.7	*	*	*	*	*	0.6	1.0	2.0	3.6	4.0	8.6
1999 .....	0.8	*	*	*	*	*	0.5	1.2	2.2	3.4	4.4	8.4
Male												
2002 .....	0.5	*	*	*	*	*	*	*	1.8	2.9	*	*
2001 .....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000 .....	0.5	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.5	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.8	*	*	*	*	*	*	1.1	2.9	2.9	5.1	*
2001 .....	1.2	*	*	*	*	*	0.9	1.7	2.6	5.4	5.4	*
2000 .....	1.0	*	*	*	*	*	0.8	1.4	2.7	4.4	4.9	*
1999 .....	1.0	*	*	*	*	*	*	1.5	2.7	4.1	4.4	11.1
Hypertensive heart disease (I11)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	9.2	*	*	*	0.1	0.7	2.7	7.2	12.5	21.3	49.6	187.0
2001 .....	8.7	*	*	*	0.1	0.6	2.4	6.3	10.9	19.8	48.7	189.5
2000 .....	8.4	*	*	*	0.1	0.5	2.2	5.6	11.0	19.6	48.2	193.1
1999 .....	8.1	*	*	*	0.1	0.5	2.1	5.3	10.7	19.1	47.7	184.9
Male												
2002 .....	8.2	*	*	*	0.1	1.1	3.8	9.8	16.9	25.2	49.4	134.6
2001 .....	7.5	*	*	*	*	0.8	3.3	8.6	14.4	23.6	48.4	134.4
2000 .....	7.2	*	*	*	0.1	0.7	2.9	7.8	14.5	23.7	46.5	139.1
1999 .....	6.8	*	*	*	0.1	0.7	2.9	7.0	13.7	21.9	46.8	133.6
Female												
2002 .....	10.2	*	*	*	*	0.4	1.6	4.6	8.4	18.1	49.8	209.7
2001 .....	9.8	*	*	*	*	0.4	1.6	4.1	7.8	16.6	48.8	212.6
2000 .....	9.7	*	*	*	*	0.4	1.5	3.5	7.8	16.2	49.4	215.1
1999 .....	9.4	*	*	*	*	0.3	1.3	3.6	7.9	16.8	48.3	205.5
Hispanic												
Both sexes												
2002 .....	3.7	*	*	*	*	0.4	1.7	4.9	12.2	22.4	51.4	165.7
2001 .....	3.3	*	*	*	*	0.3	1.7	4.9	10.0	21.0	46.9	151.9
2000 .....	2.8	*	*	*	*	0.3	1.2	3.8	8.2	18.4	46.0	143.3
1999 .....	2.7	*	*	*	*	*	1.2	3.3	8.9	16.8	44.8	152.2
Male												
2002 .....	3.9	*	*	*	*	0.6	2.6	7.4	17.0	28.5	61.1	137.2
2001 .....	3.4	*	*	*	*	*	2.7	7.0	13.8	26.4	48.3	144.7
2000 .....	2.7	*	*	*	*	*	1.6	5.4	10.7	22.4	50.6	112.9
1999 .....	2.6	*	*	*	*	*	1.8	4.7	12.5	20.9	42.8	107.8
Female												
2002 .....	3.4	*	*	*	*	*	*	2.5	7.9	17.5	44.7	179.9
2001 .....	3.1	*	*	*	*	*	*	2.7	6.5	16.8	45.9	155.5
2000 .....	2.9	*	*	*	*	*	0.8	2.1	6.1	15.3	42.9	158.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Hispanic												
Female												
1999 .....	2.9	*	*	*	*	*	*	2.0	5.8	13.6	46.1	173.7
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	10.0	*	*	*	0.1	0.8	2.8	7.4	12.4	21.0	49.3	187.1
2001 .....	9.4	*	*	*	*	0.7	2.5	6.3	10.9	19.6	48.6	190.3
2000 .....	9.2	*	*	*	0.1	0.6	2.3	5.7	11.2	19.5	48.1	194.3
1999 .....	8.8	*	*	*	0.1	0.6	2.2	5.4	10.7	19.1	47.6	185.3
Male												
2002 .....	8.9	*	*	*	0.1	1.2	4.0	10.0	16.8	24.6	48.5	133.7
2001 .....	8.1	*	*	*	*	0.9	3.4	8.6	14.2	23.3	48.3	133.4
2000 .....	7.8	*	*	*	0.1	0.7	3.0	7.9	14.6	23.5	46.0	139.6
1999 .....	7.4	*	*	*	0.1	0.8	3.1	7.2	13.6	21.7	46.7	134.2
Female												
2002 .....	11.1	*	*	*	*	0.5	1.7	4.8	8.4	18.0	49.9	210.2
2001 .....	10.7	*	*	*	*	0.4	1.7	4.2	7.8	16.5	48.8	214.0
2000 .....	10.6	*	*	*	*	0.4	1.6	3.6	8.0	16.2	49.5	216.5
1999 .....	10.2	*	*	*	*	0.3	1.4	3.7	8.0	16.9	48.2	205.8
Non-Hispanic white												
Both sexes												
2002 .....	9.1	*	*	*	*	0.4	1.9	5.1	8.8	16.1	42.2	177.0
2001 .....	8.6	*	*	*	*	0.4	1.6	4.1	7.6	14.8	42.2	180.4
2000 .....	8.3	*	*	*	*	0.3	1.5	3.6	7.5	14.5	41.2	182.5
1999 .....	7.9	*	*	*	*	0.3	1.3	3.4	7.3	14.1	40.6	175.2
Male												
2002 .....	7.6	*	*	*	*	0.7	2.8	7.3	12.2	19.0	41.2	123.5
2001 .....	6.9	*	*	*	*	0.6	2.3	5.9	10.5	18.1	41.5	122.0
2000 .....	6.6	*	*	*	*	0.4	2.1	5.4	10.3	18.0	38.8	129.5
1999 .....	6.2	*	*	*	*	0.6	2.0	4.9	9.6	16.2	39.4	123.8
Female												
2002 .....	10.5	*	*	*	*	0.2	1.0	2.9	5.5	13.6	42.9	200.1
2001 .....	10.2	*	*	*	*	0.2	0.8	2.3	4.9	11.9	42.6	204.7
2000 .....	9.9	*	*	*	*	0.2	0.8	1.9	4.9	11.6	42.7	203.9
1999 .....	9.6	*	*	*	*	*	0.6	2.0	5.0	12.4	41.4	195.7
Non-Hispanic black												
Both sexes												
2002 .....	17.7	*	*	*	*	2.9	8.8	24.8	45.1	68.3	134.2	322.1
2001 .....	16.5	*	*	*	*	2.2	8.5	23.2	39.4	66.1	125.6	328.7
2000 .....	16.6	*	*	*	*	2.2	7.5	21.6	43.1	67.6	132.7	344.1
1999 .....	16.0	*	*	*	0.4	1.9	8.1	21.3	40.1	67.4	132.7	308.5
Male												
2002 .....	18.0	*	*	*	*	4.1	11.7	31.9	61.9	84.0	141.8	264.7
2001 .....	16.3	*	*	*	*	2.7	10.5	30.4	50.1	79.8	137.4	288.7
2000 .....	16.1	*	*	*	*	2.7	9.2	28.4	56.0	81.9	139.5	275.9
1999 .....	15.6	*	*	*	*	2.2	10.3	26.8	51.1	80.6	143.2	266.2
Female												
2002 .....	17.4	*	*	*	*	1.8	6.2	18.7	31.7	57.1	129.8	344.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Non-Hispanic black												
Female												
2001 .....	16.7	*	*	*	*	1.7	6.7	17.1	30.8	56.5	118.9	343.5
2000 .....	17.1	*	*	*	*	1.8	5.9	15.8	32.7	57.5	128.9	369.4
1999 .....	16.4	*	*	*	*	1.6	6.1	16.5	31.3	58.1	126.8	324.2
Hypertensive heart and renal disease (I13)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.8	23.5
2001 .....	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.3	6.8	23.3
2000 .....	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.3	7.0	24.3
1999 .....	1.2	*	*	*	*	0.1	0.3	0.4	1.1	2.6	8.9	29.1
Male												
2002 .....	0.9	*	*	*	*	*	0.2	0.5	1.2	2.4	7.0	23.1
2001 .....	0.8	*	*	*	*	*	0.3	0.5	1.0	2.5	6.8	24.2
2000 .....	0.8	*	*	*	*	*	0.2	0.5	0.9	2.5	7.2	27.8
1999 .....	1.0	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.3
Female												
2002 .....	1.2	*	*	*	*	*	0.1	0.3	0.8	2.0	6.7	23.7
2001 .....	1.1	*	*	*	*	*	0.2	0.2	0.9	2.1	6.8	22.9
2000 .....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.9
1999 .....	1.4	*	*	*	*	*	0.2	0.3	0.9	2.4	8.5	28.2
Hispanic												
Both sexes												
2002 .....	0.4	*	*	*	*	*	*	*	1.3	2.9	8.5	22.7
2001 .....	0.4	*	*	*	*	*	*	*	1.1	3.0	7.4	26.3
2000 .....	0.4	*	*	*	*	*	*	*	*	2.5	6.3	25.2
1999 .....	0.4	*	*	*	*	*	*	*	*	3.1	8.1	23.2
Male												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	8.2	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.5	*	*	*	*	*	*	*	*	*	8.7	21.6
2001 .....	0.5	*	*	*	*	*	*	*	*	*	7.3	24.2
2000 .....	0.4	*	*	*	*	*	*	*	*	*	*	26.7
1999 .....	0.5	*	*	*	*	*	*	*	*	*	8.7	20.8
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.1	6.7	23.5
2001 .....	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.7	23.1
2000 .....	1.1	*	*	*	*	0.1	0.2	0.4	0.9	2.2	7.0	24.2
1999 .....	1.3	*	*	*	*	0.1	0.3	0.4	1.1	2.5	8.9	29.3
Male												
2002 .....	0.9	*	*	*	*	*	0.2	0.6	1.2	2.3	6.9	22.9
2001 .....	0.9	*	*	*	*	*	0.3	0.5	1.0	2.4	6.7	23.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Hypertensive heart and renal disease (I13)</b>												
<b>Non-Hispanic <sup>4</sup></b>												
<b>Male</b>												
2000 .....	0.9	*	*	*	*	*	0.2	0.5	0.9	2.4	7.0	27.9
1999 .....	1.1	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.4
<b>Female</b>												
2002 .....	1.2	*	*	*	*	*	0.2	0.3	0.8	2.0	6.5	23.7
2001 .....	1.2	*	*	*	*	*	0.2	0.2	0.9	2.0	6.7	22.8
2000 .....	1.2	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.8
1999 .....	1.5	*	*	*	*	*	0.2	0.3	0.9	2.4	8.4	28.4
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	0.9	*	*	*	*	*	*	0.1	0.4	1.2	5.6	20.8
2001 .....	0.9	*	*	*	*	*	*	0.1	0.5	1.4	5.6	20.7
2000 .....	0.9	*	*	*	*	*	0.1	0.1	0.4	1.5	5.6	21.8
1999 .....	1.1	*	*	*	*	*	*	0.2	0.5	1.4	7.3	26.1
<b>Male</b>												
2002 .....	0.7	*	*	*	*	*	*	*	0.5	1.4	5.9	20.2
2001 .....	0.7	*	*	*	*	*	*	0.2	0.4	1.5	5.8	22.2
2000 .....	0.7	*	*	*	*	*	*	0.2	0.4	1.6	6.1	24.3
1999 .....	0.9	*	*	*	*	*	*	0.2	0.5	1.6	8.3	28.7
<b>Female</b>												
2002 .....	1.1	*	*	*	*	*	*	*	0.3	1.1	5.4	21.0
2001 .....	1.1	*	*	*	*	*	*	*	0.5	1.3	5.4	20.1
2000 .....	1.1	*	*	*	*	*	*	*	0.3	1.4	5.2	20.8
1999 .....	1.2	*	*	*	*	*	*	*	0.4	1.3	6.6	25.1
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	2.5	*	*	*	*	*	1.0	2.8	6.2	10.0	19.8	57.2
2001 .....	2.3	*	*	*	*	0.4	1.2	2.1	5.1	10.1	20.2	50.6
2000 .....	2.3	*	*	*	*	*	0.8	2.2	4.7	10.0	22.6	50.9
1999 .....	2.8	*	*	*	*	*	1.5	2.3	6.2	12.7	27.3	66.0
<b>Male</b>												
2002 .....	2.3	*	*	*	*	*	1.0	3.9	7.7	10.2	19.3	62.4
2001 .....	2.2	*	*	*	*	*	1.6	3.2	6.0	12.0	18.6	45.0
2000 .....	2.1	*	*	*	*	*	1.0	2.9	5.4	11.1	20.1	64.6
1999 .....	2.6	*	*	*	*	*	1.8	3.3	7.8	13.8	26.0	60.6
<b>Female</b>												
2002 .....	2.6	*	*	*	*	*	0.9	1.8	5.0	9.9	20.1	55.2
2001 .....	2.4	*	*	*	*	*	0.8	1.2	4.3	8.7	21.1	52.6
2000 .....	2.4	*	*	*	*	*	*	1.6	4.2	9.2	24.0	45.8
1999 .....	3.1	*	*	*	*	*	1.2	1.5	4.8	12.0	28.0	68.0
<b>Ischemic heart diseases (I20-I25)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	171.4	0.6	*	0.0	0.3	2.5	17.1	63.8	177.7	458.9	1,220.2	3,775.0
2001 .....	176.3	0.6	*	0.0	0.4	2.8	16.7	64.0	184.0	477.6	1,271.3	3,958.2
2000 .....	183.1	0.8	*	*	0.3	2.3	16.7	66.1	196.7	505.1	1,324.2	4,182.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
All origins <sup>3</sup>												
Both sexes												
1999 .....	189.8	0.7	*	0.1	0.4	2.5	17.2	67.7	203.5	536.6	1,384.8	4,318.1
Male												
2002 .....	178.4	*	*	*	0.4	3.7	26.1	99.4	263.3	640.1	1,598.9	4,205.3
2001 .....	181.7	*	*	*	0.6	4.1	25.5	99.5	271.6	663.7	1,663.5	4,395.8
2000 .....	188.7	*	*	*	0.4	3.5	25.8	103.5	290.2	705.2	1,736.8	4,701.7
1999 .....	195.4	*	*	*	0.5	3.7	26.6	106.2	299.3	749.8	1,813.7	4,837.5
Female												
2002 .....	164.7	*	*	*	0.2	1.3	8.2	29.5	98.5	308.1	968.9	3,588.3
2001 .....	171.2	*	*	*	0.2	1.5	8.1	29.7	103.1	323.4	1,013.6	3,775.1
2000 .....	177.6	*	*	*	0.2	1.2	7.6	29.9	110.4	340.3	1,055.2	3,971.0
1999 .....	184.5	*	*	*	0.2	1.4	7.9	30.7	115.1	362.0	1,106.3	4,108.8
Hispanic												
Both sexes												
2002 .....	54.0	*	*	*	*	1.2	7.6	38.0	135.9	382.0	1,010.8	3,182.9
2001 .....	55.9	*	*	*	*	1.6	6.8	38.7	130.2	400.9	1,099.1	3,620.2
2000 .....	55.9	*	*	*	*	1.1	7.6	39.5	143.6	405.3	1,111.3	3,697.2
1999 .....	58.7	*	*	*	*	1.4	7.4	44.8	150.4	433.5	1,178.0	3,876.2
Male												
2002 .....	56.5	*	*	*	*	1.9	11.4	58.0	204.0	530.4	1,271.7	3,374.9
2001 .....	57.9	*	*	*	*	2.4	10.4	60.3	191.9	559.9	1,393.3	3,828.2
2000 .....	58.2	*	*	*	*	1.6	12.1	62.7	214.8	561.5	1,420.0	3,865.6
1999 .....	60.8	*	*	*	*	2.1	12.2	71.3	218.3	608.5	1,495.9	3,987.7
Female												
2002 .....	51.4	*	*	*	*	*	3.5	18.1	74.6	263.9	832.2	3,087.8
2001 .....	53.7	*	*	*	*	*	2.8	17.4	75.4	275.4	900.3	3,518.6
2000 .....	53.6	*	*	*	*	*	2.8	16.8	80.6	282.5	904.0	3,614.6
1999 .....	56.6	*	*	*	*	*	2.3	19.0	90.5	296.4	966.5	3,822.4
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	189.0	*	*	*	0.3	2.8	18.4	66.1	180.1	462.1	1,226.5	3,787.0
2001 .....	193.6	0.7	*	*	0.4	3.1	18.0	66.1	187.3	480.9	1,274.9	3,959.4
2000 .....	200.5	0.8	*	*	0.3	2.6	17.7	68.1	199.5	509.0	1,328.8	4,187.0
1999 .....	207.1	0.8	*	0.1	0.4	2.7	18.3	69.4	206.3	540.3	1,388.1	4,319.6
Male												
2002 .....	197.6	*	*	*	0.5	4.1	28.3	103.1	266.5	644.3	1,609.7	4,228.4
2001 .....	200.4	*	*	*	0.6	4.5	27.5	102.8	276.1	667.7	1,670.1	4,408.5
2000 .....	207.6	*	*	*	0.4	3.9	27.4	106.7	294.0	710.4	1,744.6	4,721.1
1999 .....	214.1	*	*	*	0.5	4.0	28.3	108.8	303.4	754.2	1,820.2	4,854.5
Female												
2002 .....	180.7	*	*	*	0.2	1.5	8.8	30.5	100.0	309.9	972.5	3,596.5
2001 .....	187.2	*	*	*	0.2	1.6	8.7	30.8	105.0	325.6	1,015.5	3,772.7
2000 .....	193.8	*	*	*	0.2	1.3	8.2	31.0	112.1	342.8	1,057.9	3,971.0
1999 .....	200.6	*	*	*	0.3	1.5	8.5	31.5	116.6	364.6	1,107.8	4,105.4
Non-Hispanic white												
Both sexes												
2002 .....	208.0	*	*	*	0.3	2.6	17.9	62.8	171.2	450.8	1,223.1	3,848.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Non-Hispanic white												
Both sexes												
2002 .....	212.6	*	*	*	0.3	2.9	17.3	62.6	177.8	467.8	1,271.3	4,022.8
2001 .....	220.2	*	*	*	0.3	2.4	16.9	64.6	189.5	496.4	1,324.5	4,261.3
2000 .....	227.2	*	*	*	0.3	2.5	17.3	65.8	196.4	527.7	1,386.5	4,390.2
Male												
2002 .....	217.2	*	*	*	0.4	4.0	28.0	100.0	256.8	636.0	1,618.3	4,324.5
2001 .....	219.6	*	*	*	0.5	4.3	26.9	99.5	266.3	656.5	1,678.7	4,506.9
2000 .....	227.6	*	*	*	0.4	3.7	26.8	103.5	283.5	701.7	1,752.3	4,833.4
1999 .....	234.2	*	*	*	0.5	3.6	27.2	105.1	292.3	745.2	1,834.9	4,954.1
Female												
2002 .....	199.1	*	*	*	0.2	1.3	7.8	26.1	90.2	293.0	958.3	3,642.5
2001 .....	205.8	*	*	*	0.2	1.4	7.6	26.4	94.2	307.7	1,001.1	3,821.4
2000 .....	213.1	*	*	*	0.2	1.0	6.9	26.4	100.9	323.6	1,043.3	4,030.7
1999 .....	220.5	*	*	*	*	1.4	7.4	27.2	106.1	345.6	1,092.9	4,165.3
Non-Hispanic black												
Both sexes												
2002 .....	137.0	*	*	*	0.5	4.1	25.8	103.6	292.4	643.9	1,433.9	3,481.1
2001 .....	141.2	*	*	*	0.9	4.8	26.4	104.6	302.1	681.4	1,487.0	3,628.7
2000 .....	144.4	*	*	*	0.4	4.4	26.1	107.7	318.9	710.1	1,534.5	3,701.7
1999 .....	147.9	*	*	*	0.8	4.1	28.2	110.8	324.7	741.6	1,559.0	3,817.9
Male												
2002 .....	139.5	*	*	*	*	5.3	36.2	146.9	413.2	850.8	1,762.3	3,572.8
2001 .....	143.2	*	*	*	1.2	6.8	37.1	149.0	418.0	903.1	1,834.9	3,791.8
2000 .....	145.7	*	*	*	*	6.1	36.3	152.2	444.2	916.5	1,887.6	3,911.2
1999 .....	150.0	*	*	*	0.9	6.0	40.3	160.0	458.4	967.5	1,873.3	4,082.5
Female												
2002 .....	134.8	*	*	*	*	2.9	16.7	66.5	195.4	497.1	1,248.1	3,446.0
2001 .....	139.4	*	*	*	*	2.9	16.9	66.5	208.9	525.0	1,291.6	3,568.3
2000 .....	143.2	*	*	*	*	2.9	17.1	69.4	218.1	565.0	1,335.7	3,623.9
1999 .....	146.0	*	*	*	0.7	2.4	17.4	68.7	217.5	582.3	1,383.0	3,719.0
Acute myocardial infarction (I21-I22)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	62.3	*	*	*	0.2	1.0	7.0	26.5	74.9	186.1	444.4	1,191.3
2001 .....	64.9	*	*	*	0.2	1.2	7.1	27.6	78.4	193.2	471.2	1,265.4
2000 .....	68.5	0.5	*	*	0.1	0.9	6.9	28.9	86.6	209.1	503.3	1,338.5
1999 .....	71.5	*	*	*	0.2	1.1	7.2	29.6	90.0	225.0	529.3	1,379.3
Male												
2002 .....	66.2	*	*	*	0.2	1.4	10.5	40.6	109.2	256.2	569.2	1,366.4
2001 .....	68.6	*	*	*	0.2	1.7	10.6	42.4	114.0	264.7	604.1	1,461.3
2000 .....	72.7	*	*	*	0.1	1.3	10.6	44.7	125.7	286.9	648.3	1,568.3
1999 .....	76.1	*	*	*	0.2	1.5	11.0	46.3	131.8	310.0	684.4	1,622.1
Female												
2002 .....	58.4	*	*	*	0.1	0.5	3.6	12.9	43.2	127.7	361.5	1,115.3
2001 .....	61.3	*	*	*	*	0.7	3.5	13.3	45.4	134.0	383.9	1,183.5
2000 .....	64.6	*	*	*	0.1	0.5	3.3	13.6	50.5	145.1	408.7	1,244.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
All origins <sup>3</sup>												
Female												
1999 .....	67.0	*	*	*	0.1	0.6	3.4	13.5	51.5	155.4	428.6	1,281.5
Hispanic												
Both sexes												
2002 .....	19.7	*	*	*	*	0.5	3.0	14.2	55.1	146.2	364.3	1,048.2
2001 .....	20.0	*	*	*	*	0.7	2.5	14.8	49.5	149.8	394.9	1,185.5
2000 .....	20.1	*	*	*	*	0.5	3.0	15.5	55.0	154.3	404.7	1,177.1
1999 .....	21.1	*	*	*	*	0.5	2.9	17.8	56.1	164.0	433.7	1,227.8
Male												
2002 .....	20.6	*	*	*	*	0.7	4.2	21.0	80.8	199.3	460.0	1,107.7
2001 .....	20.9	*	*	*	*	1.0	3.9	22.7	69.4	206.9	488.7	1,331.3
2000 .....	21.1	*	*	*	*	0.7	4.7	23.4	80.0	207.6	514.8	1,273.8
1999 .....	22.0	*	*	*	*	0.8	4.6	26.7	79.0	225.8	558.9	1,276.8
Female												
2002 .....	18.7	*	*	*	*	*	1.8	7.4	32.1	103.9	298.8	1,018.8
2001 .....	19.1	*	*	*	*	*	1.1	7.0	31.8	104.7	331.5	1,114.2
2000 .....	19.1	*	*	*	*	*	1.2	7.8	32.9	112.3	330.8	1,129.7
1999 .....	20.1	*	*	*	*	*	1.1	9.1	35.9	115.5	350.4	1,204.1
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	68.7	*	*	*	0.2	1.1	7.6	27.7	76.3	188.2	447.3	1,194.0
2001 .....	71.4	*	*	*	0.2	1.3	7.6	28.8	80.3	195.6	473.6	1,265.3
2000 .....	75.3	*	*	*	0.1	1.0	7.4	30.0	88.6	211.9	506.3	1,340.8
1999 .....	78.2	*	*	*	0.2	1.2	7.7	30.5	92.2	227.9	531.9	1,381.0
Male												
2002 .....	73.5	*	*	*	0.2	1.6	11.5	42.6	111.3	259.2	573.4	1,374.2
2001 .....	75.9	*	*	*	0.2	1.8	11.6	44.2	117.0	267.7	607.9	1,463.9
2000 .....	80.3	*	*	*	0.2	1.5	11.4	46.5	128.7	290.9	652.6	1,576.2
1999 .....	83.8	*	*	*	0.2	1.7	11.8	47.8	135.1	313.8	687.7	1,631.5
Female												
2002 .....	64.1	*	*	*	*	0.6	3.8	13.3	44.0	128.9	363.8	1,116.3
2001 .....	67.1	*	*	*	*	0.8	3.8	13.9	46.3	135.7	385.4	1,182.7
2000 .....	70.6	*	*	*	0.1	0.6	3.5	14.1	51.6	146.8	411.0	1,245.6
1999 .....	72.9	*	*	*	*	0.7	3.7	13.9	52.5	157.3	430.7	1,280.8
Non-Hispanic white												
Both sexes												
2002 .....	75.5	*	*	*	0.1	1.1	7.5	27.0	74.7	186.1	445.9	1,202.8
2001 .....	78.4	*	*	*	0.1	1.2	7.5	28.1	78.7	193.0	473.0	1,276.8
2000 .....	82.7	*	*	*	0.1	0.9	7.3	29.3	86.9	209.3	505.3	1,355.3
1999 .....	86.0	*	*	*	0.1	1.1	7.5	30.0	90.7	226.1	532.4	1,394.2
Male												
2002 .....	81.2	*	*	*	0.2	1.6	11.5	42.5	110.8	260.2	575.8	1,393.0
2001 .....	83.7	*	*	*	0.2	1.7	11.6	44.3	116.8	267.7	612.5	1,483.2
2000 .....	88.5	*	*	*	0.2	1.4	11.4	46.7	128.0	291.0	656.7	1,600.7
1999 .....	92.4	*	*	*	0.2	1.5	11.8	48.1	134.6	315.0	695.7	1,653.8
Female												
2002 .....	70.1	*	*	*	*	0.5	3.6	11.8	40.6	122.9	358.8	1,120.7

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Non-Hispanic white												
Female												
2001 .....	73.4	*	*	*	*	0.8	3.4	12.2	42.7	129.6	380.6	1,190.9
2000 .....	77.2	*	*	*	*	0.5	3.1	12.3	48.2	140.5	405.9	1,256.3
1999 .....	79.8	*	*	*	*	0.7	3.3	12.3	49.3	151.8	425.5	1,290.7
Non-Hispanic black												
Both sexes												
2002 .....	50.2	*	*	*	*	1.7	10.0	39.0	109.1	242.6	523.0	1,212.9
2001 .....	51.6	*	*	*	0.4	1.9	10.2	40.2	111.6	256.0	543.7	1,247.0
2000 .....	54.0	*	*	*	*	1.8	10.1	41.9	122.3	273.0	576.6	1,281.8
1999 .....	55.1	*	*	*	*	1.7	10.5	42.0	123.2	282.4	589.9	1,328.6
Male												
2002 .....	50.1	*	*	*	*	2.2	13.7	53.4	144.5	305.4	631.3	1,296.3
2001 .....	50.9	*	*	*	*	2.5	13.5	53.4	146.1	320.8	646.6	1,389.4
2000 .....	53.7	*	*	*	*	2.2	13.6	56.3	163.1	341.2	693.3	1,426.6
1999 .....	55.2	*	*	*	*	2.5	14.5	58.1	168.7	357.0	691.4	1,509.4
Female												
2002 .....	50.4	*	*	*	*	1.3	6.7	26.7	80.7	198.0	461.8	1,181.0
2001 .....	52.3	*	*	*	*	1.3	7.4	28.8	83.8	210.3	485.9	1,194.3
2000 .....	54.2	*	*	*	*	1.4	7.0	29.6	89.5	225.0	510.9	1,228.0
1999 .....	54.9	*	*	*	*	1.0	7.0	28.2	86.8	229.7	533.1	1,261.0
Other acute ischemic heart diseases (I24)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.2	*	*	*	*	*	0.2	0.9	1.7	3.5	7.1	21.1
2001 .....	1.2	*	*	*	*	*	0.2	0.8	1.8	3.5	6.9	21.5
2000 .....	1.2	*	*	*	*	*	0.2	0.7	2.0	3.5	7.3	21.9
1999 .....	1.2	*	*	*	*	*	0.2	0.7	1.9	3.7	7.0	21.9
Male												
2002 .....	1.2	*	*	*	*	*	0.2	1.3	2.5	4.7	8.7	21.4
2001 .....	1.2	*	*	*	*	*	0.3	1.2	2.6	4.7	8.5	20.7
2000 .....	1.3	*	*	*	*	*	0.3	1.2	2.9	4.8	9.1	20.7
1999 .....	1.2	*	*	*	*	*	0.3	1.0	2.8	4.9	8.8	22.3
Female												
2002 .....	1.1	*	*	*	*	*	0.1	0.5	1.0	2.4	5.9	21.0
2001 .....	1.1	*	*	*	*	*	0.1	0.4	1.1	2.5	5.9	21.8
2000 .....	1.1	*	*	*	*	*	0.1	0.3	1.1	2.4	6.1	22.4
1999 .....	1.1	*	*	*	*	*	*	0.4	1.1	2.6	5.8	21.7
Hispanic												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Hispanic												
Male												
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.3	*	*	*	*	*	0.2	0.9	1.8	3.6	7.3	21.8
2001 .....	1.3	*	*	*	*	*	0.2	0.8	1.9	3.6	7.1	22.0
2000 .....	1.3	*	*	*	*	*	0.2	0.8	2.1	3.6	7.5	22.2
1999 .....	1.3	*	*	*	*	*	0.2	0.7	2.0	3.8	7.2	22.2
Male												
2002 .....	1.4	*	*	*	*	*	0.3	1.4	2.6	4.9	9.0	22.1
2001 .....	1.4	*	*	*	*	*	0.3	1.3	2.7	4.8	8.7	20.9
2000 .....	1.4	*	*	*	*	*	0.3	1.3	3.1	4.9	9.3	21.0
1999 .....	1.4	*	*	*	*	*	0.3	1.1	2.9	5.2	9.0	22.9
Female												
2002 .....	1.3	*	*	*	*	*	0.1	0.5	1.1	2.5	6.1	21.6
2001 .....	1.3	*	*	*	*	*	0.1	0.4	1.1	2.6	6.0	22.4
2000 .....	1.2	*	*	*	*	*	0.1	0.3	1.2	2.5	6.3	22.7
1999 .....	1.2	*	*	*	*	*	*	0.4	1.1	2.7	6.0	21.9
Non-Hispanic white												
Both sexes												
2002 .....	1.4	*	*	*	*	*	0.2	0.8	1.7	3.5	7.2	22.0
2001 .....	1.4	*	*	*	*	*	0.2	0.8	1.7	3.4	7.0	22.0
2000 .....	1.4	*	*	*	*	*	0.2	0.7	1.9	3.4	7.3	22.5
1999 .....	1.4	*	*	*	*	*	0.2	0.6	1.8	3.6	7.0	22.7
Male												
2002 .....	1.5	*	*	*	*	*	0.3	1.2	2.5	4.6	9.0	22.7
2001 .....	1.4	*	*	*	*	*	0.3	1.2	2.5	4.5	8.7	20.9
2000 .....	1.5	*	*	*	*	*	0.3	1.2	2.8	4.6	9.0	21.3
1999 .....	1.4	*	*	*	*	*	0.2	0.9	2.6	4.9	8.7	23.6
Female												
2002 .....	1.4	*	*	*	*	*	*	0.5	1.0	2.5	6.0	21.8
2001 .....	1.4	*	*	*	*	*	*	0.3	1.0	2.5	5.9	22.4
2000 .....	1.3	*	*	*	*	*	*	0.3	1.0	2.4	6.1	23.0
1999 .....	1.3	*	*	*	*	*	*	0.3	1.1	2.6	5.8	22.4
Non-Hispanic black												
Both sexes												
2002 .....	1.2	*	*	*	*	*	*	1.7	3.0	6.1	9.4	21.5
2001 .....	1.3	*	*	*	*	*	0.5	1.6	4.0	5.8	8.5	24.3
2000 .....	1.3	*	*	*	*	*	0.4	1.5	4.3	5.6	10.8	21.9
1999 .....	1.2	*	*	*	*	*	0.5	1.9	3.6	5.8	10.9	17.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Non-Hispanic black												
Male												
2002 .....	1.4	*	*	*	*	*	*	2.8	4.4	9.4	11.7	21.9
2001 .....	1.5	*	*	*	*	*	*	2.5	5.6	8.8	10.5	24.3
2000 .....	1.5	*	*	*	*	*	*	2.3	6.3	8.7	14.2	*
1999 .....	1.5	*	*	*	*	*	0.8	2.7	6.1	8.4	12.8	*
Female												
2002 .....	1.1	*	*	*	*	*	*	0.8	2.0	3.7	8.1	21.3
2001 .....	1.1	*	*	*	*	*	*	*	2.6	3.8	7.4	24.4
2000 .....	1.1	*	*	*	*	*	*	*	2.7	3.5	8.9	22.2
1999 .....	1.0	*	*	*	*	*	*	1.1	1.6	3.9	9.8	16.4
Other forms of chronic ischemic heart disease (I20,I25)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	108.0	*	*	*	0.2	1.4	9.9	36.5	101.0	269.4	768.8	2,562.5
2001 .....	110.3	*	*	*	0.2	1.6	9.5	35.6	103.8	280.9	793.1	2,671.3
2000 .....	113.3	*	*	*	0.2	1.4	9.5	36.4	108.1	292.4	813.6	2,822.1
1999 .....	117.2	*	*	*	0.2	1.4	9.8	37.5	111.5	308.0	848.5	2,917.0
Male												
2002 .....	110.9	*	*	*	0.2	2.2	15.4	57.5	151.5	379.3	1,021.0	2,817.4
2001 .....	111.9	*	*	*	0.3	2.4	14.6	55.9	154.9	394.3	1,050.9	2,913.8
2000 .....	114.8	*	*	*	0.2	2.1	14.8	57.7	161.6	413.5	1,079.4	3,112.7
1999 .....	118.0	*	*	*	0.3	2.1	15.3	59.0	164.7	434.9	1,120.5	3,193.1
Female												
2002 .....	105.2	*	*	*	*	0.7	4.4	16.2	54.3	178.0	601.4	2,451.9
2001 .....	108.8	*	*	*	0.1	0.7	4.4	16.0	56.7	186.9	623.7	2,569.8
2000 .....	111.9	*	*	*	*	0.6	4.3	16.0	58.8	192.8	640.3	2,703.8
1999 .....	116.4	*	*	*	0.1	0.7	4.3	16.7	62.6	204.0	671.8	2,805.7
Hispanic												
Both sexes												
2002 .....	34.2	*	*	*	*	0.7	4.5	23.6	80.1	234.6	644.0	2,129.7
2001 .....	35.7	*	*	*	*	0.9	4.2	23.8	80.0	249.4	701.7	2,426.6
2000 .....	35.6	*	*	*	*	0.6	4.6	23.9	88.0	249.7	703.2	2,508.8
1999 .....	37.5	*	*	*	*	0.9	4.5	26.8	93.5	268.8	741.8	2,640.7
Male												
2002 .....	35.7	*	*	*	*	1.2	7.2	36.6	122.3	329.1	808.8	2,262.2
2001 .....	36.8	*	*	*	*	1.4	6.5	37.5	121.4	350.2	901.5	2,479.8
2000 .....	36.9	*	*	*	*	0.9	7.4	39.1	134.2	351.6	900.8	2,577.7
1999 .....	38.7	*	*	*	*	1.3	7.6	44.3	138.5	381.8	932.8	2,706.7
Female												
2002 .....	32.6	*	*	*	*	*	1.6	10.5	42.2	159.4	531.2	2,064.1
2001 .....	34.5	*	*	*	*	*	1.7	10.3	43.3	169.9	566.7	2,400.7
2000 .....	34.2	*	*	*	*	*	1.7	8.9	47.2	169.6	570.5	2,475.0
1999 .....	36.3	*	*	*	*	*	1.3	9.7	53.8	180.2	614.8	2,608.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	118.9	*	*	*	0.2	1.6	10.6	37.5	102.0	270.3	771.9	2,571.2
2001 .....	120.9	*	*	*	0.2	1.7	10.2	36.5	105.1	281.6	794.2	2,672.2
2000 .....	123.9	*	*	*	0.2	1.5	10.1	37.3	108.8	293.5	815.0	2,824.0
1999 .....	127.6	*	*	*	0.2	1.5	10.4	38.1	112.1	308.6	849.1	2,916.3
Male												
2002 .....	122.7	*	*	*	0.3	2.4	16.6	59.2	152.6	380.2	1,027.3	2,832.2
2001 .....	123.1	*	*	*	0.3	2.6	15.7	57.3	156.4	395.1	1,053.5	2,923.7
2000 .....	126.0	*	*	*	0.2	2.3	15.7	58.9	162.3	414.7	1,082.7	3,123.9
1999 .....	128.9	*	*	*	0.3	2.3	16.2	59.9	165.4	435.2	1,123.5	3,200.1
Female												
2002 .....	115.4	*	*	*	*	0.8	4.8	16.7	55.0	178.4	602.6	2,458.6
2001 .....	118.8	*	*	*	0.1	0.8	4.8	16.5	57.5	187.3	624.1	2,567.6
2000 .....	122.0	*	*	*	*	0.7	4.6	16.5	59.3	193.5	640.6	2,702.7
1999 .....	126.4	*	*	*	0.1	0.8	4.7	17.3	62.9	204.6	671.1	2,802.7
Non-Hispanic white												
Both sexes												
2002 .....	131.0	*	*	*	0.2	1.5	10.2	34.9	94.8	261.3	770.0	2,623.4
2001 .....	132.7	*	*	*	0.2	1.6	9.6	33.8	97.4	271.4	791.2	2,724.1
2000 .....	136.1	*	*	*	0.2	1.4	9.4	34.6	100.7	283.7	811.9	2,883.6
1999 .....	139.9	*	*	*	0.2	1.4	9.6	35.2	103.9	297.9	847.2	2,973.2
Male												
2002 .....	134.5	*	*	*	0.2	2.3	16.2	56.4	143.5	371.2	1,033.6	2,908.8
2001 .....	134.4	*	*	*	0.3	2.5	15.0	54.1	147.0	384.3	1,057.5	3,002.8
2000 .....	137.7	*	*	*	0.2	2.2	15.1	55.7	152.7	406.1	1,086.5	3,211.3
1999 .....	140.4	*	*	*	0.2	2.1	15.2	56.1	155.1	425.3	1,130.5	3,276.7
Female												
2002 .....	127.7	*	*	*	*	0.7	4.2	13.8	48.7	167.6	593.4	2,500.1
2001 .....	131.1	*	*	*	*	0.7	4.1	13.9	50.5	175.6	614.6	2,608.1
2000 .....	134.6	*	*	*	*	0.5	3.8	13.8	51.8	180.6	631.3	2,751.4
1999 .....	139.3	*	*	*	*	0.7	4.0	14.6	55.7	191.3	661.7	2,852.2
Non-Hispanic black												
Both sexes												
2002 .....	85.5	*	*	*	*	2.2	15.5	62.9	180.3	395.2	901.5	2,246.7
2001 .....	88.2	*	*	*	0.5	2.9	15.6	62.9	186.6	419.6	934.8	2,357.3
2000 .....	89.1	*	*	*	*	2.5	15.6	64.3	192.3	431.5	947.1	2,398.0
1999 .....	91.6	*	*	*	0.4	2.4	17.2	66.9	197.9	453.5	958.2	2,471.9
Male												
2002 .....	88.0	*	*	*	*	3.0	22.2	90.8	264.4	536.0	1,119.3	2,254.6
2001 .....	90.8	*	*	*	*	4.3	22.9	93.1	266.3	573.5	1,177.8	2,378.1
2000 .....	90.4	*	*	*	*	3.7	22.1	93.6	274.7	566.6	1,180.1	2,463.5
1999 .....	93.3	*	*	*	*	3.5	25.1	99.2	283.6	602.1	1,169.2	2,552.9
Female												
2002 .....	83.3	*	*	*	*	1.6	9.6	39.0	112.8	295.4	778.3	2,243.7
2001 .....	86.0	*	*	*	*	1.6	9.2	36.9	122.4	310.9	798.4	2,349.6
2000 .....	88.0	*	*	*	*	1.5	10.0	39.1	125.9	336.5	815.9	2,373.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Non-Hispanic black												
Female												
1999 .....	90.1	*	*	*	*	1.3	10.2	39.4	129.2	348.6	840.0	2,441.6
Atherosclerotic cardiovascular disease, so described (I25.0)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	23.6	*	*	*	*	0.5	3.4	13.4	33.1	64.7	146.2	474.5
2001 .....	23.9	*	*	*	0.1	0.5	3.2	12.9	33.3	65.6	152.7	493.6
2000 .....	24.7	*	*	*	*	0.4	3.0	13.2	34.0	68.3	158.1	534.6
1999 .....	25.8	*	*	*	*	0.4	3.3	13.0	35.3	72.7	166.2	563.4
Male												
2002 .....	25.4	*	*	*	*	0.7	5.4	21.5	51.2	91.8	185.3	491.8
2001 .....	25.3	*	*	*	*	0.7	5.0	21.0	51.4	92.4	193.9	502.0
2000 .....	26.0	*	*	*	*	0.6	4.9	21.5	52.6	97.2	199.3	550.8
1999 .....	26.8	*	*	*	*	0.6	5.3	20.9	53.7	103.0	207.0	578.6
Female												
2002 .....	21.9	*	*	*	*	0.2	1.3	5.5	16.2	42.1	120.2	467.1
2001 .....	22.5	*	*	*	*	0.2	1.4	5.1	16.7	43.3	125.5	490.1
2000 .....	23.5	*	*	*	*	0.1	1.2	5.2	16.8	44.6	131.3	528.0
1999 .....	24.8	*	*	*	*	0.2	1.3	5.3	18.3	47.9	139.8	557.2
Hispanic												
Both sexes												
2002 .....	8.2	*	*	*	*	0.4	1.9	8.9	29.7	61.1	124.1	358.6
2001 .....	8.5	*	*	*	*	0.4	1.5	10.1	29.7	65.9	132.5	417.0
2000 .....	8.4	*	*	*	*	*	1.6	9.5	30.5	65.7	128.2	452.5
1999 .....	9.0	*	*	*	*	*	1.5	10.4	33.0	72.1	147.5	469.4
Male												
2002 .....	9.7	*	*	*	*	0.6	2.9	14.8	49.3	90.8	153.9	415.0
2001 .....	10.1	*	*	*	*	0.6	2.2	17.1	47.9	100.2	181.1	440.0
2000 .....	9.9	*	*	*	*	*	2.7	16.6	50.5	92.4	180.9	493.8
1999 .....	10.6	*	*	*	*	*	2.5	17.8	50.0	115.8	194.0	485.3
Female												
2002 .....	6.5	*	*	*	*	*	*	3.1	12.0	37.6	103.7	330.7
2001 .....	6.8	*	*	*	*	*	*	3.2	13.5	38.7	99.7	405.8
2000 .....	6.8	*	*	*	*	*	*	2.6	12.8	44.6	92.8	432.3
1999 .....	7.4	*	*	*	*	*	*	3.2	18.0	37.9	116.6	461.7
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	25.9	*	*	*	*	0.5	3.6	13.7	32.9	64.3	146.4	477.8
2001 .....	26.0	*	*	*	*	0.5	3.5	13.0	33.3	65.0	152.6	494.8
2000 .....	26.9	*	*	*	*	0.4	3.2	13.4	33.8	67.9	158.3	535.9
1999 .....	27.9	*	*	*	*	0.4	3.5	13.1	35.0	72.0	165.7	564.9
Male												
2002 .....	27.8	*	*	*	*	0.7	5.8	22.0	50.7	91.0	185.6	493.3
2001 .....	27.5	*	*	*	*	0.7	5.4	21.2	51.1	91.0	193.2	503.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Non-Hispanic <sup>4</sup>												
Male												
2000 .....	28.2	*	*	*	*	0.7	5.1	21.7	52.1	96.4	198.5	551.0
1999 .....	28.9	*	*	*	*	0.6	5.6	21.0	53.3	101.1	205.6	580.2
Female												
2002 .....	24.0	*	*	*	*	0.2	1.4	5.7	16.5	42.0	120.5	471.1
2001 .....	24.6	*	*	*	*	0.3	1.5	5.2	16.8	43.3	126.0	491.4
2000 .....	25.6	*	*	*	*	0.2	1.3	5.4	16.9	44.4	132.2	529.8
1999 .....	27.0	*	*	*	*	0.2	1.4	5.4	18.2	48.2	139.8	558.8
Non-Hispanic white												
Both sexes												
2002 .....	26.8	*	*	*	*	0.4	3.3	12.4	28.9	57.2	138.2	473.5
2001 .....	26.8	*	*	*	*	0.4	3.2	11.6	28.4	57.1	144.2	490.7
2000 .....	27.6	*	*	*	*	0.4	2.9	11.9	28.9	60.2	149.0	530.2
1999 .....	28.6	*	*	*	*	0.4	3.1	11.3	29.8	63.6	156.2	558.8
Male												
2002 .....	28.2	*	*	*	*	0.7	5.4	20.3	44.8	81.3	174.4	489.2
2001 .....	27.8	*	*	*	*	0.6	5.0	19.1	44.3	80.4	182.9	499.4
2000 .....	28.4	*	*	*	*	0.6	4.8	19.6	45.2	86.3	186.1	544.7
1999 .....	29.0	*	*	*	*	0.6	5.2	18.4	46.0	90.0	193.5	572.8
Female												
2002 .....	25.3	*	*	*	*	0.2	1.1	4.6	13.8	36.6	113.9	466.8
2001 .....	25.8	*	*	*	*	0.2	1.3	4.1	13.3	37.3	118.6	487.0
2000 .....	26.9	*	*	*	*	*	1.1	4.3	13.4	38.2	124.6	524.4
1999 .....	28.2	*	*	*	*	0.2	1.1	4.2	14.6	41.6	131.8	553.2
Non-Hispanic black												
Both sexes												
2002 .....	27.7	*	*	*	*	0.8	6.0	25.7	71.9	137.2	261.1	580.4
2001 .....	28.5	*	*	*	*	0.7	5.8	25.7	78.1	144.8	266.7	595.9
2000 .....	29.3	*	*	*	*	0.7	5.3	27.3	78.9	147.2	280.2	652.2
1999 .....	30.7	*	*	*	*	0.8	6.5	28.1	82.0	158.7	291.0	670.3
Male												
2002 .....	31.9	*	*	*	*	1.1	8.9	38.8	111.8	198.7	358.4	603.9
2001 .....	32.3	*	*	*	*	1.0	8.8	39.9	118.1	209.4	352.3	609.8
2000 .....	33.2	*	*	*	*	1.1	7.9	42.3	120.8	209.3	371.5	711.6
1999 .....	34.5	*	*	*	*	1.1	9.8	43.6	123.7	228.2	378.4	714.3
Female												
2002 .....	24.0	*	*	*	*	*	3.4	14.4	39.8	93.6	206.1	571.5
2001 .....	25.0	*	*	*	*	*	3.2	13.6	45.9	99.3	218.7	590.7
2000 .....	25.8	*	*	*	*	*	2.9	14.4	45.1	103.4	228.9	630.2
1999 .....	27.1	*	*	*	*	*	3.6	14.7	48.5	109.7	242.1	653.8
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	84.4	*	*	*	0.1	1.0	6.5	23.1	68.0	204.7	622.6	2,088.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)												
All origins <sup>3</sup>												
Both sexes												
2001 .....	86.4	*	*	*	0.2	1.1	6.2	22.7	70.5	215.3	640.5	2,177.7
2000 .....	88.6	*	*	*	0.1	1.0	6.5	23.2	74.1	224.1	655.5	2,287.5
1999 .....	91.4	*	*	*	0.2	1.0	6.5	24.5	76.3	235.2	682.3	2,353.6
Male												
2002 .....	85.5	*	*	*	0.2	1.5	10.0	35.9	100.3	287.5	835.7	2,325.6
2001 .....	86.5	*	*	*	0.2	1.8	9.5	34.9	103.5	301.9	856.9	2,411.8
2000 .....	88.8	*	*	*	0.2	1.5	9.9	36.2	109.0	316.3	880.1	2,561.9
1999 .....	91.2	*	*	*	0.2	1.5	10.0	38.1	111.0	331.9	913.6	2,614.4
Female												
2002 .....	83.3	*	*	*	*	0.5	3.1	10.7	38.0	135.9	481.1	1,984.9
2001 .....	86.2	*	*	*	0.1	0.5	3.0	11.0	40.0	143.6	498.2	2,079.8
2000 .....	88.4	*	*	*	*	0.5	3.0	10.7	42.0	148.2	509.0	2,175.8
1999 .....	91.6	*	*	*	0.1	0.5	3.0	11.4	44.3	156.1	532.1	2,248.4
Hispanic												
Both sexes												
2002 .....	26.0	*	*	*	*	0.4	2.7	14.6	50.4	173.4	519.9	1,771.1
2001 .....	27.2	*	*	*	*	0.6	2.7	13.7	50.4	183.6	569.1	2,009.6
2000 .....	27.2	*	*	*	*	0.5	3.0	14.3	57.5	184.0	575.0	2,056.3
1999 .....	28.5	*	*	*	*	0.6	3.0	16.4	60.5	196.7	594.3	2,171.4
Male												
2002 .....	25.9	*	*	*	*	0.6	4.3	21.8	73.0	238.3	654.9	1,847.2
2001 .....	26.7	*	*	*	*	0.9	4.3	20.4	73.6	250.0	720.4	2,039.8
2000 .....	27.0	*	*	*	*	0.7	4.7	22.6	83.6	259.1	719.8	2,084.0
1999 .....	28.1	*	*	*	*	1.0	5.1	26.5	88.5	266.0	738.8	2,221.4
Female												
2002 .....	26.1	*	*	*	*	*	0.9	7.4	30.2	121.8	427.5	1,733.3
2001 .....	27.7	*	*	*	*	*	1.0	7.1	29.8	131.2	466.9	1,994.8
2000 .....	27.5	*	*	*	*	*	1.2	6.2	34.4	124.9	477.7	2,042.7
1999 .....	28.9	*	*	*	*	*	0.8	6.5	35.8	142.3	498.2	2,147.2
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	93.1	*	*	*	0.1	1.1	7.1	23.8	69.0	206.0	625.4	2,093.4
2001 .....	94.9	*	*	*	0.2	1.2	6.7	23.5	71.8	216.7	641.6	2,177.4
2000 .....	97.0	*	*	*	0.1	1.1	6.9	23.9	75.0	225.6	656.7	2,288.1
1999 .....	99.7	*	*	*	0.2	1.1	6.9	25.1	77.1	236.5	683.4	2,351.4
Male												
2002 .....	94.9	*	*	*	0.2	1.6	10.8	37.2	102.0	289.3	841.7	2,338.9
2001 .....	95.6	*	*	*	0.2	2.0	10.3	36.1	105.3	304.1	860.3	2,420.7
2000 .....	97.8	*	*	*	0.2	1.7	10.5	37.2	110.2	318.2	884.2	2,572.9
1999 .....	100.0	*	*	*	0.2	1.6	10.6	38.9	112.1	334.1	917.9	2,619.9
Female												
2002 .....	91.4	*	*	*	*	0.6	3.4	11.0	38.5	136.4	482.1	1,987.5
2001 .....	94.2	*	*	*	*	0.5	3.2	11.3	40.7	143.9	498.0	2,076.2
2000 .....	96.3	*	*	*	*	0.5	3.3	11.1	42.4	149.1	508.5	2,172.9
1999 .....	99.4	*	*	*	*	0.6	3.3	11.8	44.8	156.4	531.2	2,243.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)												
Non-Hispanic white												
Both sexes												
2002 .....	104.3	*	*	*	0.1	1.1	6.9	22.5	65.9	204.1	631.9	2,149.8
2001 .....	106.0	*	*	*	0.1	1.1	6.4	22.2	69.0	214.3	647.0	2,233.4
2000 .....	108.4	*	*	*	0.1	1.0	6.5	22.7	71.9	223.5	662.9	2,353.3
1999 .....	111.3	*	*	*	0.1	1.0	6.5	23.9	74.1	234.2	690.9	2,414.4
Male												
2002 .....	106.2	*	*	*	0.2	1.6	10.7	36.1	98.7	289.9	859.2	2,419.7
2001 .....	106.6	*	*	*	0.2	1.8	10.0	34.9	102.7	303.9	874.6	2,503.4
2000 .....	109.2	*	*	*	0.2	1.6	10.3	36.1	107.5	319.8	900.4	2,666.6
1999 .....	111.4	*	*	*	0.2	1.5	10.1	37.7	109.1	335.3	936.9	2,703.9
Female												
2002 .....	102.4	*	*	*	*	0.5	3.1	9.2	34.9	131.0	479.6	2,033.3
2001 .....	105.3	*	*	*	*	0.5	2.9	9.8	37.2	138.2	496.0	2,121.0
2000 .....	107.7	*	*	*	*	0.4	2.7	9.5	38.3	142.4	506.8	2,227.0
1999 .....	111.1	*	*	*	*	0.5	2.8	10.4	41.1	149.7	529.9	2,299.0
Non-Hispanic black												
Both sexes												
2002 .....	57.8	*	*	*	*	1.5	9.6	37.2	108.4	258.0	640.3	1,666.3
2001 .....	59.7	*	*	*	0.4	2.1	9.8	37.1	108.4	274.7	668.1	1,761.4
2000 .....	59.8	*	*	*	*	1.8	10.4	37.0	113.4	284.4	666.8	1,745.8
1999 .....	61.0	*	*	*	*	1.6	10.7	38.9	115.9	294.7	667.1	1,801.6
Male												
2002 .....	56.1	*	*	*	*	1.9	13.3	52.0	152.5	337.3	760.8	1,650.8
2001 .....	58.4	*	*	*	*	3.3	14.1	53.2	148.2	364.1	825.5	1,768.3
2000 .....	57.3	*	*	*	*	2.6	14.1	51.3	153.9	357.2	808.7	1,751.9
1999 .....	58.8	*	*	*	*	2.4	15.3	55.5	160.0	374.0	790.8	1,838.6
Female												
2002 .....	59.3	*	*	*	*	1.1	6.2	24.5	73.0	201.8	572.2	1,672.2
2001 .....	60.9	*	*	*	*	1.0	6.0	23.3	76.5	211.7	579.7	1,758.9
2000 .....	62.1	*	*	*	*	1.1	7.1	24.7	80.7	233.1	587.0	1,743.5
1999 .....	63.0	*	*	*	*	0.8	6.6	24.6	80.7	238.9	597.9	1,787.8
Other heart diseases (I26-I51)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	58.8	11.7	1.0	0.6	2.1	4.5	10.3	21.7	48.7	129.6	390.9	1,443.5
2001 .....	58.6	11.1	1.4	0.6	2.0	4.4	9.9	21.6	49.5	131.4	389.5	1,475.5
2000 .....	58.8	12.0	1.1	0.6	2.2	4.4	9.9	21.5	50.9	134.2	391.3	1,509.4
1999 .....	59.4	12.9	1.2	0.6	2.3	4.4	10.3	21.6	52.9	138.8	398.5	1,512.9
Male												
2002 .....	52.4	12.1	1.0	0.6	2.7	5.6	12.8	28.2	60.8	156.5	448.2	1,448.6
2001 .....	51.8	10.8	1.5	0.6	2.4	5.2	12.4	27.6	61.7	158.7	452.0	1,473.4
2000 .....	52.2	12.2	1.2	0.7	2.6	5.2	12.2	27.8	64.7	164.0	451.0	1,549.1
1999 .....	53.1	13.1	1.3	0.7	2.7	5.3	12.9	27.7	67.5	172.7	459.7	1,548.9
Female												
2002 .....	65.0	11.4	1.0	0.5	1.4	3.4	7.8	15.5	37.5	107.2	352.8	1,441.3

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
All origins <sup>3</sup>												
Female												
2001 .....	65.2	11.5	1.3	0.6	1.5	3.6	7.5	15.9	38.3	108.7	348.4	1,476.4
2000 .....	65.1	11.8	0.9	0.5	1.8	3.5	7.5	15.3	38.2	109.6	352.3	1,493.2
1999 .....	65.5	12.7	1.0	0.6	1.8	3.6	7.7	15.6	39.4	111.1	358.7	1,498.5
Hispanic												
Both sexes												
2002 .....	13.3	9.7	0.7	0.4	1.5	2.6	5.1	11.4	25.7	72.6	219.8	740.0
2001 .....	13.3	9.2	1.0	0.5	1.3	2.6	4.9	10.9	27.9	75.4	217.5	798.5
2000 .....	13.6	10.2	1.0	0.5	1.7	2.8	5.5	11.6	28.8	75.9	227.0	829.4
1999 .....	13.8	12.2	1.2	0.5	1.6	2.8	6.3	11.6	30.6	78.7	229.6	827.9
Male												
2002 .....	12.9	9.1	*	*	2.1	3.4	6.4	14.3	31.7	92.5	256.9	758.0
2001 .....	12.6	9.6	*	*	1.6	3.2	6.5	15.0	34.1	91.1	248.6	775.2
2000 .....	13.2	9.9	1.3	0.7	2.1	3.5	6.7	15.1	37.2	94.1	250.8	890.8
1999 .....	13.7	10.3	1.4	0.6	1.8	3.8	8.2	15.4	38.8	102.4	267.5	888.6
Female												
2002 .....	13.7	10.3	*	*	0.8	1.6	3.7	8.5	20.3	56.7	194.5	731.1
2001 .....	13.9	8.7	*	*	0.9	1.9	3.2	6.9	22.5	63.0	196.4	809.8
2000 .....	14.0	10.6	*	*	1.1	1.9	4.3	8.1	21.4	61.6	211.0	799.3
1999 .....	13.9	14.1	*	*	1.3	1.6	4.2	8.0	23.4	60.1	204.3	798.6
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	65.7	12.2	1.1	0.6	2.2	4.9	11.0	22.7	50.4	133.2	398.4	1,469.2
2001 .....	65.2	11.6	1.5	0.6	2.1	4.7	10.6	22.6	51.1	134.7	396.5	1,497.8
2000 .....	65.1	12.4	1.1	0.6	2.3	4.7	10.4	22.3	52.4	137.5	397.5	1,530.9
1999 .....	65.6	13.1	1.1	0.6	2.4	4.7	10.8	22.4	54.3	142.0	404.4	1,533.9
Male												
2002 .....	58.7	12.8	1.1	0.7	2.8	6.2	13.7	29.5	62.9	160.4	456.6	1,477.3
2001 .....	57.8	11.0	1.6	0.6	2.6	5.6	13.2	28.7	63.6	162.5	460.6	1,499.0
2000 .....	58.0	12.7	1.2	0.7	2.7	5.5	12.9	28.9	66.4	167.8	458.9	1,573.6
1999 .....	58.6	13.8	1.3	0.7	2.9	5.5	13.5	28.7	69.3	176.2	466.4	1,571.2
Female												
2002 .....	72.3	11.6	1.0	0.5	1.5	3.7	8.4	16.1	38.7	110.4	359.7	1,465.7
2001 .....	72.3	12.2	1.3	0.6	1.6	3.9	8.0	16.7	39.4	111.6	354.4	1,497.3
2000 .....	71.8	12.0	1.0	0.5	1.9	3.8	7.9	15.9	39.4	112.5	357.5	1,513.7
1999 .....	72.2	12.3	1.0	0.6	1.9	3.9	8.1	16.2	40.5	113.9	364.2	1,519.0
Non-Hispanic white												
Both sexes												
2002 .....	71.3	10.0	0.9	0.5	1.8	3.8	9.3	19.0	45.3	127.6	398.1	1,503.1
2001 .....	70.6	9.1	1.1	0.5	1.7	3.7	8.5	19.0	45.8	128.7	394.8	1,533.2
2000 .....	70.3	10.5	0.9	0.5	1.7	3.7	8.4	18.7	47.4	131.2	395.8	1,564.2
1999 .....	70.4	10.5	0.9	0.5	2.0	3.7	8.7	18.7	48.3	135.2	401.1	1,565.9
Male												
2002 .....	62.7	10.7	1.1	0.6	2.3	4.7	11.8	25.2	57.1	154.1	457.1	1,512.9
2001 .....	61.4	9.4	1.2	0.6	2.0	4.4	10.9	24.8	57.5	155.8	457.4	1,530.7
2000 .....	61.6	10.9	1.1	0.6	2.0	4.4	10.7	24.6	60.7	161.0	457.6	1,606.4
1999 .....	61.7	11.6	1.1	0.5	2.3	4.4	11.2	24.5	62.1	168.5	462.9	1,595.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
Non-Hispanic white												
Female												
2002 .....	79.6	9.3	0.8	0.5	1.2	2.8	6.8	13.0	34.1	105.0	358.5	1,498.8
2001 .....	79.4	8.8	1.0	0.5	1.3	3.1	6.1	13.4	34.8	105.7	353.3	1,534.2
2000 .....	78.7	10.1	0.8	0.4	1.4	2.9	6.2	12.9	34.8	106.0	355.3	1,547.2
1999 .....	78.8	9.4	0.8	0.5	1.7	3.0	6.2	13.1	35.3	107.3	360.6	1,554.1
Non-Hispanic black												
Both sexes												
2002 .....	54.3	20.9	1.6	0.9	4.2	11.8	23.4	52.6	104.1	211.2	474.7	1,285.5
2001 .....	54.7	21.3	3.0	1.0	4.4	11.0	24.5	52.2	105.7	216.1	486.4	1,292.9
2000 .....	54.6	19.6	2.0	1.1	4.9	10.8	23.7	52.8	104.5	223.6	484.3	1,319.4
1999 .....	56.2	23.2	2.0	1.0	4.7	11.0	25.3	53.5	115.3	232.9	499.9	1,314.0
Male												
2002 .....	52.6	21.8	*	0.9	5.1	14.8	28.2	67.5	130.4	259.4	543.4	1,258.0
2001 .....	53.5	18.8	3.3	0.9	5.5	13.4	29.3	63.9	134.9	261.9	593.5	1,362.9
2000 .....	53.3	19.1	2.0	1.2	5.7	12.7	28.7	68.3	132.9	269.8	565.5	1,416.1
1999 .....	56.2	22.9	2.3	1.2	6.0	13.4	30.9	68.7	149.5	291.0	584.7	1,466.6
Female												
2002 .....	55.9	19.9	1.9	0.8	3.2	9.0	19.2	39.9	83.0	177.0	435.9	1,296.0
2001 .....	55.8	23.8	2.7	1.0	3.2	8.9	20.3	42.2	82.3	183.7	426.2	1,267.0
2000 .....	55.8	20.1	1.9	1.0	4.2	9.0	19.2	39.6	81.6	191.1	438.5	1,283.4
1999 .....	56.3	23.6	1.8	0.9	3.4	8.9	20.4	40.5	87.9	192.0	452.4	1,257.0
Acute and subacute endocarditis (I33)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.6
2001 .....	0.4	*	*	*	*	0.1	0.3	0.5	0.6	1.3	2.5	2.5
2000 .....	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.4	2.8
1999 .....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
Male												
2002 .....	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.5
2001 .....	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.5	3.8
2000 .....	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.5	3.5	4.2
1999 .....	0.5	*	*	*	*	0.1	0.3	0.6	0.9	1.4	3.2	4.7
Female												
2002 .....	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	2.0	2.2
2001 .....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.1	1.8	2.0
2000 .....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	1.8	2.3
1999 .....	0.3	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.1
Hispanic												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
Hispanic												
Male												
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.7
2001 .....	0.5	*	*	*	*	0.1	0.3	0.5	0.6	1.4	2.5	2.6
2000 .....	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.5	2.9
1999 .....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
Male												
2002 .....	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.6
2001 .....	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.6	3.9
2000 .....	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.4	3.6	4.2
1999 .....	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.7
Female												
2002 .....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.3
2001 .....	0.4	*	*	*	*	*	0.3	0.3	0.5	1.2	1.8	2.0
2000 .....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.0	1.8	2.3
1999 .....	0.4	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.2
Non-Hispanic white												
Both sexes												
2002 .....	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.5	2.7
2001 .....	0.4	*	*	*	*	*	0.2	0.4	0.5	1.2	2.5	2.7
2000 .....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.1	2.4	2.8
1999 .....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	3.0
Male												
2002 .....	0.5	*	*	*	*	0.2	0.3	0.5	0.6	1.5	3.2	3.7
2001 .....	0.5	*	*	*	*	*	0.2	0.5	0.7	1.4	3.6	4.0
2000 .....	0.5	*	*	*	*	*	0.3	0.5	0.8	1.3	3.6	4.3
1999 .....	0.5	*	*	*	*	*	0.2	0.4	0.7	1.3	3.3	5.1
Female												
2002 .....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.0	2.3
2001 .....	0.4	*	*	*	*	*	0.2	0.2	0.4	1.1	1.7	2.1
2000 .....	0.4	*	*	*	*	*	0.1	0.2	0.4	1.0	1.7	2.2
1999 .....	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8	2.2
Non-Hispanic black												
Both sexes												
2002 .....	0.5	*	*	*	*	*	0.4	1.3	1.6	2.3	*	*
2001 .....	0.7	*	*	*	*	*	0.7	1.3	1.7	2.6	4.0	*
2000 .....	0.6	*	*	*	*	*	0.6	1.4	1.6	1.9	2.9	*
1999 .....	0.6	*	*	*	*	*	0.7	1.2	2.0	2.6	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
Non-Hispanic black												
Male												
2002 .....	0.6	*	*	*	*	*	*	1.6	1.9	*	*	*
2001 .....	0.8	*	*	*	*	*	0.8	1.9	2.2	3.9	*	*
2000 .....	0.7	*	*	*	*	*	0.8	1.8	*	*	*	*
1999 .....	0.7	*	*	*	*	*	1.0	1.9	2.5	*	*	*
Female												
2002 .....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2001 .....	0.5	*	*	*	*	*	0.7	*	*	*	*	*
2000 .....	0.6	*	*	*	*	*	*	1.0	1.6	*	*	*
1999 .....	0.5	*	*	*	*	*	*	*	1.6	2.9	*	*
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.3	0.6	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	2.1
2001 .....	0.3	0.8	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000 .....	0.3	0.5	0.2	0.0	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.8
1999 .....	0.3	0.7	0.2	0.1	0.1	0.1	0.3	0.3	0.5	0.8	1.3	2.0
Male												
2002 .....	0.3	*	*	*	0.1	0.2	0.2	0.4	0.5	0.9	1.2	2.4
2001 .....	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000 .....	0.3	*	*	*	0.1	0.1	0.3	0.2	0.4	0.8	1.5	2.1
1999 .....	0.3	*	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.4	2.5
Female												
2002 .....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.0
2001 .....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	2.0
2000 .....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.0	1.7
1999 .....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	1.9
Hispanic												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.3	0.7	0.2	*	0.1	0.1	0.2	0.3	0.4	0.8	1.3	2.1
2001 .....	0.3	0.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000 .....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.1	1.8
1999 .....	0.3	0.8	0.2	*	0.1	0.1	0.3	0.3	0.6	0.8	1.3	2.1
Male												
2002 .....	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.9	1.2	2.3
2001 .....	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000 .....	0.3	*	*	*	0.1	0.2	0.3	0.3	0.4	0.9	1.4	2.1
1999 .....	0.3	*	*	*	*	0.2	0.3	0.3	0.6	0.9	1.4	2.6
Female												
2002 .....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.0
2001 .....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	1.9
2000 .....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.6	1.0	1.7
1999 .....	0.3	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	1.9
Non-Hispanic white												
Both sexes												
2002 .....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.2	2.1
2001 .....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000 .....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.1	1.8
1999 .....	0.3	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.3	2.0
Male												
2002 .....	0.3	*	*	*	*	0.2	0.2	0.3	0.4	0.9	1.2	2.3
2001 .....	0.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.2	2.0
2000 .....	0.3	*	*	*	*	*	0.2	0.2	0.4	0.8	1.4	2.1
1999 .....	0.3	*	*	*	*	*	0.2	0.3	0.7	0.8	1.4	2.6
Female												
2002 .....	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.1
2001 .....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	1.3	1.8
2000 .....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.0	1.7
1999 .....	0.3	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	1.8
Non-Hispanic black												
Both sexes												
2002 .....	0.4	*	*	*	*	*	*	0.6	0.9	*	2.1	*
2001 .....	0.4	*	*	*	*	*	0.4	0.7	*	*	*	*
2000 .....	0.4	*	*	*	*	*	0.4	0.5	*	*	*	*
1999 .....	0.5	*	*	*	*	*	0.5	0.5	*	1.2	*	*
Male												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.5	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.5	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.4	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Non-Hispanic black												
Female												
1999 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
Heart failure (I50)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	19.6	0.6	*	*	0.1	0.3	0.6	2.5	8.8	33.8	132.8	645.0
2001 .....	20.0	0.6	*	*	0.1	0.2	0.7	2.4	9.4	34.3	136.3	674.6
2000 .....	19.8	0.8	*	0.1	0.1	0.2	0.6	2.2	9.1	34.8	136.1	684.9
1999 .....	19.7	*	*	*	0.1	0.2	0.6	2.1	8.9	35.4	137.2	681.9
Male												
2002 .....	15.3	*	*	*	*	0.3	0.8	3.1	10.5	39.2	149.0	621.4
2001 .....	15.5	*	*	*	*	0.3	0.8	3.1	11.1	40.3	153.8	646.4
2000 .....	15.3	1.1	*	*	0.1	0.2	0.7	2.7	10.9	41.1	153.7	672.2
1999 .....	15.2	*	*	*	0.1	0.2	0.7	2.7	10.8	42.5	152.0	667.0
Female												
2002 .....	23.7	*	*	*	*	0.2	0.5	1.8	7.2	29.2	122.0	655.3
2001 .....	24.3	*	*	*	*	0.2	0.6	1.8	7.8	29.3	124.7	686.4
2000 .....	24.1	*	*	*	*	0.1	0.5	1.6	7.5	29.6	124.6	690.0
1999 .....	24.0	*	*	*	*	0.2	0.5	1.6	7.1	29.6	127.6	687.9
Hispanic												
Both sexes												
2002 .....	3.6	*	*	*	*	*	*	1.2	4.6	19.9	75.2	316.0
2001 .....	3.7	*	*	*	*	*	0.4	1.3	5.6	18.8	72.9	349.5
2000 .....	3.6	*	*	*	*	*	*	0.7	5.8	18.9	72.1	369.6
1999 .....	3.7	*	*	*	*	*	*	1.0	5.3	19.5	73.5	377.4
Male												
2002 .....	3.0	*	*	*	*	*	*	1.1	5.8	23.1	85.8	296.2
2001 .....	3.0	*	*	*	*	*	*	1.8	6.0	21.8	75.6	333.3
2000 .....	2.9	*	*	*	*	*	*	*	7.7	23.0	74.2	380.9
1999 .....	3.0	*	*	*	*	*	*	*	6.1	24.6	82.9	358.0
Female												
2002 .....	4.4	*	*	*	*	*	*	1.3	3.6	17.3	68.0	325.8
2001 .....	4.5	*	*	*	*	*	*	*	5.2	16.4	71.0	357.4
2000 .....	4.3	*	*	*	*	*	*	*	4.2	15.6	70.7	364.0
1999 .....	4.4	*	*	*	*	*	*	*	4.6	15.5	67.3	386.8
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	22.0	*	*	*	0.1	0.3	0.7	2.6	9.1	34.6	135.3	656.8
2001 .....	22.4	*	*	*	0.1	0.3	0.7	2.5	9.6	35.2	138.8	685.6
2000 .....	22.1	0.8	*	*	0.1	0.2	0.6	2.3	9.3	35.7	138.5	694.8
1999 .....	21.8	*	*	*	0.1	0.2	0.6	2.2	9.1	36.3	139.4	691.0
Male												
2002 .....	17.3	*	*	*	*	0.4	0.9	3.3	10.9	40.2	151.8	634.7
2001 .....	17.4	*	*	*	*	0.3	0.9	3.2	11.4	41.4	157.2	658.0
2000 .....	17.2	*	*	*	0.1	0.2	0.8	2.9	11.0	42.1	156.9	683.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Non-Hispanic <sup>4</sup>												
Male												
1999 .....	16.9	*	*	*	0.1	0.2	0.8	2.8	11.0	43.5	154.5	677.1
Female												
2002 .....	26.5	*	*	*	*	0.2	0.5	1.9	7.5	29.9	124.3	666.4
2001 .....	27.1	*	*	*	*	0.2	0.6	1.9	8.0	30.1	126.8	697.0
2000 .....	26.7	*	*	*	*	0.2	0.5	1.7	7.7	30.5	126.5	699.5
1999 .....	26.5	*	*	*	*	0.2	0.5	1.6	7.2	30.4	129.7	696.5
Non-Hispanic white												
Both sexes												
2002 .....	24.7	*	*	*	*	0.2	0.5	2.0	8.2	33.3	135.9	677.3
2001 .....	25.1	*	*	*	*	0.2	0.5	2.1	8.7	33.9	139.7	706.8
2000 .....	24.7	*	*	*	*	0.1	0.5	1.9	8.4	34.3	139.2	714.3
1999 .....	24.3	*	*	*	*	0.1	0.5	1.7	8.1	34.9	139.1	709.1
Male												
2002 .....	19.2	*	*	*	*	0.2	0.7	2.6	9.8	38.6	153.2	656.2
2001 .....	19.3	*	*	*	*	0.2	0.7	2.7	10.5	40.0	158.1	679.1
2000 .....	19.0	*	*	*	*	0.2	0.6	2.4	10.1	41.0	157.9	701.6
1999 .....	18.6	*	*	*	*	*	0.6	2.2	10.1	42.2	154.7	690.8
Female												
2002 .....	30.0	*	*	*	*	0.2	0.4	1.4	6.7	28.8	124.2	686.4
2001 .....	30.7	*	*	*	*	0.2	0.4	1.4	6.9	28.8	127.5	718.4
2000 .....	30.2	*	*	*	*	*	0.3	1.3	6.8	28.7	126.9	719.4
1999 .....	29.9	*	*	*	*	*	0.3	1.2	6.3	28.8	128.8	716.4
Non-Hispanic black												
Both sexes												
2002 .....	14.2	*	*	*	*	0.9	1.9	7.4	19.0	54.1	158.3	534.5
2001 .....	14.3	*	*	*	*	0.6	2.0	6.6	20.0	55.5	159.1	551.2
2000 .....	14.1	*	*	*	*	0.5	1.7	5.9	19.0	56.2	156.3	564.8
1999 .....	14.3	*	*	*	*	0.6	1.7	6.5	19.2	57.3	164.8	557.4
Male												
2002 .....	12.0	*	*	*	*	1.1	2.3	9.2	23.3	65.9	176.4	489.0
2001 .....	12.0	*	*	*	*	0.9	2.3	7.6	22.6	66.3	185.4	547.5
2000 .....	11.6	*	*	*	*	*	1.8	7.3	21.8	62.9	176.5	589.4
1999 .....	12.2	*	*	*	*	*	1.9	8.6	23.0	66.4	180.4	609.7
Female												
2002 .....	16.3	*	*	*	*	*	1.5	5.7	15.5	45.8	148.1	551.8
2001 .....	16.4	*	*	*	*	*	1.7	5.6	18.0	47.9	144.4	552.6
2000 .....	16.3	*	*	*	*	*	1.6	4.7	16.8	51.5	145.0	555.6
1999 .....	16.2	*	*	*	*	*	1.5	4.8	16.1	50.9	156.0	537.9
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	38.5	10.5	0.8	0.5	1.9	4.0	9.3	18.6	38.8	93.7	254.4	793.7
2001 .....	37.9	9.7	1.1	0.5	1.8	4.0	8.8	18.5	39.1	95.1	249.5	796.5
2000 .....	38.3	10.6	0.9	0.5	2.0	4.0	8.8	18.6	40.8	97.4	251.6	820.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
All origins <sup>3</sup>												
Both sexes												
1999 .....	39.0	11.9	0.9	0.5	2.1	4.1	9.2	18.7	42.8	101.3	257.6	826.1
Male												
2002 .....	36.3	10.9	0.8	0.5	2.5	5.0	11.5	24.1	49.0	114.8	294.9	821.3
2001 .....	35.6	9.1	1.2	0.5	2.2	4.7	11.1	23.6	49.4	116.1	293.4	821.3
2000 .....	36.1	10.5	1.0	0.6	2.3	4.8	10.9	24.3	52.6	120.7	292.4	870.7
1999 .....	37.1	12.1	1.0	0.6	2.5	4.8	11.6	24.2	55.3	127.9	303.1	874.7
Female												
2002 .....	40.6	10.0	0.8	0.4	1.3	3.1	7.1	13.2	29.3	76.2	227.6	781.7
2001 .....	40.2	10.4	1.0	0.5	1.3	3.2	6.5	13.5	29.6	77.7	220.7	786.1
2000 .....	40.4	10.7	0.7	0.4	1.6	3.2	6.8	13.1	29.9	78.3	225.0	799.3
1999 .....	40.9	11.6	0.8	0.5	1.7	3.3	6.9	13.5	31.3	79.6	228.0	806.5
Hispanic												
Both sexes												
2002 .....	9.3	8.6	0.6	0.3	1.4	2.4	4.7	9.6	19.7	50.9	141.6	419.6
2001 .....	9.3	8.0	0.8	0.4	1.2	2.2	4.3	9.2	21.6	55.3	141.6	444.6
2000 .....	9.6	8.7	0.7	0.5	1.6	2.5	4.9	10.2	22.0	54.9	152.1	457.2
1999 .....	9.8	11.5	0.9	0.4	1.3	2.5	5.5	10.1	24.6	57.2	151.9	447.6
Male												
2002 .....	9.6	8.0	*	*	2.0	3.1	6.0	12.2	24.2	67.1	168.2	455.1
2001 .....	9.3	7.9	*	*	1.5	2.8	5.7	12.6	27.3	67.7	170.4	438.1
2000 .....	9.9	8.1	*	0.7	2.1	3.3	6.0	13.7	28.6	68.8	172.6	505.9
1999 .....	10.3	9.8	*	*	1.5	3.4	7.2	13.7	31.9	75.6	179.4	526.2
Female												
2002 .....	9.0	9.3	*	*	0.8	1.5	3.3	7.0	15.8	38.0	123.4	402.0
2001 .....	9.2	8.0	*	*	0.9	1.6	2.8	5.8	16.5	45.5	122.1	447.7
2000 .....	9.3	9.3	*	*	1.0	1.7	3.8	6.8	16.1	44.0	138.3	433.3
1999 .....	9.2	13.3	*	*	1.1	1.5	3.7	6.6	18.1	42.9	133.6	409.7
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	42.9	10.9	0.8	0.5	2.0	4.4	9.9	19.4	40.2	96.5	259.4	807.6
2001 .....	42.1	10.1	1.2	0.5	1.9	4.3	9.4	19.3	40.4	97.5	253.9	807.8
2000 .....	42.3	11.0	0.9	0.5	2.0	4.3	9.3	19.3	42.0	99.8	255.4	831.4
1999 .....	43.0	11.9	0.9	0.5	2.2	4.3	9.6	19.5	44.0	103.6	261.3	837.9
Male												
2002 .....	40.6	11.6	0.8	0.6	2.6	5.5	12.3	25.3	50.8	117.7	300.6	836.7
2001 .....	39.6	9.3	1.4	0.5	2.4	5.1	11.8	24.5	50.9	118.8	298.5	835.3
2000 .....	40.0	10.9	1.0	0.6	2.4	5.1	11.5	25.1	54.1	123.4	297.1	883.9
1999 .....	40.9	12.7	1.0	0.6	2.6	5.1	12.1	25.0	56.7	130.5	307.4	886.8
Female												
2002 .....	45.1	10.1	0.8	0.4	1.4	3.3	7.6	13.8	30.3	78.7	232.2	795.0
2001 .....	44.5	11.0	1.0	0.5	1.4	3.5	6.9	14.2	30.6	79.7	224.6	796.3
2000 .....	44.5	11.0	0.8	0.4	1.7	3.4	7.1	13.6	30.9	80.4	228.3	810.2
1999 .....	44.9	11.1	0.9	0.5	1.8	3.6	7.2	14.1	32.2	81.5	231.4	818.4

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Non-Hispanic white												
Both sexes												
2002 .....	45.8	9.2	0.7	0.5	1.6	3.4	8.4	16.5	36.2	92.3	258.6	820.9
2001 .....	44.7	8.1	0.9	0.4	1.5	3.4	7.6	16.3	36.2	92.9	251.4	821.8
2000 .....	44.9	9.6	0.8	0.4	1.6	3.4	7.6	16.3	38.0	95.0	253.1	845.3
1999 .....	45.4	9.8	0.8	0.5	1.9	3.4	7.8	16.4	39.0	98.3	258.4	851.7
Male												
2002 .....	42.7	9.8	0.8	0.5	2.1	4.2	10.7	21.9	46.2	113.1	299.5	850.7
2001 .....	41.3	8.1	1.0	0.5	1.9	4.0	9.9	21.3	45.8	113.8	294.4	845.6
2000 .....	41.7	9.6	0.8	0.5	1.8	4.1	9.6	21.5	49.3	117.9	294.7	898.3
1999 .....	42.4	10.8	0.9	0.5	2.1	4.1	10.1	21.5	50.6	124.1	303.5	897.0
Female												
2002 .....	48.9	8.5	0.6	0.4	1.1	2.5	6.2	11.2	26.6	74.6	231.1	808.1
2001 .....	47.9	8.1	0.7	0.4	1.1	2.7	5.3	11.5	27.2	75.2	222.8	811.9
2000 .....	47.9	9.6	0.7	0.3	1.3	2.7	5.6	11.1	27.4	75.7	225.7	824.0
1999 .....	48.2	8.7	0.6	0.5	1.6	2.8	5.5	11.4	28.1	76.7	228.9	833.7
Non-Hispanic black												
Both sexes												
2002 .....	39.1	17.8	1.2	0.7	3.9	10.5	20.9	43.4	82.7	153.6	312.3	747.1
2001 .....	39.3	18.2	2.4	0.9	3.8	10.0	21.4	43.7	83.3	157.1	322.1	737.9
2000 .....	39.5	16.1	1.5	0.9	4.5	9.8	21.0	45.1	83.1	164.4	323.9	748.9
1999 .....	40.9	20.9	1.5	0.9	4.4	10.1	22.3	45.2	93.3	171.8	331.4	752.7
Male												
2002 .....	39.6	19.7	*	0.8	4.7	13.3	25.0	55.8	104.0	189.7	363.1	763.6
2001 .....	40.3	16.0	2.7	0.9	5.0	11.9	25.9	53.5	109.1	190.6	401.2	813.1
2000 .....	40.5	15.0	1.7	1.0	5.1	11.7	25.4	58.6	108.9	202.9	384.1	821.9
1999 .....	42.8	20.8	*	1.1	5.6	12.1	27.4	57.6	123.1	220.6	401.5	855.7
Female												
2002 .....	38.7	15.8	*	*	3.0	8.0	17.2	32.8	65.5	127.9	283.5	740.8
2001 .....	38.5	20.5	2.1	0.9	2.7	8.3	17.3	35.3	62.5	133.5	277.6	710.1
2000 .....	38.6	17.3	*	0.8	3.8	8.1	17.0	33.4	62.4	137.3	290.0	721.7
1999 .....	39.2	21.1	*	*	3.2	8.2	17.9	34.7	69.5	137.3	292.2	714.2
Essential (primary) hypertension and hypertensive renal disease (I10,I12)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	7.0	*	*	*	0.1	0.2	0.8	2.3	5.7	16.0	48.2	180.4
2001 .....	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.5	47.7	171.9
2000 .....	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9
1999 .....	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1
Male												
2002 .....	5.4	*	*	*	*	0.2	1.0	2.8	6.6	17.1	47.3	153.4
2001 .....	5.2	*	*	*	*	0.3	0.9	3.0	6.5	16.6	45.8	150.8
2000 .....	5.0	*	*	*	*	0.2	1.0	3.0	7.0	15.8	45.2	142.4
1999 .....	4.7	*	*	*	*	0.2	0.9	2.7	6.4	16.5	42.9	132.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)												
All origins <sup>3</sup>												
Female												
2002 .....	8.6	*	*	*	*	0.2	0.6	1.8	4.9	15.1	48.8	192.1
2001 .....	8.3	*	*	*	*	0.2	0.6	1.8	5.2	14.5	48.9	180.8
2000 .....	7.8	*	*	*	*	0.2	0.6	1.7	4.9	14.6	45.7	171.3
1999 .....	7.4	*	*	*	*	0.2	0.5	1.7	4.7	14.1	43.9	160.0
Hispanic												
Both sexes												
2002 .....	2.4	*	*	*	*	*	0.5	1.1	6.2	15.5	47.2	148.6
2001 .....	2.4	*	*	*	*	*	*	1.8	4.5	16.5	48.9	160.1
2000 .....	2.3	*	*	*	*	*	0.5	1.8	6.4	15.0	45.8	148.0
1999 .....	2.2	*	*	*	*	*	*	2.2	5.0	15.2	48.3	129.8
Male												
2002 .....	1.9	*	*	*	*	*	*	1.2	7.1	15.3	46.4	132.2
2001 .....	1.9	*	*	*	*	*	*	2.3	3.9	16.8	51.0	129.5
2000 .....	2.1	*	*	*	*	*	0.8	2.8	8.1	15.2	52.6	141.1
1999 .....	1.9	*	*	*	*	*	*	2.4	5.2	18.5	51.6	131.6
Female												
2002 .....	3.0	*	*	*	*	*	*	*	5.5	15.7	47.8	156.7
2001 .....	2.9	*	*	*	*	*	*	1.2	5.0	16.2	47.4	175.0
2000 .....	2.5	*	*	*	*	*	*	*	4.9	14.9	41.3	151.3
1999 .....	2.5	*	*	*	*	*	*	1.9	4.8	12.7	46.1	128.9
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	7.7	*	*	*	*	0.2	0.8	2.4	5.7	15.9	48.1	181.1
2001 .....	7.4	*	*	*	0.1	0.3	0.8	2.4	5.8	15.3	47.5	172.0
2000 .....	7.0	*	*	*	*	0.2	0.8	2.4	5.8	15.1	45.3	162.9
1999 .....	6.6	*	*	*	*	0.2	0.7	2.2	5.6	15.1	43.2	152.3
Male												
2002 .....	5.9	*	*	*	*	0.3	1.0	3.0	6.6	17.1	47.2	153.9
2001 .....	5.7	*	*	*	*	0.3	1.0	3.0	6.6	16.5	45.3	151.1
2000 .....	5.4	*	*	*	*	0.2	1.0	3.0	6.9	15.7	44.7	142.0
1999 .....	5.1	*	*	*	*	0.2	0.9	2.7	6.5	16.4	42.4	131.9
Female												
2002 .....	9.4	*	*	*	*	0.2	0.6	1.9	4.8	15.0	48.7	192.8
2001 .....	9.0	*	*	*	*	0.3	0.6	1.9	5.1	14.4	48.9	180.7
2000 .....	8.5	*	*	*	*	0.2	0.6	1.8	4.9	14.6	45.8	171.3
1999 .....	8.0	*	*	*	*	0.2	0.6	1.7	4.7	14.1	43.7	160.5
Non-Hispanic white												
Both sexes												
2002 .....	7.4	*	*	*	*	0.1	0.4	1.5	3.9	12.5	42.2	170.6
2001 .....	7.0	*	*	*	*	0.1	0.4	1.4	3.9	11.7	41.8	162.8
2000 .....	6.5	*	*	*	*	0.1	0.4	1.3	3.8	11.6	39.9	152.7
1999 .....	6.1	*	*	*	*	0.1	0.3	1.2	3.6	11.8	37.3	141.7
Male												
2002 .....	5.5	*	*	*	*	*	0.5	2.0	4.7	13.5	41.0	145.4
2001 .....	5.1	*	*	*	*	0.2	0.5	1.7	4.5	12.8	39.6	142.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)												
Non-Hispanic white												
Male												
2000 .....	4.9	*	*	*	*	*	0.6	1.8	4.7	12.4	39.5	131.8
1999 .....	4.6	*	*	*	*	0.2	0.5	1.6	4.5	13.0	36.5	119.8
Female												
2002 .....	9.2	*	*	*	*	*	0.3	1.1	3.1	11.5	42.9	181.4
2001 .....	8.8	*	*	*	*	*	0.3	1.0	3.3	10.8	43.2	171.5
2000 .....	8.1	*	*	*	*	*	0.3	0.9	3.1	10.9	40.1	161.1
1999 .....	7.6	*	*	*	*	*	0.2	0.9	2.8	10.8	37.8	150.5
Non-Hispanic black												
Both sexes												
2002 .....	11.5	*	*	*	*	0.9	3.4	9.2	21.4	48.3	120.0	312.4
2001 .....	11.3	*	*	*	*	1.1	3.2	10.1	22.4	48.4	113.9	291.5
2000 .....	11.0	*	*	*	*	0.7	3.2	9.9	23.3	49.5	108.9	285.9
1999 .....	10.7	*	*	*	*	0.7	3.2	9.4	21.8	47.5	112.9	272.2
Male												
2002 .....	9.9	*	*	*	*	1.0	4.3	11.3	24.0	52.6	129.5	276.8
2001 .....	10.0	*	*	*	*	1.2	3.9	13.1	25.8	55.5	119.7	277.2
2000 .....	9.5	*	*	*	*	*	3.8	12.1	28.4	51.5	110.5	268.9
1999 .....	9.4	*	*	*	*	*	3.8	12.0	24.8	52.3	119.1	268.6
Female												
2002 .....	13.0	*	*	*	*	0.7	2.6	7.4	19.4	45.3	114.6	326.1
2001 .....	12.4	*	*	*	*	1.0	2.5	7.5	19.7	43.4	110.7	296.9
2000 .....	12.4	*	*	*	*	*	2.6	8.0	19.3	48.1	108.0	292.2
1999 .....	11.8	*	*	*	*	0.7	2.7	7.2	19.4	44.1	109.4	273.6
Cerebrovascular diseases (I60-I69)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	56.4	2.9	0.3	0.2	0.4	1.4	5.4	15.1	37.2	120.3	431.0	1,445.9
2001 .....	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.1	38.0	123.4	443.9	1,500.2
2000 .....	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2
1999 .....	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8
Male												
2002 .....	44.2	3.2	0.4	0.2	0.5	1.4	5.3	16.7	42.7	135.0	445.9	1,317.9
2001 .....	45.2	3.1	0.3	0.2	0.5	1.6	5.7	16.7	43.4	140.4	467.3	1,380.2
2000 .....	46.9	3.8	*	0.2	0.5	1.5	5.8	17.5	47.2	145.0	490.8	1,484.3
1999 .....	47.1	3.3	0.3	0.2	0.5	1.5	5.8	16.7	46.5	147.3	500.4	1,512.8
Female												
2002 .....	68.2	2.5	0.3	0.2	0.3	1.4	5.5	13.6	32.1	108.1	421.2	1,501.5
2001 .....	69.2	2.3	0.4	0.2	0.5	1.5	5.4	13.6	32.9	109.3	428.6	1,550.4
2000 .....	71.8	2.7	0.4	0.2	0.5	1.5	5.7	14.5	35.3	115.1	442.1	1,632.0
1999 .....	72.3	2.1	0.3	0.2	0.5	1.4	5.6	13.8	35.1	117.2	449.8	1,656.0
Hispanic												
Both sexes												
2002 .....	16.6	2.4	*	0.3	0.4	1.3	4.4	15.3	35.9	102.9	319.9	898.0
2001 .....	17.4	*	*	*	0.6	1.4	5.0	15.9	35.4	106.2	341.2	1,042.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Cerebrovascular diseases (I60-I69)</b>												
<b>Hispanic</b>												
<b>Both sexes</b>												
2000 .....	17.5	*	*	*	0.5	1.7	5.2	15.2	39.8	113.2	344.1	1,082.9
1999 .....	17.4	*	*	*	0.5	1.6	4.9	15.7	37.7	116.9	348.3	1,077.6
<b>Male</b>												
2002 .....	15.0	*	*	*	*	1.5	5.1	18.6	45.0	124.6	338.5	856.7
2001 .....	15.7	*	*	*	0.7	1.6	5.3	18.7	43.5	127.2	386.3	1,005.6
2000 .....	15.8	*	*	*	*	1.9	5.5	18.1	48.8	136.1	392.9	1,029.9
1999 .....	16.1	*	*	*	0.6	1.8	5.7	19.5	46.1	140.8	395.2	1,136.6
<b>Female</b>												
2002 .....	18.4	*	*	*	*	1.0	3.7	12.0	27.6	85.6	307.2	918.5
2001 .....	19.1	*	*	*	*	1.2	4.7	13.1	28.2	89.6	310.7	1,061.2
2000 .....	19.4	*	*	*	*	1.4	4.9	12.4	31.9	95.2	311.3	1,108.9
1999 .....	18.8	*	*	*	*	1.4	4.0	11.9	30.3	98.1	317.1	1,049.2
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	62.4	3.0	0.3	0.2	0.4	1.5	5.5	15.0	37.2	121.3	435.5	1,465.0
2001 .....	63.2	2.8	0.3	0.2	0.5	1.5	5.6	14.9	38.0	124.3	447.7	1,514.0
2000 .....	65.4	3.6	0.3	0.2	0.5	1.5	5.8	15.9	40.9	129.2	465.0	1,604.5
1999 .....	65.7	2.9	0.3	0.2	0.5	1.4	5.8	15.1	40.6	131.2	473.8	1,630.4
<b>Male</b>												
2002 .....	48.9	3.5	0.4	0.2	0.5	1.4	5.3	16.4	42.3	135.4	450.1	1,336.0
2001 .....	49.7	3.3	0.3	0.2	0.5	1.5	5.7	16.4	43.2	140.9	469.8	1,392.9
2000 .....	51.5	4.2	*	0.2	0.5	1.5	5.8	17.3	46.8	145.1	493.7	1,499.9
1999 .....	51.5	3.7	0.3	0.2	0.5	1.4	5.8	16.4	46.3	147.3	503.6	1,524.8
<b>Female</b>												
2002 .....	75.3	2.5	*	0.2	0.3	1.5	5.8	13.7	32.4	109.5	425.8	1,520.7
2001 .....	76.1	2.4	0.4	0.2	0.5	1.5	5.5	13.5	33.2	110.4	433.2	1,564.4
2000 .....	78.7	2.8	0.4	0.2	0.5	1.5	5.8	14.6	35.5	116.1	446.4	1,646.8
1999 .....	79.2	2.0	*	0.2	0.5	1.4	5.8	13.9	35.4	118.1	454.4	1,672.7
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	67.0	2.2	0.3	0.2	0.4	1.2	4.1	11.1	30.8	110.8	427.5	1,479.2
2001 .....	67.6	1.9	0.3	0.2	0.4	1.3	4.2	10.8	31.2	113.0	439.5	1,525.0
2000 .....	70.0	2.6	0.3	0.2	0.5	1.2	4.3	12.0	34.2	119.3	455.7	1,616.8
1999 .....	70.4	2.4	0.3	0.2	0.4	1.1	4.3	11.2	33.4	120.7	465.1	1,648.0
<b>Male</b>												
2002 .....	51.3	2.7	*	0.2	0.5	1.1	4.0	12.1	34.5	123.2	441.1	1,345.9
2001 .....	52.0	2.2	*	0.2	0.4	1.3	4.3	11.9	35.1	128.0	460.5	1,399.0
2000 .....	53.9	3.3	*	0.2	0.5	1.2	4.3	13.0	38.7	133.1	482.3	1,505.9
1999 .....	53.9	3.1	*	0.2	0.4	1.1	4.4	12.0	38.0	135.0	492.8	1,535.1
<b>Female</b>												
2002 .....	82.1	*	*	0.2	0.3	1.2	4.3	10.1	27.2	100.2	418.4	1,536.7
2001 .....	82.6	*	*	*	0.3	1.3	4.1	9.8	27.4	100.3	425.6	1,577.4
2000 .....	85.5	1.9	*	0.2	0.4	1.2	4.3	10.9	29.9	107.6	438.3	1,661.6
1999 .....	86.1	*	*	*	0.4	1.1	4.3	10.5	29.1	108.8	447.0	1,693.0

See footnotes at end of table.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Non-Hispanic black												
Both sexes												
2002 .....	51.7	5.6	*	0.4	0.6	3.2	13.3	41.6	89.7	218.7	558.8	1,398.0
2001 .....	52.9	6.9	*	*	0.9	3.0	13.6	43.5	92.8	226.0	567.2	1,465.5
2000 .....	54.3	8.0	*	*	0.7	3.0	14.9	44.3	95.2	226.3	593.8	1,545.8
1999 .....	53.9	4.7	*	*	0.7	2.9	14.7	42.2	97.7	235.7	596.4	1,501.5
Male												
2002 .....	45.1	*	*	*	0.7	2.9	12.7	47.5	112.7	267.3	595.2	1,266.8
2001 .....	46.2	8.3	*	*	0.7	3.1	13.5	49.9	113.9	273.5	620.5	1,359.4
2000 .....	47.7	9.2	*	*	*	2.8	15.1	50.4	116.9	272.1	664.9	1,479.5
1999 .....	47.5	*	*	*	*	2.9	15.0	49.6	119.5	279.1	656.7	1,455.9
Female												
2002 .....	57.7	*	*	*	*	3.5	13.8	36.6	71.3	184.2	538.2	1,448.1
2001 .....	59.0	*	*	*	1.1	2.9	13.6	38.1	75.9	192.5	537.2	1,504.8
2000 .....	60.3	*	*	*	0.8	3.2	14.8	39.2	77.8	194.1	553.8	1,570.4
1999 .....	59.8	*	*	*	0.8	2.8	14.4	35.9	80.2	205.1	562.7	1,518.6
Atherosclerosis (I70)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	4.8	*	*	*	*	*	0.1	0.5	1.8	7.1	31.9	168.1
2001 .....	4.9	*	*	*	*	*	0.1	0.5	1.9	7.3	33.1	179.1
2000 .....	5.1	*	*	*	*	*	0.1	0.5	1.9	7.7	34.6	188.8
1999 .....	5.4	*	*	*	*	*	0.1	0.5	2.0	8.0	35.9	202.0
Male												
2002 .....	3.7	*	*	*	*	*	0.1	0.6	2.4	8.8	35.5	156.6
2001 .....	3.7	*	*	*	*	*	0.1	0.6	2.4	9.0	35.9	167.2
2000 .....	3.9	*	*	*	*	*	0.2	0.6	2.5	9.6	39.7	176.5
1999 .....	4.0	*	*	*	*	*	0.1	0.7	2.7	9.6	40.0	189.0
Female												
2002 .....	5.9	*	*	*	*	*	0.1	0.3	1.3	5.7	29.5	173.1
2001 .....	6.2	*	*	*	*	*	*	0.3	1.4	5.9	31.2	184.1
2000 .....	6.3	*	*	*	*	*	*	0.3	1.2	6.2	31.3	193.8
1999 .....	6.7	*	*	*	*	*	*	0.3	1.4	6.7	33.2	207.2
Hispanic												
Both sexes												
2002 .....	1.1	*	*	*	*	*	*	*	1.0	4.2	22.7	108.6
2001 .....	1.1	*	*	*	*	*	*	*	*	4.0	23.1	130.7
2000 .....	1.2	*	*	*	*	*	*	*	*	5.9	24.5	132.7
1999 .....	1.2	*	*	*	*	*	*	*	1.5	5.2	27.1	129.1
Male												
2002 .....	0.9	*	*	*	*	*	*	*	*	5.2	26.7	102.1
2001 .....	0.9	*	*	*	*	*	*	*	*	4.8	27.3	127.6
2000 .....	1.0	*	*	*	*	*	*	*	*	8.2	32.0	104.8
1999 .....	1.1	*	*	*	*	*	*	*	*	6.8	31.3	151.0
Female												
2002 .....	1.3	*	*	*	*	*	*	*	*	3.5	20.0	111.9
2001 .....	1.4	*	*	*	*	*	*	*	*	3.5	20.2	132.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Atherosclerosis (I70)</b>												
Hispanic												
Female												
2000 .....	1.4	*	*	*	*	*	*	*	*	4.0	19.5	146.4
1999 .....	1.4	*	*	*	*	*	*	*	*	4.0	24.3	118.5
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	5.4	*	*	*	*	*	0.1	0.5	1.9	7.3	32.2	170.2
2001 .....	5.5	*	*	*	*	*	0.1	0.5	1.9	7.5	33.4	180.5
2000 .....	5.7	*	*	*	*	*	0.1	0.5	1.9	7.8	35.0	190.3
1999 .....	5.9	*	*	*	*	*	0.1	0.5	2.1	8.1	36.1	203.9
Male												
2002 .....	4.1	*	*	*	*	*	0.1	0.6	2.5	9.0	35.8	158.7
2001 .....	4.1	*	*	*	*	*	0.1	0.6	2.4	9.2	36.1	168.2
2000 .....	4.3	*	*	*	*	*	0.2	0.7	2.6	9.7	39.9	178.7
1999 .....	4.4	*	*	*	*	*	0.2	0.7	2.7	9.7	40.3	189.9
Female												
2002 .....	6.6	*	*	*	*	*	*	0.3	1.4	5.9	29.8	175.1
2001 .....	6.8	*	*	*	*	*	*	0.3	1.5	6.0	31.6	185.5
2000 .....	7.0	*	*	*	*	*	*	0.3	1.3	6.3	31.8	195.1
1999 .....	7.4	*	*	*	*	*	*	0.3	1.5	6.9	33.4	209.6
Non-Hispanic white												
Both sexes												
2002 .....	6.1	*	*	*	*	*	0.1	0.4	1.8	7.3	32.5	177.0
2001 .....	6.3	*	*	*	*	*	0.1	0.5	1.8	7.5	34.1	187.4
2000 .....	6.5	*	*	*	*	*	0.1	0.4	1.8	7.8	35.5	196.8
1999 .....	6.7	*	*	*	*	*	0.1	0.5	2.0	8.2	36.7	209.9
Male												
2002 .....	4.6	*	*	*	*	*	0.1	0.5	2.3	9.0	35.9	165.7
2001 .....	4.7	*	*	*	*	*	0.1	0.6	2.3	9.2	36.6	175.2
2000 .....	4.9	*	*	*	*	*	0.2	0.6	2.6	9.8	40.6	183.6
1999 .....	4.9	*	*	*	*	*	0.1	0.7	2.6	9.6	40.4	194.6
Female												
2002 .....	7.6	*	*	*	*	*	*	0.3	1.3	5.9	30.2	181.9
2001 .....	7.9	*	*	*	*	*	*	0.3	1.3	6.1	32.5	192.6
2000 .....	8.0	*	*	*	*	*	*	0.2	1.1	6.2	32.1	202.1
1999 .....	8.5	*	*	*	*	*	*	0.3	1.4	6.9	34.3	215.9
Non-Hispanic black												
Both sexes												
2002 .....	2.8	*	*	*	*	*	*	1.0	3.0	8.1	37.0	118.0
2001 .....	2.7	*	*	*	*	*	*	0.7	3.4	8.9	30.7	126.1
2000 .....	3.0	*	*	*	*	*	*	1.1	3.3	9.1	35.6	140.2
1999 .....	3.1	*	*	*	*	*	*	0.7	3.3	9.3	35.0	155.8
Male												
2002 .....	2.4	*	*	*	*	*	*	1.4	4.3	10.2	44.8	109.4
2001 .....	2.2	*	*	*	*	*	*	1.0	3.9	11.0	37.2	113.2
2000 .....	2.5	*	*	*	*	*	*	1.5	4.1	11.1	40.4	150.3
1999 .....	2.6	*	*	*	*	*	*	*	4.5	12.0	44.4	166.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Atherosclerosis (I70)</b>												
Non-Hispanic black												
Female												
2002 .....	3.2	*	*	*	*	*	*	*	2.0	6.6	32.5	121.2
2001 .....	3.2	*	*	*	*	*	*	*	3.0	7.4	27.1	130.9
2000 .....	3.5	*	*	*	*	*	*	*	2.7	7.7	32.8	136.5
1999 .....	3.5	*	*	*	*	*	*	*	2.3	7.4	29.8	151.9
<b>Other diseases of circulatory system (I71-I78)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	8.6	*	*	*	0.2	0.4	1.0	2.7	8.6	29.1	68.7	148.4
2001 .....	8.9	*	*	*	0.2	0.4	1.1	2.6	8.7	29.2	71.8	158.2
2000 .....	9.3	*	*	*	0.2	0.4	1.0	2.8	9.2	32.2	74.6	163.2
1999 .....	9.2	*	*	*	0.1	0.4	1.1	2.7	9.3	31.5	75.8	165.2
Male												
2002 .....	9.4	*	*	*	0.2	0.5	1.3	3.7	11.9	39.8	92.3	187.4
2001 .....	9.6	*	*	*	0.2	0.6	1.5	3.6	12.0	40.3	97.0	197.3
2000 .....	10.2	*	*	*	0.2	0.5	1.2	3.8	13.1	45.4	102.9	214.9
1999 .....	10.1	*	*	*	0.2	0.5	1.3	3.7	13.3	44.4	104.2	207.5
Female												
2002 .....	7.9	*	*	*	0.1	0.3	0.7	1.7	5.5	20.1	53.0	131.5
2001 .....	8.2	*	*	*	0.1	0.3	0.7	1.7	5.6	19.9	55.3	141.8
2000 .....	8.3	*	*	*	0.1	0.3	0.8	1.8	5.6	21.4	56.2	142.2
1999 .....	8.4	*	*	*	*	0.2	0.8	1.7	5.6	21.0	57.3	148.2
Hispanic												
Both sexes												
2002 .....	2.2	*	*	*	*	0.3	0.8	1.7	5.9	14.8	41.0	92.6
2001 .....	2.0	*	*	*	*	*	0.7	2.1	4.8	13.7	39.0	95.0
2000 .....	2.3	*	*	*	*	0.4	0.7	1.8	5.8	20.2	42.5	102.8
1999 .....	2.1	*	*	*	*	*	0.7	1.3	5.7	16.4	44.0	110.8
Male												
2002 .....	2.3	*	*	*	*	*	1.0	2.2	7.1	19.9	56.6	102.1
2001 .....	2.2	*	*	*	*	*	0.9	2.7	6.3	20.0	53.2	108.6
2000 .....	2.5	*	*	*	*	*	0.8	2.1	8.0	25.7	62.0	131.0
1999 .....	2.4	*	*	*	*	*	0.9	1.7	7.7	22.2	62.0	163.9
Female												
2002 .....	2.0	*	*	*	*	*	*	1.1	4.9	10.8	30.4	87.9
2001 .....	1.8	*	*	*	*	*	*	1.6	3.4	8.8	29.3	88.4
2000 .....	2.1	*	*	*	*	*	*	1.5	3.9	15.8	29.4	89.0
1999 .....	1.9	*	*	*	*	*	*	*	4.0	11.9	31.9	85.3
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	9.6	*	*	*	0.2	0.4	1.0	2.8	8.8	30.0	69.8	150.3
2001 .....	9.9	*	*	*	0.2	0.5	1.2	2.6	8.9	30.1	73.2	160.1
2000 .....	10.2	*	*	*	0.2	0.4	1.0	2.8	9.4	32.9	75.8	165.0
1999 .....	10.2	*	*	*	0.1	0.4	1.1	2.8	9.5	32.3	76.9	166.8
Male												
2002 .....	10.5	*	*	*	0.2	0.5	1.3	3.9	12.3	41.1	93.9	190.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
Non-Hispanic <sup>4</sup>												
Male												
2001 .....	10.7	*	*	*	0.2	0.6	1.6	3.6	12.3	41.5	98.8	200.5
2000 .....	11.3	*	*	*	0.2	0.5	1.3	3.9	13.5	46.5	104.4	217.9
1999 .....	11.2	*	*	*	0.2	0.6	1.4	3.8	13.7	45.5	105.8	209.0
Female												
2002 .....	8.8	*	*	*	*	0.3	0.7	1.7	5.5	20.7	53.9	132.8
2001 .....	9.1	*	*	*	*	0.3	0.8	1.7	5.8	20.6	56.3	143.3
2000 .....	9.2	*	*	*	0.1	0.2	0.8	1.8	5.7	21.7	57.1	143.6
1999 .....	9.3	*	*	*	*	0.2	0.8	1.8	5.7	21.5	58.2	149.9
Non-Hispanic white												
Both sexes												
2002 .....	10.6	*	*	*	0.1	0.3	0.9	2.3	8.4	30.4	70.9	150.9
2001 .....	10.8	*	*	*	0.2	0.4	1.0	2.3	8.5	30.5	73.5	160.1
2000 .....	11.2	*	*	*	0.1	0.3	0.8	2.4	9.0	33.5	76.3	164.8
1999 .....	11.3	*	*	*	0.1	0.4	1.0	2.5	9.4	32.9	77.7	166.7
Male												
2002 .....	11.6	*	*	*	0.2	0.4	1.2	3.3	11.8	42.0	96.3	194.6
2001 .....	11.8	*	*	*	0.2	0.6	1.3	3.1	12.0	42.4	100.3	204.7
2000 .....	12.6	*	*	*	0.2	0.5	1.0	3.5	13.2	47.9	106.3	222.7
1999 .....	12.5	*	*	*	0.2	0.5	1.3	3.4	13.8	47.0	107.9	212.8
Female												
2002 .....	9.6	*	*	*	*	0.3	0.6	1.4	5.2	20.5	53.9	132.1
2001 .....	9.9	*	*	*	*	0.3	0.6	1.5	5.3	20.4	55.7	141.6
2000 .....	9.9	*	*	*	*	0.2	0.6	1.4	5.2	21.3	56.6	141.5
1999 .....	10.1	*	*	*	*	0.2	0.7	1.5	5.3	21.1	57.8	148.3
Non-Hispanic black												
Both sexes												
2002 .....	6.9	*	*	*	*	0.9	2.0	6.0	13.6	31.7	69.1	157.6
2001 .....	7.3	*	*	*	*	0.8	2.5	5.2	14.1	32.3	80.5	175.7
2000 .....	7.3	*	*	*	*	0.7	2.2	5.7	14.5	33.4	78.9	180.0
1999 .....	7.0	*	*	*	*	0.6	2.0	5.4	12.5	32.1	77.3	180.1
Male												
2002 .....	7.0	*	*	*	*	1.2	2.7	8.2	19.5	39.2	78.8	168.5
2001 .....	7.2	*	*	*	*	1.0	3.5	7.4	17.9	40.3	97.2	168.6
2000 .....	7.2	*	*	*	*	0.9	2.8	7.6	18.7	41.8	96.9	179.7
1999 .....	6.7	*	*	*	*	0.9	2.4	7.4	16.0	40.0	90.7	184.2
Female												
2002 .....	6.9	*	*	*	*	*	1.4	4.2	8.9	26.3	63.7	153.4
2001 .....	7.4	*	*	*	*	*	1.6	3.3	11.0	26.6	71.1	178.4
2000 .....	7.5	*	*	*	*	*	1.6	4.0	11.2	27.5	68.8	180.1
1999 .....	7.2	*	*	*	*	*	1.7	3.6	9.7	26.6	69.9	178.6
Aortic aneurysm and dissection (I71)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	5.1	*	*	*	0.1	0.3	0.7	1.8	5.5	19.1	43.3	69.4
2001 .....	5.3	*	*	*	0.1	0.3	0.7	1.7	5.5	19.2	45.7	77.5

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
All origins <sup>3</sup>												
Both sexes												
2000 .....	5.6	*	*	*	0.1	0.3	0.7	1.8	6.2	21.8	47.2	79.2
1999 .....	5.7	*	*	*	0.1	0.2	0.7	1.7	6.0	21.7	48.6	80.3
Male												
2002 .....	6.4	*	*	*	0.2	0.4	1.0	2.7	8.5	28.5	62.7	106.6
2001 .....	6.5	*	*	*	0.2	0.4	1.2	2.6	8.5	28.7	66.4	114.8
2000 .....	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.9	70.6	127.3
1999 .....	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.7	72.7	125.3
Female												
2002 .....	4.0	*	*	*	*	0.1	0.3	0.8	2.7	11.2	30.5	53.3
2001 .....	4.2	*	*	*	*	0.1	0.3	0.8	2.8	11.3	32.1	61.9
2000 .....	4.3	*	*	*	*	0.1	0.4	0.9	3.0	12.7	32.0	59.6
1999 .....	4.4	*	*	*	*	*	0.4	0.8	2.8	12.7	32.9	62.2
Hispanic												
Both sexes												
2002 .....	1.1	*	*	*	*	*	0.4	1.0	3.8	7.5	21.2	31.0
2001 .....	1.1	*	*	*	*	*	0.4	1.2	2.5	8.2	22.9	31.9
2000 .....	1.3	*	*	*	*	*	0.4	1.3	3.4	11.2	27.1	41.8
1999 .....	1.1	*	*	*	*	*	0.5	0.9	3.4	9.2	24.0	40.7
Male												
2002 .....	1.4	*	*	*	*	*	0.7	1.6	5.5	12.0	32.0	43.5
2001 .....	1.5	*	*	*	*	*	0.7	1.8	3.8	14.2	34.9	53.3
2000 .....	1.6	*	*	*	*	*	*	1.8	5.2	16.0	42.3	64.5
1999 .....	1.5	*	*	*	*	*	*	1.5	5.3	14.7	33.9	79.8
Female												
2002 .....	0.8	*	*	*	*	*	*	*	2.3	4.0	13.8	24.9
2001 .....	0.7	*	*	*	*	*	*	*	*	3.5	14.8	21.4
2000 .....	1.0	*	*	*	*	*	*	*	*	7.5	16.8	30.7
1999 .....	0.8	*	*	*	*	*	*	*	*	4.8	17.3	21.8
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	5.7	*	*	*	0.1	0.3	0.7	1.8	5.6	19.8	44.3	70.8
2001 .....	6.0	*	*	*	0.1	0.3	0.8	1.7	5.7	19.8	46.7	79.0
2000 .....	6.2	*	*	*	0.1	0.3	0.7	1.8	6.3	22.4	47.9	80.3
1999 .....	6.3	*	*	*	0.1	0.3	0.8	1.8	6.2	22.4	49.5	81.7
Male												
2002 .....	7.2	*	*	*	0.2	0.4	1.1	2.9	8.7	29.6	64.1	109.1
2001 .....	7.3	*	*	*	0.2	0.5	1.2	2.7	8.8	29.5	67.7	117.0
2000 .....	7.8	*	*	*	*	0.4	1.0	2.7	9.8	33.8	71.6	129.6
1999 .....	7.8	*	*	*	*	0.4	1.1	2.8	9.8	33.6	74.3	127.0
Female												
2002 .....	4.4	*	*	*	*	0.1	0.4	0.9	2.7	11.6	31.2	54.3
2001 .....	4.7	*	*	*	*	0.2	0.3	0.8	2.9	11.8	32.9	63.2
2000 .....	4.7	*	*	*	*	0.1	0.4	0.9	3.1	13.0	32.5	60.4
1999 .....	4.8	*	*	*	*	*	0.4	0.9	2.9	13.2	33.4	63.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Aortic aneurysm and dissection (I71)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	6.5	*	*	*	0.1	0.2	0.6	1.5	5.5	20.7	46.0	72.8
2001 .....	6.7	*	*	*	0.1	0.3	0.6	1.5	5.6	20.7	47.9	81.3
2000 .....	7.0	*	*	*	*	0.2	0.6	1.6	6.3	23.4	49.3	82.0
1999 .....	7.1	*	*	*	0.1	0.2	0.7	1.6	6.3	23.4	51.1	83.7
<b>Male</b>												
2002 .....	8.0	*	*	*	*	0.3	0.9	2.4	8.6	31.0	67.1	114.7
2001 .....	8.2	*	*	*	0.2	0.5	1.0	2.3	8.7	30.9	70.0	122.9
2000 .....	8.8	*	*	*	*	0.3	0.8	2.5	9.8	35.6	74.2	135.3
1999 .....	8.9	*	*	*	*	0.4	1.0	2.4	10.1	35.4	77.5	131.8
<b>Female</b>												
2002 .....	4.9	*	*	*	*	*	0.3	0.7	2.6	11.9	31.9	54.7
2001 .....	5.2	*	*	*	*	*	0.3	0.6	2.7	12.0	33.3	64.0
2000 .....	5.2	*	*	*	*	*	0.3	0.7	2.9	13.1	32.9	60.5
1999 .....	5.3	*	*	*	*	*	0.4	0.7	2.7	13.3	33.9	64.5
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	3.3	*	*	*	*	0.6	1.4	4.3	7.5	14.8	30.2	50.5
2001 .....	3.5	*	*	*	*	0.4	1.6	3.6	7.7	14.7	38.0	55.9
2000 .....	3.6	*	*	*	*	0.5	1.5	3.3	8.2	17.3	36.8	63.0
1999 .....	3.4	*	*	*	*	0.4	1.5	3.9	6.5	16.8	34.0	61.1
<b>Male</b>												
2002 .....	3.9	*	*	*	*	1.0	2.3	6.6	12.1	19.3	34.9	49.2
2001 .....	3.9	*	*	*	*	*	2.6	5.4	11.3	19.9	47.4	49.7
2000 .....	4.0	*	*	*	*	*	2.3	4.7	11.6	22.6	47.2	70.5
1999 .....	3.7	*	*	*	*	*	2.0	6.0	9.3	22.1	38.8	77.3
<b>Female</b>												
2002 .....	2.8	*	*	*	*	*	*	2.2	3.8	11.6	27.5	51.0
2001 .....	3.1	*	*	*	*	*	0.7	2.0	4.8	11.1	32.7	58.2
2000 .....	3.3	*	*	*	*	*	0.9	2.1	5.5	13.5	30.9	60.2
1999 .....	3.1	*	*	*	*	*	1.0	2.0	4.3	13.1	31.3	55.1
<b>Other diseases of arteries, arterioles and capillaries (I72-I78)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	3.5	*	*	*	0.1	0.1	0.3	0.9	3.1	10.0	25.3	79.0
2001 .....	3.5	*	*	*	*	0.1	0.4	0.9	3.2	10.0	26.1	80.7
2000 .....	3.6	*	*	*	0.1	0.1	0.3	1.0	3.1	10.4	27.4	84.0
1999 .....	3.6	*	*	*	*	0.1	0.3	0.9	3.3	9.8	27.2	84.9
<b>Male</b>												
2002 .....	3.0	*	*	*	*	*	0.3	1.0	3.4	11.3	29.6	80.8
2001 .....	3.1	*	*	*	*	0.1	0.3	0.9	3.5	11.6	30.6	82.5
2000 .....	3.2	*	*	*	*	0.1	0.3	1.1	3.6	12.5	32.3	87.6
1999 .....	3.1	*	*	*	*	0.1	0.3	1.0	3.8	11.7	31.5	82.2
<b>Female</b>												
2002 .....	4.0	*	*	*	*	0.2	0.3	0.8	2.8	8.9	22.5	78.2
2001 .....	4.0	*	*	*	*	0.1	0.4	0.9	2.9	8.6	23.1	79.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All origins <sup>3</sup>												
Female												
2002 .....	4.0	*	*	*	*	0.1	0.4	0.9	2.6	8.7	24.2	82.6
1999 .....	4.1	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.5	86.0
Hispanic												
Both sexes												
2002 .....	1.1	*	*	*	*	*	0.3	0.7	2.1	7.3	19.8	61.5
2001 .....	0.9	*	*	*	*	*	*	0.9	2.2	5.5	16.0	63.2
2000 .....	1.0	*	*	*	*	*	*	*	2.4	8.9	15.4	61.0
1999 .....	1.0	*	*	*	*	*	*	*	2.4	7.2	20.0	70.2
Male												
2002 .....	0.9	*	*	*	*	*	*	*	*	7.8	24.6	58.6
2001 .....	0.8	*	*	*	*	*	*	*	2.5	5.8	18.3	55.2
2000 .....	0.9	*	*	*	*	*	*	*	2.7	9.7	19.7	66.5
1999 .....	0.9	*	*	*	*	*	*	*	*	7.5	28.2	84.1
Female												
2002 .....	1.3	*	*	*	*	*	*	*	2.6	6.8	16.6	63.0
2001 .....	1.1	*	*	*	*	*	*	*	*	5.4	14.5	67.0
2000 .....	1.1	*	*	*	*	*	*	*	*	8.3	12.5	58.4
1999 .....	1.1	*	*	*	*	*	*	*	2.4	7.1	14.6	63.4
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	3.9	*	*	*	0.1	0.1	0.3	0.9	3.2	10.2	25.6	79.5
2001 .....	3.9	*	*	*	*	0.1	0.4	0.9	3.2	10.3	26.5	81.1
2000 .....	4.0	*	*	*	0.1	0.1	0.3	1.0	3.1	10.5	27.9	84.7
1999 .....	3.9	*	*	*	*	0.1	0.3	1.0	3.3	9.9	27.5	85.1
Male												
2002 .....	3.4	*	*	*	*	*	0.3	1.0	3.5	11.5	29.8	81.7
2001 .....	3.4	*	*	*	*	0.1	0.3	0.9	3.5	12.0	31.1	83.5
2000 .....	3.5	*	*	*	*	0.1	0.2	1.2	3.6	12.7	32.8	88.2
1999 .....	3.4	*	*	*	*	0.1	0.3	1.0	3.9	11.9	31.5	82.0
Female												
2002 .....	4.3	*	*	*	*	0.2	0.3	0.8	2.8	9.0	22.8	78.5
2001 .....	4.4	*	*	*	*	0.1	0.4	0.9	2.9	8.8	23.5	80.1
2000 .....	4.4	*	*	*	*	*	0.4	0.9	2.6	8.7	24.6	83.3
1999 .....	4.5	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.8	86.4
Non-Hispanic white												
Both sexes												
2002 .....	4.1	*	*	*	*	0.1	0.3	0.8	2.9	9.7	24.9	78.2
2001 .....	4.2	*	*	*	*	0.1	0.3	0.8	2.9	9.8	25.6	78.8
2000 .....	4.3	*	*	*	*	0.1	0.3	0.9	2.8	10.1	27.1	82.8
1999 .....	4.2	*	*	*	*	0.1	0.3	0.9	3.1	9.6	26.5	83.0
Male												
2002 .....	3.6	*	*	*	*	*	0.2	0.9	3.2	11.0	29.3	79.9
2001 .....	3.6	*	*	*	*	*	0.2	0.8	3.3	11.5	30.3	81.8
2000 .....	3.8	*	*	*	*	*	0.2	1.0	3.3	12.3	32.1	87.4
1999 .....	3.6	*	*	*	*	*	0.3	1.0	3.6	11.6	30.4	81.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Non-Hispanic white												
Female												
2002 .....	4.7	*	*	*	*	*	0.3	0.7	2.6	8.6	21.9	77.4
2001 .....	4.7	*	*	*	*	*	0.4	0.9	2.6	8.3	22.4	77.6
2000 .....	4.7	*	*	*	*	*	0.3	0.7	2.2	8.2	23.8	81.0
1999 .....	4.8	*	*	*	*	*	0.4	0.8	2.5	7.9	24.0	83.9
Non-Hispanic black												
Both sexes												
2002 .....	3.6	*	*	*	*	*	0.6	1.8	6.2	16.8	39.0	107.1
2001 .....	3.8	*	*	*	*	*	0.9	1.7	6.4	17.5	42.5	119.9
2000 .....	3.7	*	*	*	*	*	0.6	2.4	6.3	16.1	42.1	117.0
1999 .....	3.6	*	*	*	*	*	0.6	1.5	5.9	15.3	43.3	119.0
Male												
2002 .....	3.1	*	*	*	*	*	*	1.6	7.4	19.9	44.0	119.2
2001 .....	3.3	*	*	*	*	*	0.9	2.0	6.6	20.5	49.8	119.0
2000 .....	3.2	*	*	*	*	*	*	2.9	7.1	19.2	49.7	109.2
1999 .....	3.0	*	*	*	*	*	*	1.4	6.7	17.9	51.9	107.0
Female												
2002 .....	4.0	*	*	*	*	*	0.8	1.9	5.1	14.6	36.1	102.4
2001 .....	4.4	*	*	*	*	*	0.9	1.4	6.3	15.5	38.4	120.2
2000 .....	4.2	*	*	*	*	*	0.7	1.9	5.7	14.0	37.9	119.9
1999 .....	4.2	*	*	*	*	*	0.8	1.6	5.4	13.5	38.5	123.5
Other disorders of circulatory system (I80-I99)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.2	3.8	9.8	24.4
2001 .....	1.6	0.5	*	*	0.1	0.3	0.6	1.2	2.2	4.6	9.4	26.1
2000 .....	1.6	0.7	*	*	0.1	0.3	0.7	1.4	2.1	4.3	9.4	25.6
1999 .....	1.4	0.6	*	*	0.1	0.2	0.6	1.0	1.9	3.8	9.1	23.4
Male												
2002 .....	1.4	*	*	*	0.1	0.3	0.7	1.4	2.4	3.9	9.5	20.5
2001 .....	1.4	*	*	*	*	0.3	0.7	1.4	2.5	4.9	9.8	22.0
2000 .....	1.4	*	*	*	*	0.3	0.8	1.6	2.3	4.7	9.7	22.2
1999 .....	1.3	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.6	20.8
Female												
2002 .....	1.9	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.1	26.2
2001 .....	1.9	*	*	*	0.1	0.4	0.6	1.0	2.0	4.4	9.2	27.8
2000 .....	1.8	*	*	*	0.1	0.4	0.6	1.1	2.0	4.0	9.2	26.9
1999 .....	1.6	*	*	*	*	0.2	0.5	1.0	1.8	3.4	8.8	24.5
Hispanic												
Both sexes												
2002 .....	0.6	*	*	*	*	*	0.4	1.0	1.4	3.1	6.3	18.3
2001 .....	0.6	*	*	*	*	*	*	0.9	*	4.2	7.4	32.5
2000 .....	0.6	*	*	*	*	*	0.6	1.1	1.3	3.3	8.7	24.6
1999 .....	0.5	*	*	*	*	*	*	1.1	1.5	4.1	5.4	24.6
Male												
2002 .....	0.5	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
Hispanic												
Male												
2001 .....	0.5	*	*	*	*	*	*	1.3	*	4.0	8.9	*
2000 .....	0.6	*	*	*	*	*	0.8	*	*	*	10.8	*
1999 .....	0.5	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.7	*	*	*	*	*	*	*	*	3.2	6.2	23.2
2001 .....	0.6	*	*	*	*	*	*	*	*	4.4	6.3	33.5
2000 .....	0.7	*	*	*	*	*	*	*	*	*	7.3	23.7
1999 .....	0.6	*	*	*	*	*	*	*	*	4.3	*	23.9
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.8	0.6	*	*	0.1	0.4	0.7	1.4	2.3	3.9	10.0	24.7
2001 .....	1.8	*	*	*	0.1	0.4	0.7	1.2	2.3	4.6	9.5	25.8
2000 .....	1.8	0.7	*	*	0.1	0.4	0.7	1.4	2.2	4.4	9.4	25.6
1999 .....	1.6	0.7	*	*	0.1	0.3	0.6	1.0	1.9	3.7	9.3	23.3
Male												
2002 .....	1.5	*	*	*	*	0.4	0.7	1.5	2.5	3.9	9.6	21.1
2001 .....	1.5	*	*	*	*	0.3	0.7	1.4	2.6	4.9	9.8	21.6
2000 .....	1.5	*	*	*	*	0.3	0.8	1.6	2.3	4.8	9.6	22.1
1999 .....	1.4	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.7	20.5
Female												
2002 .....	2.1	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.2	26.2
2001 .....	2.0	*	*	*	0.1	0.4	0.6	1.0	2.1	4.3	9.2	27.6
2000 .....	2.0	*	*	*	*	0.4	0.6	1.2	2.1	4.1	9.3	27.0
1999 .....	1.8	*	*	*	*	0.3	0.5	0.9	1.8	3.3	9.0	24.4
Non-Hispanic white												
Both sexes												
2002 .....	1.8	*	*	*	*	0.3	0.6	1.2	2.1	3.6	9.6	24.5
2001 .....	1.8	*	*	*	*	0.3	0.6	1.0	2.1	4.3	9.2	25.7
2000 .....	1.8	*	*	*	*	0.3	0.6	1.2	1.9	4.0	9.2	25.5
1999 .....	1.6	*	*	*	*	0.2	0.5	0.9	1.8	3.5	9.0	23.0
Male												
2002 .....	1.5	*	*	*	*	0.3	0.6	1.3	2.2	3.7	9.1	20.5
2001 .....	1.5	*	*	*	*	0.2	0.6	1.2	2.3	4.6	9.4	21.2
2000 .....	1.5	*	*	*	*	0.2	0.7	1.5	2.0	4.5	9.4	21.8
1999 .....	1.4	*	*	*	*	0.2	0.6	0.9	2.0	3.9	9.3	20.3
Female												
2002 .....	2.2	*	*	*	*	0.3	0.6	1.1	1.9	3.6	9.9	26.3
2001 .....	2.1	*	*	*	*	0.3	0.5	0.8	1.9	4.1	9.0	27.6
2000 .....	2.0	*	*	*	*	0.3	0.5	1.0	1.9	3.6	9.0	27.0
1999 .....	1.8	*	*	*	*	0.2	0.5	0.8	1.7	3.1	8.8	24.1
Non-Hispanic black												
Both sexes												
2002 .....	2.2	*	*	*	*	1.0	1.7	2.7	5.0	6.9	16.8	29.9
2001 .....	2.0	*	*	*	*	0.8	1.6	2.3	4.9	7.9	14.3	29.7
2000 .....	2.1	*	*	*	*	0.9	1.5	2.8	4.6	9.0	14.4	29.9
1999 .....	1.8	*	*	*	*	0.5	1.5	2.4	3.0	6.8	14.0	28.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Other disorders of circulatory system (I80-I99)</b>												
<b>Non-Hispanic black</b>												
<b>Male</b>												
2002 .....	2.1	*	*	*	*	1.1	1.8	3.0	5.6	7.8	18.5	31.7
2001 .....	2.0	*	*	*	*	*	1.8	2.4	6.5	9.2	15.9	28.9
2000 .....	2.0	*	*	*	*	0.9	1.8	3.5	5.3	9.1	13.3	30.5
1999 .....	1.7	*	*	*	*	*	1.9	2.9	2.7	8.4	17.2	*
<b>Female</b>												
2002 .....	2.3	*	*	*	*	1.0	1.6	2.4	4.5	6.2	15.9	29.3
2001 .....	2.1	*	*	*	*	0.9	1.4	2.1	3.7	7.0	13.5	29.9
2000 .....	2.3	*	*	*	*	1.0	1.3	2.3	4.0	8.8	15.1	29.7
1999 .....	1.8	*	*	*	*	*	1.1	2.0	3.2	5.6	12.3	30.6
<b>Influenza and pneumonia (J10-J18)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	22.8	6.5	0.7	0.2	0.4	0.9	2.2	4.8	11.2	37.5	156.9	696.6
2001 .....	21.8	7.4	0.7	0.2	0.5	0.9	2.2	4.6	10.7	36.3	148.5	685.6
2000 .....	23.2	7.6	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.1	160.3	744.1
1999 .....	22.8	8.4	0.8	0.2	0.5	0.8	2.4	4.6	11.0	37.2	157.0	751.8
<b>Male</b>												
2002 .....	20.4	7.8	0.8	0.2	0.5	0.9	2.4	5.8	13.6	46.4	194.7	805.8
2001 .....	19.6	8.1	0.7	0.2	0.5	1.0	2.5	5.7	13.1	45.0	186.4	808.4
2000 .....	20.8	8.4	0.7	0.2	0.5	1.0	2.8	5.8	14.4	48.3	199.8	892.0
1999 .....	20.3	8.8	0.8	0.3	0.5	1.0	2.6	5.9	12.8	46.9	196.5	894.7
<b>Female</b>												
2002 .....	25.1	5.1	0.6	0.3	0.4	0.8	1.9	3.8	9.0	30.1	131.8	649.2
2001 .....	23.9	6.7	0.7	0.2	0.4	0.8	1.9	3.5	8.5	29.1	123.7	634.2
2000 .....	25.6	6.8	0.7	0.2	0.5	0.8	1.9	3.7	9.5	31.5	134.6	683.9
1999 .....	25.3	8.0	0.9	0.2	0.5	0.7	2.1	3.4	9.4	29.3	131.3	694.2
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	7.3	5.8	*	*	*	0.6	1.3	3.4	10.5	33.8	142.6	566.0
2001 .....	7.4	7.0	1.1	*	*	0.3	1.3	3.6	9.1	35.4	137.2	660.3
2000 .....	7.4	7.8	*	0.3	0.3	0.7	1.6	3.8	11.2	37.6	136.3	643.6
1999 .....	6.6	8.5	*	*	0.4	0.4	1.5	3.5	9.5	31.7	118.5	614.6
<b>Male</b>												
2002 .....	7.0	7.7	*	*	*	0.7	1.7	4.7	13.9	43.8	173.2	655.9
2001 .....	6.9	7.4	*	*	*	*	1.7	5.0	11.4	40.7	173.5	765.6
2000 .....	7.2	7.6	*	*	*	0.8	2.1	5.3	14.8	46.0	180.4	763.9
1999 .....	6.3	10.3	*	*	*	*	2.1	4.9	10.4	41.1	148.1	720.3
<b>Female</b>												
2002 .....	7.6	*	*	*	*	*	0.9	2.1	7.4	25.9	121.7	521.4
2001 .....	7.9	6.5	*	*	*	*	1.0	2.1	7.1	31.2	112.7	608.8
2000 .....	7.7	8.0	*	*	*	0.7	1.0	2.4	7.9	31.0	106.6	584.6
1999 .....	7.0	6.5	*	*	*	*	0.8	2.2	8.6	24.3	98.9	563.6
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	25.1	6.7	0.7	0.2	0.4	0.9	2.3	4.9	11.3	37.6	157.1	700.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Influenza and pneumonia (J10-J18)</b>												
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	23.8	7.5	0.6	0.2	0.5	1.0	2.3	4.7	10.7	36.1	148.6	684.8
2001 .....	25.4	7.5	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.0	160.7	745.7
2000 .....	25.0	8.4	0.9	0.2	0.5	0.9	2.4	4.7	11.0	37.4	158.0	754.0
1999 .....	25.0	8.4	0.9	0.2	0.5	0.9	2.4	4.7	11.0	37.4	158.0	754.0
<b>Male</b>												
2002 .....	22.5	7.8	0.8	0.2	0.5	1.0	2.5	5.8	13.5	46.3	195.1	810.7
2001 .....	21.5	8.2	0.6	0.2	0.5	1.1	2.6	5.7	13.1	44.9	186.5	808.2
2000 .....	22.7	8.5	0.7	0.2	0.6	1.1	2.9	5.7	14.3	48.2	199.7	894.9
1999 .....	22.2	8.4	0.9	0.3	0.5	1.1	2.7	5.9	12.9	47.0	197.7	897.9
<b>Female</b>												
2002 .....	27.5	5.4	0.7	0.3	0.4	0.8	2.0	4.0	9.2	30.3	131.8	652.6
2001 .....	26.1	6.7	0.6	0.2	0.4	0.8	2.0	3.6	8.5	28.9	123.8	633.5
2000 .....	27.9	6.5	0.6	0.2	0.5	0.8	2.0	3.8	9.6	31.4	135.3	685.4
1999 .....	27.6	8.4	0.9	0.2	0.5	0.8	2.2	3.5	9.4	29.5	132.2	696.4
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	27.9	4.9	0.5	0.2	0.4	0.8	2.0	4.4	10.3	36.7	156.9	712.7
2001 .....	26.2	4.9	0.5	0.2	0.4	0.8	1.9	4.0	9.9	35.0	147.2	693.9
2000 .....	27.9	5.0	0.5	0.2	0.5	0.8	2.1	4.1	10.7	36.8	160.2	759.8
1999 .....	27.5	5.3	0.6	0.2	0.4	0.8	2.0	4.0	10.0	36.1	158.8	766.9
<b>Male</b>												
2002 .....	24.6	6.1	0.5	0.2	0.5	0.9	2.2	5.2	12.1	44.9	193.2	820.6
2001 .....	23.1	5.8	0.6	0.2	0.5	0.9	2.1	4.8	11.7	42.7	182.5	813.9
2000 .....	24.5	5.8	0.5	0.2	0.5	0.9	2.4	4.7	12.6	44.7	197.5	908.1
1999 .....	23.9	5.5	0.5	0.3	0.4	0.9	2.1	4.8	11.5	44.6	197.3	909.3
<b>Female</b>												
2002 .....	31.1	3.6	0.5	0.3	0.3	0.8	1.8	3.7	8.6	29.7	132.5	666.1
2001 .....	29.2	3.9	0.5	0.2	0.4	0.8	1.7	3.3	8.1	28.5	123.7	644.0
2000 .....	31.3	4.2	0.6	0.2	0.4	0.8	1.8	3.4	9.0	30.1	135.7	700.0
1999 .....	31.0	5.1	0.8	0.2	0.4	0.6	1.9	3.1	8.7	29.0	133.5	710.2
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	16.1	13.3	1.7	*	0.6	1.6	4.5	8.9	21.6	50.9	167.7	579.8
2001 .....	16.0	16.9	1.0	*	0.8	1.7	4.9	9.9	19.1	48.8	169.5	583.1
2000 .....	16.9	17.5	1.2	*	0.8	1.8	4.6	10.3	22.3	60.8	172.2	604.5
1999 .....	16.7	20.0	1.7	*	0.9	2.0	5.5	10.8	20.9	54.7	158.3	634.7
<b>Male</b>												
2002 .....	15.9	14.2	1.8	*	0.7	1.8	5.2	11.5	29.6	66.5	231.2	695.8
2001 .....	16.3	16.6	*	*	0.8	2.1	5.9	13.5	26.6	70.0	242.5	718.4
2000 .....	17.3	18.8	1.7	*	0.8	2.4	5.8	14.3	30.3	84.1	235.4	763.2
1999 .....	16.9	20.1	2.1	*	0.8	2.3	6.7	15.3	27.3	77.9	215.1	803.4
<b>Female</b>												
2002 .....	16.2	12.3	*	*	*	1.4	3.9	6.6	15.2	39.8	131.8	535.5
2001 .....	15.6	17.1	*	*	0.8	1.3	4.0	6.9	13.0	33.9	128.5	533.0
2000 .....	16.5	16.2	*	*	0.8	1.2	3.5	6.9	15.9	44.4	136.7	545.6
1999 .....	16.5	20.0	*	*	0.9	1.8	4.5	7.0	15.8	38.4	126.4	571.6

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Influenza (J10-J11)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	8.8
2001 .....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.2
2000 .....	0.6	*	*	*	*	*	0.0	0.1	0.4	1.1	4.0	20.2
1999 .....	0.6	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.2
<b>Male</b>												
2002 .....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	8.9
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2000 .....	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.0	20.7
1999 .....	0.4	*	*	*	*	*	*	*	0.3	0.9	4.2	19.9
<b>Female</b>												
2002 .....	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.7
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000 .....	0.8	*	*	*	*	*	*	0.1	0.4	0.9	4.0	19.9
1999 .....	0.7	*	*	*	*	*	*	*	0.3	0.9	3.4	21.7
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	9.0
2001 .....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.3
2000 .....	0.7	*	*	*	*	*	0.0	0.1	0.4	1.1	4.1	20.5
1999 .....	0.7	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.6
<b>Male</b>												
2002 .....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	9.2
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2000 .....	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.1	21.1
1999 .....	0.5	*	*	*	*	*	*	*	0.3	0.8	4.3	20.7
<b>Female</b>												
2002 .....	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.9
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000 .....	0.9	*	*	*	*	*	*	0.1	0.4	1.0	4.1	20.3
1999 .....	0.8	*	*	*	*	*	*	*	0.3	0.9	3.4	22.0

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Influenza (J10-J11)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.6	9.6
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000 .....	0.8	*	*	*	*	*	*	0.1	0.4	1.2	4.3	21.7
1999 .....	0.8	*	*	*	*	*	*	0.1	0.3	0.9	3.9	23.3
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	*	*	*	0.4	1.7	10.0
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2000 .....	0.6	*	*	*	*	*	*	*	0.4	1.4	4.3	22.0
1999 .....	0.6	*	*	*	*	*	*	*	0.3	0.9	4.6	22.5
<b>Female</b>												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	1.4	9.5
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000 .....	1.0	*	*	*	*	*	*	0.1	0.5	1.0	4.3	21.6
1999 .....	1.0	*	*	*	*	*	*	*	0.3	1.0	3.5	23.5
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	9.5
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
<b>Pneumonia (J12-J18)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	22.5	6.3	0.7	0.2	0.4	0.9	2.1	4.7	11.1	37.2	155.5	687.8
2001 .....	21.7	7.2	0.7	0.2	0.4	0.8	2.2	4.6	10.6	36.2	148.1	683.4
2000 .....	22.6	7.4	0.6	0.2	0.4	0.9	2.3	4.6	11.5	38.0	156.4	723.9
1999 .....	22.2	8.1	0.8	0.2	0.4	0.8	2.3	4.6	10.7	36.4	153.3	730.6
<b>Male</b>												
2002 .....	20.2	7.6	0.8	0.2	0.4	0.9	2.4	5.7	13.4	46.0	193.1	796.9
2001 .....	19.5	7.8	0.7	0.2	0.5	0.9	2.5	5.7	13.0	44.9	186.0	806.2
2000 .....	20.3	8.1	0.6	0.2	0.5	1.0	2.7	5.6	14.0	47.1	195.8	871.3
1999 .....	19.8	8.5	0.7	0.2	0.4	0.9	2.6	5.8	12.5	46.0	192.3	874.7
<b>Female</b>												
2002 .....	24.8	5.0	0.6	0.2	0.3	0.8	1.9	3.8	9.0	29.8	130.6	640.5
2001 .....	23.8	6.7	0.7	0.2	0.4	0.7	1.9	3.5	8.4	29.0	123.2	632.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
All origins <sup>3</sup>												
Female												
2000 .....	24.8	6.6	0.6	0.2	0.4	0.8	1.9	3.6	9.1	30.6	130.6	663.9
1999 .....	24.6	7.7	0.8	0.2	0.4	0.7	2.0	3.4	9.1	28.4	128.0	672.5
Hispanic												
Both sexes												
2002 .....	7.2	5.5	*	*	*	0.6	1.3	3.4	10.5	33.6	142.1	563.2
2001 .....	7.3	7.0	1.0	*	*	0.3	1.3	3.5	9.1	35.3	136.3	660.3
2000 .....	7.3	7.5	*	*	0.3	0.7	1.6	3.8	11.0	37.2	134.3	633.0
1999 .....	6.5	7.9	*	*	0.4	0.4	1.5	3.4	9.2	30.9	116.7	607.6
Male												
2002 .....	6.9	7.3	*	*	*	0.6	1.7	4.7	13.9	43.6	172.8	654.2
2001 .....	6.8	7.4	*	*	*	1.7	5.0	11.4	40.7	172.6	765.6	
2000 .....	7.1	7.6	*	*	*	0.7	2.1	5.2	14.6	45.6	178.0	751.8
1999 .....	6.1	10.3	*	*	*	*	2.0	4.7	9.9	39.5	147.0	718.2
Female												
2002 .....	7.5	*	*	*	*	*	0.9	2.1	7.4	25.7	121.1	518.1
2001 .....	7.8	6.5	*	*	*	*	0.9	2.1	7.1	31.0	111.8	608.8
2000 .....	7.6	7.4	*	*	*	*	1.0	2.3	7.9	30.7	105.0	574.7
1999 .....	6.8	5.4	*	*	*	*	0.8	2.2	8.5	24.1	96.4	554.2
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	24.8	6.5	0.7	0.2	0.4	0.9	2.3	4.9	11.1	37.3	155.6	691.2
2001 .....	23.8	7.2	0.6	0.2	0.5	0.9	2.3	4.6	10.6	36.0	148.2	682.5
2000 .....	24.7	7.3	0.6	0.2	0.5	0.9	2.4	4.6	11.4	37.9	156.6	725.2
1999 .....	24.3	8.1	0.8	0.2	0.4	0.9	2.4	4.6	10.7	36.5	154.2	732.4
Male												
2002 .....	22.3	7.6	0.8	0.2	0.5	1.0	2.5	5.8	13.4	45.9	193.5	801.5
2001 .....	21.4	7.8	0.6	0.2	0.5	1.0	2.6	5.7	13.0	44.8	186.1	805.9
2000 .....	22.2	8.2	0.7	0.2	0.5	1.0	2.8	5.6	13.9	46.8	195.7	873.8
1999 .....	21.7	8.0	0.8	0.2	0.5	1.1	2.6	5.8	12.5	46.2	193.4	877.2
Female												
2002 .....	27.2	5.3	0.6	0.2	0.4	0.8	2.0	4.0	9.1	30.1	130.6	643.7
2001 .....	26.0	6.6	0.6	0.2	0.4	0.8	2.0	3.6	8.5	28.8	123.3	631.3
2000 .....	27.1	6.4	0.5	0.1	0.4	0.8	2.0	3.7	9.2	30.4	131.2	665.2
1999 .....	26.8	8.2	0.9	0.2	0.4	0.8	2.2	3.5	9.1	28.5	128.8	674.4
Non-Hispanic white												
Both sexes												
2002 .....	27.6	4.7	0.5	0.2	0.4	0.8	2.0	4.4	10.2	36.4	155.3	703.1
2001 .....	26.1	4.6	0.5	0.2	0.4	0.8	1.9	4.0	9.8	34.9	146.7	691.5
2000 .....	27.1	4.8	0.5	0.2	0.4	0.8	2.1	3.9	10.3	35.6	155.9	738.1
1999 .....	26.8	5.2	0.6	0.2	0.4	0.7	1.9	3.9	9.7	35.2	154.8	743.7
Male												
2002 .....	24.3	6.0	0.5	0.2	0.5	0.9	2.2	5.1	11.9	44.5	191.5	810.6
2001 .....	23.0	5.3	0.6	0.2	0.5	0.9	2.1	4.8	11.6	42.6	182.0	811.5
2000 .....	23.8	5.4	0.4	0.2	0.5	0.8	2.4	4.6	12.2	43.3	193.2	886.1
1999 .....	23.4	5.4	*	0.2	0.4	0.9	2.0	4.7	11.1	43.8	192.7	886.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Pneumonia (J12-J18)</b>												
<b>Non-Hispanic white</b>												
Female												
2002 .....	30.7	3.4	0.5	0.2	0.3	0.8	1.8	3.7	8.5	29.5	131.1	656.6
2001 .....	29.1	3.8	0.4	0.2	0.3	0.7	1.7	3.2	8.0	28.3	123.3	641.7
2000 .....	30.3	4.2	0.5	*	0.3	0.8	1.7	3.3	8.5	29.1	131.4	678.4
1999 .....	30.0	4.9	0.7	0.2	0.3	0.6	1.8	3.0	8.4	28.0	130.0	686.6
<b>Non-Hispanic black</b>												
Both sexes												
2002 .....	16.0	13.1	1.5	*	0.6	1.6	4.5	8.9	21.4	50.5	167.3	577.4
2001 .....	15.9	16.9	1.0	*	0.8	1.7	4.9	9.9	19.0	48.7	169.1	581.5
2000 .....	16.6	17.2	1.2	*	0.8	1.8	4.5	10.1	22.1	59.9	170.8	595.0
1999 .....	16.5	19.3	1.7	*	0.9	2.0	5.4	10.7	20.7	54.0	156.6	628.9
Male												
2002 .....	15.8	13.9	1.6	*	*	1.8	5.2	11.5	29.3	66.0	230.6	693.6
2001 .....	16.3	16.6	*	*	0.8	2.0	5.9	13.5	26.4	69.7	242.5	716.1
2000 .....	17.1	18.8	1.7	*	0.7	2.3	5.7	14.0	30.1	83.4	233.9	750.3
1999 .....	16.8	19.1	2.0	*	0.8	2.3	6.6	15.1	27.2	77.0	213.3	799.9
Female												
2002 .....	16.1	12.3	*	*	*	1.4	3.9	6.6	15.1	39.5	131.5	533.0
2001 .....	15.6	17.1	*	*	0.8	1.3	4.0	6.9	13.0	33.9	127.9	531.7
2000 .....	16.3	15.5	*	*	0.8	1.2	3.5	6.8	15.6	43.3	135.3	537.3
1999 .....	16.3	19.6	*	*	0.9	1.7	4.4	6.9	15.5	37.7	124.8	565.0
<b>Other acute lower respiratory infections (J20-J22)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.1	1.3	0.1	*	*	*	*	*	*	0.2	0.5	3.3
2001 .....	0.1	1.2	*	*	*	*	*	*	*	0.2	0.5	3.3
2000 .....	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999 .....	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.0
Male												
2002 .....	0.1	1.9	*	*	*	*	*	*	*	*	0.5	2.8
2001 .....	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2000 .....	0.1	1.3	*	*	*	*	*	*	*	0.3	0.7	3.8
1999 .....	0.1	1.9	*	*	*	*	*	*	*	*	1.1	4.0
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.5
2001 .....	0.2	1.2	*	*	*	*	*	*	*	*	0.5	3.7
2000 .....	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.8
1999 .....	0.2	1.5	*	*	*	*	*	*	*	*	0.8	6.8
<b>Hispanic</b>												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
Hispanic												
Male												
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.1	1.3	0.2	*	*	*	*	*	*	0.2	0.5	3.4
2001 .....	0.1	1.3	*	*	*	*	*	*	*	0.2	0.4	3.3
2000 .....	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999 .....	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.1
Male												
2002 .....	0.1	2.1	*	*	*	*	*	*	*	*	0.6	2.9
2001 .....	0.1	1.2	*	*	*	*	*	*	*	*	*	2.4
2000 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.7
1999 .....	0.2	2.0	*	*	*	*	*	*	*	*	1.1	4.1
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.6
2001 .....	0.2	1.4	*	*	*	*	*	*	*	*	0.5	3.7
2000 .....	0.2	*	*	*	*	*	*	*	*	0.2	0.7	4.8
1999 .....	0.3	1.5	*	*	*	*	*	*	*	*	0.8	6.9
Non-Hispanic white												
Both sexes												
2002 .....	0.2	1.1	*	*	*	*	*	*	*	0.2	0.5	3.5
2001 .....	0.1	1.0	*	*	*	*	*	*	*	0.1	0.5	3.5
2000 .....	0.2	*	*	*	*	*	*	*	*	0.3	0.7	4.7
1999 .....	0.2	1.1	*	*	*	*	*	*	*	0.2	0.9	6.4
Male												
2002 .....	0.1	1.8	*	*	*	*	*	*	*	*	0.6	3.0
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	2.5
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.7
1999 .....	0.2	*	*	*	*	*	*	*	*	*	1.1	4.1
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	0.5	3.7
2001 .....	0.2	*	*	*	*	*	*	*	*	*	0.5	3.9
2000 .....	0.2	*	*	*	*	*	*	*	*	0.3	0.7	5.1
1999 .....	0.3	*	*	*	*	*	*	*	*	*	0.8	7.3
Non-Hispanic black												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	4.2	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
Non-Hispanic black												
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Acute bronchitis and bronchiolitis (J20-J21)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.1	1.2	0.1	*	*	*	*	*	*	0.1	0.3	2.0
2001 .....	0.1	1.2	*	*	*	*	*	*	*	0.1	0.3	2.1
2000 .....	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999 .....	0.1	1.6	0.2	*	*	*	*	*	*	0.2	0.6	3.6
Male												
2002 .....	0.1	1.8	*	*	*	*	*	*	*	*	*	1.7
2001 .....	0.1	1.2	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	1.2	*	*	*	*	*	*	*	*	0.6	2.4
1999 .....	0.1	1.9	*	*	*	*	*	*	*	*	0.7	2.6
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2001 .....	0.1	1.2	*	*	*	*	*	*	*	*	0.4	2.4
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.7
1999 .....	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.1	1.3	0.2	*	*	*	*	*	*	0.1	0.3	2.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2001 .....	0.1	1.3	*	*	*	*	*	*	*	0.1	0.3	2.1
2000 .....	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999 .....	0.1	1.7	0.2	*	*	*	*	*	*	0.2	0.6	3.6
Male												
2002 .....	0.1	2.0	*	*	*	*	*	*	*	*	*	1.7
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
1999 .....	0.1	2.0	*	*	*	*	*	*	*	*	0.7	2.6
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2001 .....	0.1	1.4	*	*	*	*	*	*	*	*	0.4	2.4
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
1999 .....	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0
Non-Hispanic white												
Both sexes												
2002 .....	0.1	1.0	*	*	*	*	*	*	*	0.1	0.3	2.1
2001 .....	0.1	1.0	*	*	*	*	*	*	*	*	0.3	2.2
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	2.7
1999 .....	0.2	1.1	*	*	*	*	*	*	*	0.1	0.6	3.9
Male												
2002 .....	0.1	1.7	*	*	*	*	*	*	*	*	*	1.7
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.6	2.3
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.8	2.6
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.8
1999 .....	0.2	*	*	*	*	*	*	*	*	*	0.5	4.4
Non-Hispanic black												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	4.2	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.9
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4
Male												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.2	1.9
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4
Male												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Unspecified acute lower respiratory infection (J22)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.4
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.5
<b>Male</b>												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	3.0
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Chronic lower respiratory diseases (J40-J47)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	43.3	1.0	0.4	0.3	0.5	0.8	2.2	8.7	42.4	163.0	386.7	637.6
2001 .....	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.5	44.1	167.9	379.8	644.7
2000 .....	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6
1999 .....	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0
<b>Male</b>												
2002 .....	42.9	1.1	0.6	0.4	0.6	0.8	2.2	9.1	45.2	184.8	480.8	894.8
2001 .....	42.7	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.9	191.3	475.1	916.9
2000 .....	43.5	1.2	0.4	0.4	0.6	0.8	1.9	9.0	47.8	195.2	488.5	967.9
1999 .....	45.6	*	0.4	0.4	0.6	0.8	1.8	8.6	52.3	210.7	513.2	996.7
<b>Female</b>												
2002 .....	43.7	*	0.3	0.3	0.4	0.7	2.3	8.2	39.8	144.9	324.1	526.0
2001 .....	43.7	*	*	0.2	0.4	0.7	2.3	8.1	41.5	148.5	317.3	530.8

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
All origins <sup>3</sup>												
Female												
2002 .....	43.2	*	0.3	0.3	0.4	0.7	2.2	8.3	41.0	148.2	319.2	518.5
1999 .....	43.4	*	0.3	0.2	0.5	0.9	2.1	8.4	43.1	149.8	322.9	504.6
Hispanic												
Both sexes												
2002 .....	7.9	*	*	*	*	0.4	1.2	3.5	13.9	53.8	176.0	457.3
2001 .....	7.7	*	*	*	*	0.3	1.0	3.6	14.0	55.8	169.5	480.2
2000 .....	7.6	*	*	*	*	0.3	1.1	3.3	14.3	53.1	181.9	481.1
1999 .....	8.4	*	*	*	*	0.5	1.6	3.9	14.6	62.6	194.8	536.0
Male												
2002 .....	8.1	*	*	*	*	*	1.0	3.8	17.5	69.2	243.3	602.4
2001 .....	7.8	*	*	*	*	*	0.7	3.2	16.1	75.5	224.0	676.1
2000 .....	8.0	*	*	*	*	*	0.9	3.4	18.2	72.4	250.3	671.1
1999 .....	8.9	*	*	*	*	*	1.4	3.6	17.6	81.9	272.7	836.8
Female												
2002 .....	7.6	*	*	*	*	*	1.4	3.1	10.6	41.5	129.8	385.5
2001 .....	7.5	*	*	*	*	*	1.2	4.1	12.1	40.3	132.7	384.4
2000 .....	7.2	*	*	*	*	*	1.3	3.3	10.8	38.0	136.0	387.8
1999 .....	8.0	*	*	*	*	*	1.8	4.2	12.0	47.5	142.9	391.0
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	48.6	1.1	0.4	0.4	0.5	0.8	2.4	9.2	44.6	170.1	396.0	643.5
2001 .....	48.4	1.1	0.3	0.3	0.5	0.8	2.3	8.9	46.3	174.8	388.6	649.4
2000 .....	48.3	1.0	0.3	0.4	0.5	0.8	2.2	9.0	46.3	176.1	393.8	653.2
1999 .....	49.4	1.0	0.4	0.4	0.6	0.9	2.0	8.9	49.8	183.5	405.1	648.0
Male												
2002 .....	48.4	1.2	0.6	0.4	0.6	0.9	2.4	9.6	47.3	191.9	491.2	906.0
2001 .....	48.0	*	0.4	0.3	0.6	0.8	2.2	9.3	49.0	198.1	485.7	924.8
2000 .....	48.7	*	0.4	0.4	0.6	0.9	2.1	9.4	49.7	201.9	497.6	978.0
1999 .....	50.8	*	0.4	0.5	0.7	0.8	1.9	9.0	54.5	217.4	521.9	1,000.2
Female												
2002 .....	48.9	*	*	0.3	0.4	0.8	2.4	8.7	42.0	151.8	332.9	530.3
2001 .....	48.7	*	*	0.2	0.4	0.8	2.4	8.5	43.8	155.4	324.9	534.8
2000 .....	48.0	*	*	0.3	0.4	0.7	2.3	8.7	43.2	154.8	326.2	521.9
1999 .....	48.0	*	*	0.2	0.5	0.9	2.1	8.7	45.4	155.7	329.3	507.0
Non-Hispanic white												
Both sexes												
2002 .....	56.4	0.9	0.2	0.2	0.4	0.6	2.0	9.0	46.7	181.6	416.3	673.1
2001 .....	55.9	*	*	0.2	0.3	0.5	1.9	8.7	48.6	185.9	408.6	676.9
2000 .....	55.7	*	*	0.2	0.4	0.6	1.8	8.6	48.2	186.5	413.2	680.8
1999 .....	56.6	*	0.2	0.2	0.4	0.6	1.6	8.5	51.8	193.6	422.4	675.6
Male												
2002 .....	55.1	*	*	0.2	0.4	0.6	2.0	9.3	48.3	200.4	506.7	935.4
2001 .....	54.6	*	*	0.2	0.3	0.5	1.9	9.1	50.5	206.1	500.9	951.5
2000 .....	55.1	*	*	0.2	0.4	0.5	1.7	8.9	50.8	208.8	513.6	1,008.6
1999 .....	57.3	*	*	0.3	0.5	0.5	1.5	8.5	55.6	224.9	534.9	1,033.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Chronic lower respiratory diseases (J40-J47)</b>												
<b>Non-Hispanic white</b>												
<b>Female</b>												
2002 .....	57.7	*	*	*	0.4	0.6	2.1	8.6	45.1	165.5	355.7	559.8
2001 .....	57.2	*	*	*	0.3	0.6	2.0	8.3	46.8	168.8	347.3	562.7
2000 .....	56.2	*	*	0.2	0.3	0.6	1.9	8.3	45.8	167.6	347.2	548.7
1999 .....	55.9	*	*	*	0.4	0.6	1.8	8.5	48.3	167.4	348.8	532.8
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	21.4	*	1.2	1.1	1.3	2.0	5.2	13.1	41.6	115.0	250.5	370.8
2001 .....	21.1	*	0.9	0.8	1.3	2.5	4.9	12.7	41.3	120.4	236.3	379.3
2000 .....	21.5	*	0.9	1.1	1.4	2.2	4.9	14.5	44.7	123.2	236.5	376.2
1999 .....	22.6	*	*	1.0	1.4	2.7	4.7	14.1	46.7	129.9	265.3	377.0
<b>Male</b>												
2002 .....	24.9	*	1.6	1.2	2.0	2.3	5.7	14.6	53.1	160.0	395.7	653.1
2001 .....	24.5	*	*	1.0	1.7	2.8	4.7	13.5	50.4	170.1	385.6	682.6
2000 .....	25.1	*	*	1.3	1.8	2.8	4.9	15.2	55.3	178.9	372.4	696.3
1999 .....	27.0	*	*	1.4	1.9	2.7	4.7	15.6	60.4	186.1	438.4	707.2
<b>Female</b>												
2002 .....	18.2	*	*	1.1	0.7	1.7	4.8	11.8	32.3	83.0	168.5	263.0
2001 .....	18.0	*	*	0.7	1.0	2.2	5.0	11.9	34.0	85.3	152.5	266.9
2000 .....	18.2	*	*	0.8	1.0	1.6	4.9	13.8	36.1	84.1	159.9	257.3
1999 .....	18.6	*	*	0.6	0.9	2.7	4.7	12.8	35.8	90.3	168.4	253.6
<b>Bronchitis, chronic and unspecified (J40-J42)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	0.3	0.6	*	*	*	*	*	0.1	0.2	0.7	2.2	8.1
2001 .....	0.3	0.5	*	*	*	*	0.0	0.1	0.2	0.7	2.1	9.1
2000 .....	0.4	*	*	*	*	*	0.0	0.1	0.3	0.9	2.7	11.8
1999 .....	0.4	0.5	*	*	*	*	*	0.1	0.4	0.9	2.8	10.7
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.5	8.8
2001 .....	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.3	9.2
2000 .....	0.4	*	*	*	*	*	*	*	0.3	1.0	3.6	12.9
1999 .....	0.4	*	*	*	*	*	*	*	0.4	1.1	3.3	11.9
<b>Female</b>												
2002 .....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.1	7.8
2001 .....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	1.9	9.1
2000 .....	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999 .....	0.5	*	*	*	*	*	*	*	0.4	0.8	2.5	10.3
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	11.1
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
Hispanic												
Male												
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.4	0.6	*	*	*	*	*	0.1	0.2	0.8	2.2	8.0
2001 .....	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.1	9.1
2000 .....	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.7	11.8
1999 .....	0.5	*	*	*	*	*	*	0.1	0.4	1.0	2.8	10.7
Male												
2002 .....	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.5	8.8
2001 .....	0.3	*	*	*	*	*	*	0.1	0.3	1.0	2.3	9.3
2000 .....	0.4	*	*	*	*	*	*	*	0.3	0.9	3.6	12.8
1999 .....	0.4	*	*	*	*	*	*	*	0.5	1.1	3.3	11.9
Female												
2002 .....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.0	7.6
2001 .....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.0	9.1
2000 .....	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999 .....	0.5	*	*	*	*	*	*	*	0.4	0.8	2.4	10.2
Non-Hispanic white												
Both sexes												
2002 .....	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.3	8.4
2001 .....	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.2	9.7
2000 .....	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.8	12.6
1999 .....	0.5	*	*	*	*	*	*	0.1	0.5	1.0	2.8	11.2
Male												
2002 .....	0.4	*	*	*	*	*	*	*	0.2	1.0	2.6	9.3
2001 .....	0.3	*	*	*	*	*	*	*	0.3	1.0	2.3	9.6
2000 .....	0.4	*	*	*	*	*	*	*	0.3	0.9	3.8	13.8
1999 .....	0.4	*	*	*	*	*	*	*	0.5	1.2	3.2	12.4
Female												
2002 .....	0.5	*	*	*	*	*	*	*	0.2	0.7	2.1	8.0
2001 .....	0.5	*	*	*	*	*	*	*	0.2	0.6	2.1	9.7
2000 .....	0.6	*	*	*	*	*	*	*	0.3	0.8	2.1	12.2
1999 .....	0.6	*	*	*	*	*	*	*	0.4	0.8	2.5	10.7
Non-Hispanic black												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	2.2	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	2.9	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
Non-Hispanic black												
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Emphysema (J43)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	5.4	*	*	*	*	*	0.2	1.1	6.4	24.0	48.0	59.3
2001 .....	5.7	*	*	*	*	*	0.2	1.1	7.2	26.2	50.4	62.1
2000 .....	5.9	*	*	*	*	*	0.2	1.3	7.4	27.8	52.7	63.3
1999 .....	6.4	*	*	*	*	*	0.2	1.3	8.5	30.7	56.9	62.1
Male												
2002 .....	5.7	*	*	*	*	*	0.3	1.4	7.2	28.1	63.5	90.0
2001 .....	6.0	*	*	*	*	*	0.2	1.4	7.9	30.8	65.3	101.2
2000 .....	6.3	*	*	*	*	*	0.2	1.5	8.1	33.0	69.1	104.2
1999 .....	6.9	*	*	*	*	*	0.2	1.5	9.5	37.8	76.4	105.4
Female												
2002 .....	5.0	*	*	*	*	*	0.1	0.9	5.6	20.6	37.6	46.0
2001 .....	5.4	*	*	*	*	*	0.1	0.9	6.6	22.5	40.6	45.8
2000 .....	5.6	*	*	*	*	*	0.2	1.0	6.8	23.6	41.9	46.7
1999 .....	5.8	*	*	*	*	*	0.2	1.1	7.5	24.9	44.2	44.7
Hispanic												
Both sexes												
2002 .....	0.8	*	*	*	*	*	*	*	1.5	6.7	18.8	36.6
2001 .....	0.8	*	*	*	*	*	*	*	1.4	6.6	20.4	39.4
2000 .....	0.8	*	*	*	*	*	*	*	1.5	6.0	22.3	31.2
1999 .....	0.9	*	*	*	*	*	*	*	1.3	7.1	23.5	44.2
Male												
2002 .....	0.8	*	*	*	*	*	*	*	*	9.2	26.7	53.5
2001 .....	0.9	*	*	*	*	*	*	*	*	9.2	34.0	60.9
2000 .....	0.9	*	*	*	*	*	*	*	*	9.1	32.5	58.4
1999 .....	1.0	*	*	*	*	*	*	*	*	11.2	36.0	77.6
Female												
2002 .....	0.7	*	*	*	*	*	*	*	*	4.7	13.5	28.2
2001 .....	0.6	*	*	*	*	*	*	*	*	4.6	11.2	28.9
2000 .....	0.6	*	*	*	*	*	*	*	*	3.6	15.5	*
1999 .....	0.7	*	*	*	*	*	*	*	*	4.0	15.3	28.1
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	6.1	*	*	*	*	*	0.2	1.2	6.8	25.1	49.3	60.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2001 .....	6.4	*	*	*	*	*	0.2	1.2	7.6	27.5	51.7	62.8
2000 .....	6.7	*	*	*	*	*	0.2	1.3	7.9	29.1	53.8	64.3
1999 .....	7.1	*	*	*	*	*	0.2	1.3	9.0	32.0	58.1	62.5
Male												
2002 .....	6.5	*	*	*	*	*	0.3	1.5	7.7	29.3	65.2	91.3
2001 .....	6.8	*	*	*	*	*	0.3	1.5	8.4	32.1	66.6	102.7
2000 .....	7.1	*	*	*	*	*	0.2	1.6	8.5	34.3	70.6	105.9
1999 .....	7.8	*	*	*	*	*	0.2	1.5	10.0	39.2	77.8	106.1
Female												
2002 .....	5.7	*	*	*	*	*	0.1	1.0	6.0	21.7	38.8	46.6
2001 .....	6.1	*	*	*	*	*	0.1	1.0	7.0	23.6	41.9	46.3
2000 .....	6.3	*	*	*	*	*	0.2	1.1	7.3	24.8	42.9	47.5
1999 .....	6.5	*	*	*	*	*	0.2	1.1	8.1	26.1	45.3	45.0
Non-Hispanic white												
Both sexes												
2002 .....	7.1	*	*	*	*	*	0.2	1.3	7.3	27.0	52.3	63.1
2001 .....	7.5	*	*	*	*	*	0.2	1.3	8.2	29.6	54.9	65.9
2000 .....	7.8	*	*	*	*	*	0.2	1.3	8.4	31.2	57.0	66.9
1999 .....	8.3	*	*	*	*	*	0.2	1.3	9.7	34.5	61.2	65.2
Male												
2002 .....	7.4	*	*	*	*	*	0.3	1.5	8.0	30.6	67.7	94.3
2001 .....	7.8	*	*	*	*	*	0.3	1.5	8.7	33.8	69.6	106.6
2000 .....	8.1	*	*	*	*	*	0.2	1.4	8.9	35.8	73.9	107.3
1999 .....	8.9	*	*	*	*	*	0.2	1.4	10.4	41.1	80.6	110.0
Female												
2002 .....	6.8	*	*	*	*	*	*	1.1	6.7	24.0	41.9	49.7
2001 .....	7.2	*	*	*	*	*	0.1	1.0	7.7	26.1	45.2	48.9
2000 .....	7.4	*	*	*	*	*	0.2	1.1	8.0	27.3	45.8	50.6
1999 .....	7.7	*	*	*	*	*	0.2	1.2	9.0	29.0	48.5	47.3
Non-Hispanic black												
Both sexes												
2002 .....	2.3	*	*	*	*	*	0.5	1.2	4.6	16.2	28.2	32.1
2001 .....	2.2	*	*	*	*	*	*	1.4	5.6	15.6	23.6	35.9
2000 .....	2.5	*	*	*	*	*	*	2.0	6.1	17.1	27.0	37.5
1999 .....	2.7	*	*	*	*	*	0.4	1.9	6.3	17.3	31.8	37.8
Male												
2002 .....	3.1	*	*	*	*	*	*	1.8	6.7	25.3	49.5	65.6
2001 .....	2.8	*	*	*	*	*	*	1.6	7.9	23.0	40.2	69.3
2000 .....	3.3	*	*	*	*	*	*	3.0	8.2	27.4	41.3	89.2
1999 .....	3.6	*	*	*	*	*	*	3.0	9.1	29.3	55.7	73.7
Female												
2002 .....	1.6	*	*	*	*	*	*	*	2.8	9.8	16.1	19.2
2001 .....	1.7	*	*	*	*	*	*	1.1	3.7	10.4	14.3	23.5
2000 .....	1.8	*	*	*	*	*	*	1.3	4.5	9.8	18.9	18.3
1999 .....	1.8	*	*	*	*	*	*	1.0	4.0	8.9	18.4	24.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Asthma (J45-J46)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	1.5	*	0.3	0.3	0.4	0.6	1.1	1.5	2.0	3.2	6.4	14.7
2001 .....	1.5	*	0.2	0.2	0.4	0.6	1.0	1.5	2.2	3.5	6.4	16.1
2000 .....	1.6	*	0.2	0.3	0.4	0.6	1.0	1.6	2.2	4.0	6.9	16.7
1999 .....	1.7	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.2	7.6	17.3
<b>Male</b>												
2002 .....	1.1	*	0.4	0.3	0.5	0.6	0.9	1.2	1.6	2.4	5.0	10.9
2001 .....	1.1	*	0.3	0.3	0.4	0.6	0.9	1.1	1.5	2.7	4.5	12.1
2000 .....	1.2	*	0.3	0.4	0.5	0.7	0.9	1.3	1.7	3.0	5.0	12.9
1999 .....	1.2	*	*	0.4	0.5	0.6	0.8	1.1	1.8	3.2	5.7	12.7
<b>Female</b>												
2002 .....	1.8	*	*	0.3	0.3	0.5	1.2	1.8	2.4	3.9	7.3	16.3
2001 .....	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.8	4.1	7.6	17.8
2000 .....	2.0	*	*	0.3	0.3	0.5	1.2	2.0	2.7	4.8	8.1	18.2
1999 .....	2.1	*	*	0.2	0.4	0.7	1.2	2.2	3.0	5.0	8.8	19.2
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	0.7	*	*	*	*	0.3	0.7	1.2	1.9	3.8	7.0	12.2
2001 .....	0.7	*	*	*	*	*	0.6	1.4	2.4	3.8	7.9	13.8
2000 .....	0.8	*	*	*	*	*	0.6	1.3	2.8	5.6	7.9	16.6
1999 .....	0.9	*	*	*	*	0.3	0.9	1.6	2.4	5.8	10.4	17.5
<b>Male</b>												
2002 .....	0.6	*	*	*	*	*	*	1.2	*	*	*	*
2001 .....	0.5	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.7	*	*	*	*	*	*	1.3	*	5.3	*	*
1999 .....	0.7	*	*	*	*	*	0.9	1.4	*	*	*	*
<b>Female</b>												
2002 .....	0.9	*	*	*	*	*	1.0	1.2	2.2	5.3	7.6	*
2001 .....	1.0	*	*	*	*	*	*	2.0	2.9	3.9	9.4	18.6
2000 .....	1.0	*	*	*	*	*	*	1.3	3.2	5.8	9.2	*
1999 .....	1.2	*	*	*	*	*	1.0	1.9	3.1	7.1	13.2	*
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	1.6	*	0.3	0.3	0.5	0.6	1.1	1.5	2.0	3.1	6.3	14.7
2001 .....	1.6	*	0.2	0.3	0.4	0.7	1.1	1.5	2.2	3.4	6.3	16.1
2000 .....	1.7	*	0.2	0.3	0.5	0.7	1.1	1.6	2.2	3.9	6.8	16.6
1999 .....	1.8	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.1	7.4	17.3
<b>Male</b>												
2002 .....	1.2	*	0.4	0.3	0.6	0.7	0.9	1.2	1.6	2.4	4.9	11.1
2001 .....	1.1	*	*	0.3	0.4	0.6	0.9	1.2	1.5	2.6	4.4	12.2
2000 .....	1.2	*	*	0.4	0.6	0.8	0.9	1.3	1.6	2.9	4.9	12.7
1999 .....	1.3	*	*	0.4	0.6	0.7	0.8	1.1	1.8	3.1	5.6	12.6
<b>Female</b>												
2002 .....	2.0	*	*	0.3	0.3	0.6	1.2	1.9	2.4	3.8	7.2	16.3
2001 .....	2.1	*	*	0.2	0.4	0.7	1.2	1.9	2.8	4.1	7.5	17.7
2000 .....	2.1	*	*	0.3	0.4	0.6	1.2	2.0	2.6	4.7	8.0	18.2
1999 .....	2.2	*	*	0.2	0.4	0.7	1.2	2.2	3.0	4.9	8.6	19.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Asthma (J45-J46)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	1.4	*	*	0.1	0.3	0.5	0.7	1.1	1.5	2.6	6.1	14.7
2001 .....	1.4	*	*	0.1	0.2	0.4	0.7	1.1	1.5	2.9	5.8	15.7
2000 .....	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.5	3.2	6.4	15.8
1999 .....	1.5	*	*	0.2	0.4	0.4	0.7	1.2	1.9	3.4	7.1	16.5
<b>Male</b>												
2002 .....	0.9	*	*	0.2	0.3	0.5	0.5	0.8	1.1	1.8	4.5	10.8
2001 .....	0.9	*	*	0.2	0.2	0.3	0.6	0.8	1.0	2.1	3.9	11.4
2000 .....	1.0	*	*	0.2	0.3	0.5	0.6	0.9	1.0	2.3	4.6	11.1
1999 .....	1.0	*	*	0.2	0.4	0.4	0.5	0.7	1.4	2.4	5.2	10.6
<b>Female</b>												
2002 .....	1.8	*	*	*	0.3	0.5	0.9	1.3	1.8	3.2	7.1	16.4
2001 .....	1.9	*	*	*	0.2	0.4	0.8	1.4	2.1	3.6	7.1	17.5
2000 .....	1.9	*	*	0.2	0.3	0.4	0.9	1.3	2.0	4.0	7.6	17.7
1999 .....	2.1	*	*	*	0.3	0.5	0.9	1.6	2.3	4.2	8.3	18.8
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	3.0	*	0.9	1.1	1.3	1.6	3.5	4.9	6.6	8.0	7.9	11.8
2001 .....	3.1	*	*	0.8	1.2	2.2	3.3	4.7	6.9	7.7	10.7	16.5
2000 .....	3.3	*	*	1.1	1.3	2.0	3.2	5.5	6.9	9.4	10.4	22.3
1999 .....	3.3	*	*	1.0	1.3	2.4	3.0	5.4	7.0	9.8	10.6	20.7
<b>Male</b>												
2002 .....	2.8	*	*	1.1	1.9	2.0	3.7	4.0	5.9	7.5	6.7	*
2001 .....	2.7	*	*	0.9	1.4	2.5	3.0	4.1	4.7	6.8	8.7	*
2000 .....	2.8	*	*	1.3	1.6	2.5	3.0	3.7	6.1	7.9	6.5	24.7
1999 .....	2.9	*	*	1.4	1.7	2.7	2.8	4.0	5.5	9.2	8.4	*
<b>Female</b>												
2002 .....	3.1	*	*	1.0	*	1.3	3.4	5.7	7.2	8.4	8.6	12.5
2001 .....	3.4	*	*	0.7	0.9	2.0	3.6	5.1	8.6	8.4	11.8	17.1
2000 .....	3.7	*	*	0.8	1.0	1.5	3.4	7.0	7.6	10.4	12.7	21.4
1999 .....	3.6	*	*	0.6	0.9	2.2	3.2	6.5	8.1	10.2	11.7	21.3
<b>Other chronic lower respiratory diseases (J44,J47)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	36.1	*	*	*	*	0.1	0.9	5.9	33.8	135.1	330.1	555.5
2001 .....	35.7	*	*	*	0.1	0.1	0.9	5.7	34.5	137.5	321.0	557.4
2000 .....	35.4	*	*	*	*	0.1	0.8	5.6	34.3	136.7	323.8	556.7
1999 .....	36.0	*	*	*	*	0.1	0.7	5.5	36.2	141.4	330.6	555.8
<b>Male</b>												
2002 .....	35.7	*	*	*	*	0.1	1.0	6.4	36.2	153.4	409.8	785.1
2001 .....	35.3	*	*	*	*	0.1	0.9	6.2	37.2	157.0	403.0	794.5
2000 .....	35.6	*	*	*	*	*	0.8	6.1	37.7	158.3	410.9	838.0
1999 .....	37.1	*	*	*	*	*	0.8	5.9	40.5	168.6	427.9	866.7
<b>Female</b>												
2002 .....	36.5	*	*	*	*	0.1	0.9	5.4	31.5	119.8	277.2	455.8
2001 .....	36.0	*	*	*	*	0.1	1.0	5.2	32.0	121.3	267.1	458.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
All origins <sup>3</sup>												
Female												
2002 .....	35.2	*	*	*	*	*	0.8	5.2	31.2	119.0	267.1	442.2
1999 .....	35.0	*	*	*	*	0.2	0.7	5.0	32.2	119.1	267.4	430.5
Hispanic												
Both sexes												
2002 .....	6.3	*	*	*	*	*	0.4	2.1	10.4	43.1	147.8	397.4
2001 .....	6.0	*	*	*	*	*	*	1.9	10.0	45.2	139.6	418.9
2000 .....	5.9	*	*	*	*	*	0.4	1.7	9.9	40.7	149.1	422.7
1999 .....	6.5	*	*	*	*	*	0.4	1.9	10.8	49.3	158.1	462.3
Male												
2002 .....	6.6	*	*	*	*	*	*	2.2	14.0	57.9	208.9	535.4
2001 .....	6.3	*	*	*	*	*	*	2.1	12.2	62.5	181.5	605.7
2000 .....	6.2	*	*	*	*	*	*	1.6	13.6	56.6	209.0	584.5
1999 .....	7.0	*	*	*	*	*	*	1.6	13.9	65.9	228.9	735.4
Female												
2002 .....	5.9	*	*	*	*	*	*	1.9	7.1	31.3	106.0	329.1
2001 .....	5.8	*	*	*	*	*	*	1.8	8.1	31.5	111.2	327.7
2000 .....	5.6	*	*	*	*	*	*	1.7	6.6	28.2	109.0	343.3
1999 .....	5.9	*	*	*	*	*	*	2.1	8.1	36.3	111.0	330.7
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	40.6	*	*	*	*	0.1	1.0	6.3	35.6	141.0	338.1	560.8
2001 .....	40.0	*	*	*	0.1	0.1	1.0	6.0	36.2	143.2	328.6	561.3
2000 .....	39.5	*	*	*	*	0.1	0.9	6.0	36.0	142.3	330.5	560.5
1999 .....	40.0	*	*	*	*	0.1	0.8	5.8	37.9	146.4	336.8	557.6
Male												
2002 .....	40.4	*	*	*	*	0.1	1.1	6.9	37.8	159.2	418.5	794.9
2001 .....	39.8	*	*	*	*	0.1	1.0	6.6	38.9	162.6	412.4	800.6
2000 .....	40.0	*	*	*	*	*	0.9	6.5	39.3	163.8	418.4	846.6
1999 .....	41.4	*	*	*	*	*	0.8	6.3	42.2	173.9	435.1	869.6
Female												
2002 .....	40.8	*	*	*	*	0.1	1.0	5.8	33.4	125.8	284.9	459.8
2001 .....	40.1	*	*	*	*	*	1.0	5.5	33.8	127.1	273.6	461.7
2000 .....	39.1	*	*	*	*	*	0.9	5.5	32.9	124.5	273.2	444.8
1999 .....	38.7	*	*	*	*	0.2	0.7	5.2	33.9	123.9	272.9	432.7
Non-Hispanic white												
Both sexes												
2002 .....	47.5	*	*	*	*	0.1	1.0	6.5	37.7	151.2	355.7	586.9
2001 .....	46.6	*	*	*	*	0.1	1.0	6.2	38.6	152.6	345.7	585.7
2000 .....	46.0	*	*	*	*	*	0.8	6.1	37.9	151.2	347.1	585.5
1999 .....	46.3	*	*	*	*	0.1	0.7	5.9	39.9	154.8	351.4	582.8
Male												
2002 .....	46.4	*	*	*	*	*	1.1	6.9	39.0	167.0	431.9	821.0
2001 .....	45.6	*	*	*	*	*	1.0	6.7	40.5	169.3	425.2	823.9
2000 .....	45.7	*	*	*	*	*	0.9	6.4	40.5	169.8	431.4	876.5
1999 .....	47.1	*	*	*	*	*	0.8	6.2	43.4	180.3	445.9	900.5

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
Non-Hispanic white												
Female												
2002 .....	48.6	*	*	*	*	*	1.0	6.2	36.4	137.7	304.6	485.7
2001 .....	47.6	*	*	*	*	*	1.0	5.7	36.8	138.5	292.9	486.6
2000 .....	46.3	*	*	*	*	*	0.8	5.8	35.5	135.5	291.6	468.2
1999 .....	45.5	*	*	*	*	*	0.7	5.5	36.6	133.4	289.5	456.1
Non-Hispanic black												
Both sexes												
2002 .....	15.9	*	*	*	*	*	1.2	6.8	30.2	90.3	213.1	322.7
2001 .....	15.6	*	*	*	*	*	1.3	6.5	28.7	96.1	200.8	323.4
2000 .....	15.4	*	*	*	*	*	1.3	6.8	31.4	95.8	196.8	312.6
1999 .....	16.4	*	*	*	*	*	1.2	6.6	33.2	101.9	220.1	314.0
Male												
2002 .....	18.7	*	*	*	*	*	1.3	8.5	40.3	126.6	337.9	572.1
2001 .....	18.8	*	*	*	*	*	1.4	7.5	37.6	139.1	334.9	592.5
2000 .....	18.7	*	*	*	*	*	1.3	8.5	40.7	142.7	322.4	578.9
1999 .....	20.3	*	*	*	*	*	1.5	8.3	45.4	146.5	369.3	610.9
Female												
2002 .....	13.3	*	*	*	*	*	1.1	5.3	22.1	64.6	142.4	227.4
2001 .....	12.7	*	*	*	*	*	1.3	5.7	21.5	65.8	125.5	223.7
2000 .....	12.4	*	*	*	*	*	1.3	5.3	24.0	62.8	126.1	213.7
1999 .....	12.9	*	*	*	*	*	1.0	5.0	23.3	70.4	136.6	203.0
Pneumoconioses and chemical effects (J60-J66,J68)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.4	*	*	*	*	*	*	*	0.3	1.2	3.8	6.8
2001 .....	0.4	*	*	*	*	*	*	*	0.3	1.5	4.1	6.2
2000 .....	0.4	*	*	*	*	*	*	*	0.2	1.5	3.9	7.5
1999 .....	0.4	*	*	*	*	*	*	*	0.2	1.5	4.1	6.1
Male												
2002 .....	0.7	*	*	*	*	*	*	*	0.5	2.6	9.0	21.6
2001 .....	0.8	*	*	*	*	*	*	*	0.5	3.1	9.9	20.2
2000 .....	0.8	*	*	*	*	*	*	*	0.5	3.2	9.6	24.9
1999 .....	0.8	*	*	*	*	*	*	*	0.5	3.2	10.1	20.4
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
Hispanic												
Male												
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.4	*	*	*	*	*	*	*	0.3	1.2	3.9	6.9
2001 .....	0.4	*	*	*	*	*	*	*	0.3	1.5	4.2	6.2
2000 .....	0.5	*	*	*	*	*	*	*	0.2	1.6	4.0	7.6
1999 .....	0.4	*	*	*	*	*	*	*	0.3	1.5	4.3	6.2
Male												
2002 .....	0.8	*	*	*	*	*	*	*	0.5	2.6	9.3	21.9
2001 .....	0.9	*	*	*	*	*	*	*	0.5	3.3	10.2	20.5
2000 .....	0.9	*	*	*	*	*	*	*	0.5	3.3	9.7	25.7
1999 .....	0.9	*	*	*	*	*	*	*	0.5	3.4	10.5	21.1
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	0.5	*	*	*	*	*	*	*	0.3	1.4	4.1	7.3
2001 .....	0.5	*	*	*	*	*	*	*	0.3	1.6	4.5	6.6
2000 .....	0.5	*	*	*	*	*	*	*	0.3	1.7	4.1	8.1
1999 .....	0.5	*	*	*	*	*	*	*	0.2	1.6	4.6	6.4
Male												
2002 .....	1.0	*	*	*	*	*	*	*	0.6	2.8	9.8	23.2
2001 .....	1.0	*	*	*	*	*	*	*	0.5	3.5	10.8	21.6
2000 .....	1.1	*	*	*	*	*	*	*	0.5	3.6	10.1	27.2
1999 .....	1.0	*	*	*	*	*	*	*	0.5	3.6	11.1	21.8
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	2.3	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	2.2	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	2.6	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Pneumoconioses and chemical effects (J60-J66,J68)</b>												
<b>Non-Hispanic black</b>												
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	5.9	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	6.0	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	6.5	*
1999 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Pneumonitis due to solids and liquids (J69)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.3	186.0
2001 .....	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000 .....	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999 .....	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6
<b>Male</b>												
2002 .....	6.3	*	*	*	0.1	0.2	0.5	1.1	3.1	13.0	63.8	268.5
2001 .....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.1	270.3
2000 .....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999 .....	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.1	59.7	258.4
<b>Female</b>												
2002 .....	5.9	*	*	*	*	0.2	0.3	0.6	2.0	7.1	34.7	150.2
2001 .....	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.6
2000 .....	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999 .....	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.1	142.2
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	1.4	*	*	*	*	*	*	*	1.8	5.3	29.0	131.4
2001 .....	1.5	*	*	*	*	*	*	*	1.8	6.1	27.4	161.3
2000 .....	1.4	*	*	*	*	*	*	0.6	1.9	5.9	27.9	137.4
1999 .....	1.2	*	*	*	*	*	*	*	1.6	5.5	24.6	130.5
<b>Male</b>												
2002 .....	1.4	*	*	*	*	*	*	*	2.6	6.1	43.1	167.3
2001 .....	1.5	*	*	*	*	*	*	*	2.5	8.4	36.2	220.9
2000 .....	1.4	*	*	*	*	*	*	*	2.5	7.4	42.8	169.3
1999 .....	1.1	*	*	*	*	*	*	*	*	6.8	33.9	159.6
<b>Female</b>												
2002 .....	1.4	*	*	*	*	*	*	*	*	4.6	19.4	113.6
2001 .....	1.5	*	*	*	*	*	*	*	*	4.3	21.5	132.2
2000 .....	1.4	*	*	*	*	*	*	*	*	4.8	17.8	121.7
1999 .....	1.3	*	*	*	*	*	*	*	*	4.5	18.4	116.5
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	6.8	*	*	*	0.1	0.2	0.5	0.9	2.6	10.1	47.1	187.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Pneumonitis due to solids and liquids (J69)</b>												
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2001 .....	6.7	*	*	*	0.1	0.2	0.4	1.0	2.7	10.2	46.5	190.0
2000 .....	6.5	*	*	*	0.1	0.2	0.4	1.0	2.5	10.6	45.1	189.0
1999 .....	6.0	*	*	*	0.1	0.2	0.4	0.8	2.5	9.7	41.7	176.8
<b>Male</b>												
2002 .....	7.0	*	*	*	0.1	0.2	0.6	1.2	3.1	13.4	64.7	272.1
2001 .....	6.8	*	*	*	0.1	0.2	0.6	1.3	3.2	13.6	64.2	271.6
2000 .....	6.8	*	*	*	0.1	0.2	0.5	1.2	3.1	14.8	62.8	279.7
1999 .....	6.3	*	*	*	0.1	0.3	0.5	1.0	3.3	13.3	60.6	261.6
<b>Female</b>												
2002 .....	6.6	*	*	*	*	0.2	0.3	0.6	2.1	7.3	35.4	151.4
2001 .....	6.6	*	*	*	*	0.2	0.3	0.8	2.2	7.4	34.9	156.0
2000 .....	6.3	*	*	*	*	0.1	0.3	0.8	2.0	7.1	33.6	152.3
1999 .....	5.8	*	*	*	*	0.2	0.3	0.7	1.8	6.8	29.4	142.9
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	7.7	*	*	*	0.1	0.2	0.4	0.9	2.5	9.7	47.3	191.3
2001 .....	7.5	*	*	*	0.1	0.2	0.4	0.9	2.6	9.9	46.6	191.2
2000 .....	7.3	*	*	*	0.1	0.2	0.4	0.9	2.3	10.0	45.0	192.5
1999 .....	6.7	*	*	*	0.1	0.2	0.4	0.7	2.4	9.2	41.2	179.3
<b>Male</b>												
2002 .....	7.9	*	*	*	*	0.3	0.6	1.1	3.0	12.9	64.9	278.4
2001 .....	7.6	*	*	*	*	0.3	0.6	1.2	3.1	13.2	64.4	276.9
2000 .....	7.5	*	*	*	*	0.2	0.5	1.1	2.9	13.7	62.7	288.1
1999 .....	6.9	*	*	*	*	0.2	0.4	0.9	3.0	12.4	59.6	267.4
<b>Female</b>												
2002 .....	7.4	*	*	*	*	0.2	0.3	0.6	2.0	7.1	35.6	153.7
2001 .....	7.4	*	*	*	*	0.2	0.3	0.7	2.1	7.0	34.7	155.5
2000 .....	7.1	*	*	*	*	*	0.3	0.7	1.8	7.0	33.4	153.9
1999 .....	6.5	*	*	*	*	0.2	0.3	0.6	1.7	6.5	29.3	144.2
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	4.2	*	*	*	*	*	0.7	1.4	3.8	15.2	52.8	167.3
2001 .....	4.4	*	*	*	*	*	0.6	1.6	4.1	15.4	53.6	187.3
2000 .....	4.3	*	*	*	*	*	0.7	1.9	4.3	17.9	52.0	160.6
1999 .....	4.3	*	*	*	*	*	0.7	1.5	4.5	16.4	52.6	168.5
<b>Male</b>												
2002 .....	4.4	*	*	*	*	*	0.9	2.0	4.9	22.3	74.1	237.4
2001 .....	4.3	*	*	*	*	*	0.8	1.9	5.5	20.8	75.6	243.7
2000 .....	4.5	*	*	*	*	*	0.8	2.7	5.2	28.4	75.0	210.2
1999 .....	4.7	*	*	*	*	*	0.9	1.8	6.6	25.3	81.6	229.4
<b>Female</b>												
2002 .....	4.1	*	*	*	*	*	*	0.9	3.0	10.2	40.8	140.5
2001 .....	4.6	*	*	*	*	*	*	1.4	2.9	11.6	41.3	166.4
2000 .....	4.1	*	*	*	*	*	*	1.2	3.6	10.4	39.1	142.2
1999 .....	4.0	*	*	*	*	*	*	1.2	2.9	10.2	36.4	145.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39, J67,J70-J98)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	8.7	6.9	0.5	0.2	0.3	0.5	1.2	3.2	9.2	28.7	68.0	133.5
2001 .....	9.2	7.3	0.6	0.2	0.4	0.5	1.3	3.5	9.6	30.1	73.2	142.7
2000 .....	9.1	7.2	0.6	0.2	0.3	0.6	1.4	3.5	9.9	29.6	72.0	141.7
1999 .....	9.0	6.6	0.6	0.2	0.3	0.6	1.3	3.3	9.5	29.4	70.7	147.6
Male												
2002 .....	8.5	7.9	0.7	0.2	0.3	0.4	1.3	3.3	10.5	35.3	83.9	156.6
2001 .....	8.9	9.0	0.6	0.2	0.4	0.5	1.4	3.8	10.9	36.5	89.3	169.1
2000 .....	9.0	7.9	0.6	0.2	0.4	0.6	1.5	3.8	11.4	35.7	90.5	179.7
1999 .....	8.9	7.5	0.7	0.2	0.3	0.6	1.4	3.5	10.6	36.2	89.9	182.3
Female												
2002 .....	8.8	5.8	0.4	0.2	0.3	0.5	1.1	3.1	8.0	23.1	57.5	123.4
2001 .....	9.4	5.5	0.6	0.2	0.3	0.6	1.2	3.1	8.4	24.8	62.6	131.6
2000 .....	9.2	6.4	0.6	0.2	0.3	0.6	1.2	3.2	8.6	24.7	59.9	126.2
1999 .....	9.1	5.8	0.5	0.2	0.2	0.7	1.1	3.1	8.5	23.7	58.3	133.6
Hispanic												
Both sexes												
2002 .....	3.3	5.2	*	*	*	0.4	0.7	2.7	8.3	26.1	58.0	112.0
2001 .....	3.4	5.0	*	*	*	*	0.7	3.1	8.9	25.3	65.5	134.4
2000 .....	3.5	5.6	*	*	*	0.5	1.1	2.6	9.8	25.4	64.2	135.4
1999 .....	3.4	5.0	*	*	*	0.4	0.9	3.2	9.6	26.0	64.0	134.0
Male												
2002 .....	3.2	4.9	*	*	*	*	0.8	2.6	9.8	34.6	64.0	108.8
2001 .....	3.4	5.0	*	*	*	*	0.9	3.4	10.6	30.2	78.3	167.6
2000 .....	3.3	5.8	*	*	*	*	1.3	3.0	11.9	29.3	81.1	133.0
1999 .....	3.3	5.2	*	*	*	*	1.1	3.1	11.6	30.5	82.4	161.8
Female												
2002 .....	3.4	5.4	*	*	*	*	*	2.8	6.9	19.3	54.0	113.6
2001 .....	3.5	5.0	*	*	*	*	*	2.8	7.5	21.4	56.8	118.2
2000 .....	3.6	5.3	*	*	*	*	0.9	2.2	7.8	22.2	52.8	136.5
1999 .....	3.5	*	*	*	*	*	*	3.3	7.8	22.4	51.7	120.6
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	9.5	7.3	0.5	0.2	0.3	0.5	1.2	3.2	9.2	28.8	68.4	134.1
2001 .....	10.0	7.8	0.6	0.2	0.4	0.6	1.3	3.5	9.7	30.3	73.3	142.7
2000 .....	9.8	7.5	0.6	0.2	0.4	0.7	1.4	3.6	9.9	29.8	72.1	141.6
1999 .....	9.7	6.9	0.6	0.2	0.3	0.7	1.3	3.3	9.4	29.5	70.8	147.7
Male												
2002 .....	9.4	8.7	0.7	0.2	0.3	0.4	1.3	3.4	10.5	35.3	84.7	158.5
2001 .....	9.8	10.0	0.6	0.2	0.4	0.6	1.4	3.9	10.9	36.7	89.5	169.1
2000 .....	9.8	8.4	0.6	0.2	0.4	0.6	1.5	3.9	11.3	36.0	90.7	181.3
1999 .....	9.6	7.9	0.8	0.2	0.4	0.7	1.4	3.6	10.5	36.4	89.9	182.7
Female												
2002 .....	9.6	5.9	0.4	0.2	0.3	0.5	1.2	3.1	8.0	23.3	57.5	123.6
2001 .....	10.2	5.5	0.6	0.2	0.3	0.6	1.3	3.1	8.5	25.0	62.7	131.8
2000 .....	9.9	6.6	0.5	0.2	0.3	0.7	1.3	3.3	8.6	24.7	60.1	125.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39, J67,J70-J98)												
Non-Hispanic <sup>4</sup>												
Female												
1999 .....	9.8	5.9	0.5	0.2	0.2	0.7	1.2	3.1	8.5	23.8	58.4	133.7
Non-Hispanic white												
Both sexes												
2002 .....	10.5	4.7	0.4	0.2	0.3	0.4	1.1	2.9	8.8	29.3	70.3	138.6
2001 .....	11.0	4.8	0.5	0.2	0.3	0.5	1.1	3.0	9.3	30.9	75.6	145.9
2000 .....	10.8	5.4	0.5	0.2	0.3	0.6	1.2	3.2	9.6	30.2	73.8	145.7
1999 .....	10.6	4.9	0.6	0.2	0.3	0.6	1.1	3.0	9.1	29.7	72.3	150.7
Male												
2002 .....	10.3	5.6	0.5	*	0.3	0.4	1.1	3.0	10.1	36.0	87.0	164.1
2001 .....	10.8	6.2	0.4	0.2	0.4	0.4	1.3	3.4	10.5	37.7	92.4	173.4
2000 .....	10.7	6.6	0.5	0.2	0.3	0.6	1.3	3.5	11.0	36.4	92.6	185.9
1999 .....	10.5	5.6	0.6	0.2	0.4	0.7	1.2	3.2	10.0	37.0	92.0	186.1
Female												
2002 .....	10.7	3.7	*	0.2	0.2	0.5	1.0	2.7	7.7	23.6	59.1	127.6
2001 .....	11.3	3.4	0.5	0.2	0.3	0.6	0.9	2.6	8.2	25.1	64.4	134.5
2000 .....	11.0	4.1	0.5	*	0.3	0.6	1.1	3.0	8.4	25.0	61.5	129.5
1999 .....	10.7	4.0	0.5	0.2	0.2	0.5	0.9	2.7	8.2	23.6	59.4	136.6
Non-Hispanic black												
Both sexes												
2002 .....	6.2	17.3	1.2	0.4	0.7	0.9	2.4	6.2	13.8	27.1	52.3	90.7
2001 .....	6.9	19.0	1.1	*	0.7	1.3	2.9	7.5	13.9	29.6	56.9	110.8
2000 .....	6.7	17.7	0.8	*	0.7	1.1	2.7	6.9	13.7	30.6	58.6	99.8
1999 .....	6.7	15.8	1.1	*	0.4	1.4	2.9	6.6	13.6	31.3	58.7	112.8
Male												
2002 .....	6.3	20.3	*	*	*	0.9	2.8	6.5	16.4	32.5	67.1	102.8
2001 .....	6.6	24.3	*	*	*	1.4	2.6	7.8	15.7	32.9	69.0	131.7
2000 .....	6.8	17.1	*	*	0.8	1.2	3.2	8.0	15.6	36.2	74.0	135.0
1999 .....	6.6	18.0	*	*	*	1.1	2.9	7.3	15.9	35.6	73.5	146.2
Female												
2002 .....	6.2	14.2	*	*	0.7	0.9	2.0	5.9	11.7	23.3	43.9	86.1
2001 .....	7.2	13.4	*	*	*	1.2	3.2	7.2	12.4	27.2	50.0	103.1
2000 .....	6.6	18.3	*	*	*	1.0	2.3	6.0	12.1	26.6	49.8	86.8
1999 .....	6.9	13.6	*	*	*	1.7	2.9	5.9	11.9	28.2	50.4	100.4
Peptic ulcer (K25-K28)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.4	*	*	*	*	0.1	0.3	0.7	1.5	3.6	10.1	28.3
2001 .....	1.6	*	*	*	*	0.1	0.3	0.7	1.7	3.9	11.4	33.4
2000 .....	1.6	*	*	*	*	*	0.3	0.8	1.8	4.1	11.4	35.6
1999 .....	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	12.3	34.6
Male												
2002 .....	1.4	*	*	*	*	*	0.3	0.9	1.9	4.6	12.2	31.6
2001 .....	1.5	*	*	*	*	0.1	0.4	1.0	2.2	4.8	13.0	34.9
2000 .....	1.6	*	*	*	*	*	0.4	1.2	2.4	5.3	13.7	40.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
All origins <sup>3</sup>												
Male												
1999 .....	1.6	*	*	*	*	*	0.3	0.9	2.3	6.1	14.8	37.5
Female												
2002 .....	1.5	*	*	*	*	*	0.2	0.5	1.2	2.9	8.7	26.9
2001 .....	1.7	*	*	*	*	*	0.1	0.5	1.1	3.2	10.4	32.7
2000 .....	1.6	*	*	*	*	*	0.1	0.5	1.3	3.0	10.0	33.5
1999 .....	1.7	*	*	*	*	*	0.2	0.5	1.3	3.3	10.6	33.5
Hispanic												
Both sexes												
2002 .....	0.5	*	*	*	*	*	*	0.5	1.6	3.3	8.8	20.5
2001 .....	0.4	*	*	*	*	*	*	*	1.2	2.9	9.7	23.8
2000 .....	0.5	*	*	*	*	*	*	*	1.6	3.3	9.9	30.5
1999 .....	0.5	*	*	*	*	*	*	*	1.5	3.6	9.2	26.7
Male												
2002 .....	0.6	*	*	*	*	*	*	*	2.5	5.4	11.1	*
2001 .....	0.5	*	*	*	*	*	*	*	*	*	16.5	*
2000 .....	0.5	*	*	*	*	*	*	*	2.9	4.4	*	*
1999 .....	0.6	*	*	*	*	*	*	*	*	4.8	14.1	*
Female												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	7.3	17.4
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	25.1
2000 .....	0.5	*	*	*	*	*	*	*	*	*	10.9	29.7
1999 .....	0.4	*	*	*	*	*	*	*	*	*	*	22.9
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.6	*	*	*	*	0.1	0.3	0.7	1.5	3.7	10.2	28.6
2001 .....	1.7	*	*	*	*	0.1	0.3	0.7	1.7	4.0	11.5	33.6
2000 .....	1.8	*	*	*	*	*	0.3	0.8	1.8	4.1	11.5	35.7
1999 .....	1.8	*	*	*	*	*	0.3	0.7	1.8	4.6	12.4	34.8
Male												
2002 .....	1.5	*	*	*	*	*	0.4	0.9	1.9	4.5	12.3	31.7
2001 .....	1.6	*	*	*	*	0.1	0.4	1.0	2.2	4.8	12.8	35.5
2000 .....	1.8	*	*	*	*	*	0.4	1.2	2.4	5.4	13.9	41.0
1999 .....	1.7	*	*	*	*	*	0.4	1.0	2.3	6.1	14.8	37.7
Female												
2002 .....	1.6	*	*	*	*	*	0.2	0.5	1.2	3.0	8.8	27.3
2001 .....	1.8	*	*	*	*	*	0.2	0.5	1.2	3.3	10.6	32.8
2000 .....	1.8	*	*	*	*	*	0.1	0.5	1.3	3.1	9.9	33.5
1999 .....	1.9	*	*	*	*	*	0.2	0.5	1.3	3.3	10.8	33.7
Non-Hispanic white												
Both sexes												
2002 .....	1.7	*	*	*	*	*	0.2	0.6	1.4	3.6	10.4	29.4
2001 .....	1.9	*	*	*	*	*	0.2	0.7	1.5	3.9	11.6	34.7
2000 .....	1.9	*	*	*	*	*	0.2	0.7	1.8	4.0	11.6	36.3
1999 .....	2.0	*	*	*	*	*	0.2	0.6	1.7	4.5	12.4	35.6
Male												
2002 .....	1.6	*	*	*	*	*	0.3	0.8	1.7	4.4	12.4	32.6

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Peptic ulcer (K25-K28)</b>												
<b>Non-Hispanic white</b>												
<b>Male</b>												
2001 .....	1.7	*	*	*	*	*	0.3	0.8	2.0	4.5	12.7	36.3
2000 .....	1.9	*	*	*	*	*	0.3	1.1	2.3	5.2	13.7	41.0
1999 .....	1.8	*	*	*	*	*	0.3	0.8	2.1	5.9	14.9	38.6
<b>Female</b>												
2002 .....	1.8	*	*	*	*	*	0.2	0.5	1.1	3.0	9.0	27.9
2001 .....	2.1	*	*	*	*	*	0.1	0.5	1.1	3.4	10.9	34.0
2000 .....	2.0	*	*	*	*	*	0.1	0.4	1.3	3.1	10.2	34.4
1999 .....	2.1	*	*	*	*	*	0.2	0.4	1.3	3.2	10.8	34.4
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	1.0	*	*	*	*	*	0.6	1.2	2.6	4.2	7.9	18.8
2001 .....	1.1	*	*	*	*	*	0.6	1.4	2.8	4.5	9.0	20.9
2000 .....	1.2	*	*	*	*	*	0.5	1.7	2.6	4.9	10.1	26.7
1999 .....	1.3	*	*	*	*	*	0.8	1.5	2.8	5.7	11.0	25.5
<b>Male</b>												
2002 .....	1.2	*	*	*	*	*	0.8	2.0	3.9	6.0	11.1	*
2001 .....	1.4	*	*	*	*	*	1.0	2.3	4.3	7.0	11.7	24.3
2000 .....	1.4	*	*	*	*	*	0.8	2.4	3.8	7.2	13.9	31.7
1999 .....	1.5	*	*	*	*	*	1.2	2.4	4.5	8.3	12.2	28.5
<b>Female</b>												
2002 .....	0.9	*	*	*	*	*	*	*	1.6	2.9	6.1	18.4
2001 .....	0.9	*	*	*	*	*	*	*	1.5	2.7	7.4	19.7
2000 .....	1.0	*	*	*	*	*	*	1.1	1.7	3.3	8.0	24.9
1999 .....	1.1	*	*	*	*	*	*	*	*	3.9	10.3	24.4
<b>Diseases of appendix (K35-K38)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.9	2.0
2001 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.1
2000 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
1999 .....	0.1	*	*	0.1	*	*	0.1	0.1	0.2	0.4	0.8	1.5
<b>Male</b>												
2002 .....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.3	2.2
2001 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.8
2000 .....	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999 .....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.9
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Diseases of appendix (K35-K38)</b>												
<b>Hispanic</b>												
Both sexes												
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
Both sexes												
2002 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	2.1
2001 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.2
2000 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.7
1999 .....	0.1	*	*	*	*	*	0.0	0.1	0.2	0.4	0.8	1.5
Male												
2002 .....	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.3	2.3
2001 .....	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.0	3.0
2000 .....	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999 .....	0.2	*	*	*	*	*	*	0.2	0.3	0.4	1.0	3.0
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9
<b>Non-Hispanic white</b>												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.0	2.1
2001 .....	0.2	*	*	*	*	*	*	0.1	0.1	0.5	0.7	2.1
2000 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.8
1999 .....	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.6
Male												
2002 .....	0.2	*	*	*	*	*	*	0.2	0.3	0.6	1.4	2.1
2001 .....	0.2	*	*	*	*	*	*	*	0.2	0.7	1.0	2.6
2000 .....	0.2	*	*	*	*	*	*	*	0.3	0.4	1.2	2.1
1999 .....	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.9	3.1
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.1
2001 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.4	2.0
2000 .....	0.2	*	*	*	*	*	*	*	*	0.3	0.7	1.6
1999 .....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Diseases of appendix (K35-K38)</b>												
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Hernia (K40-K46)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	0.6	0.7	*	*	*	*	0.1	0.2	0.5	1.2	3.8	12.8
2001 .....	0.5	1.1	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.6
2000 .....	0.5	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.1
1999 .....	0.5	1.2	*	*	*	*	0.1	0.2	0.4	1.2	3.7	12.7
<b>Male</b>												
2002 .....	0.4	*	*	*	*	*	*	0.2	0.4	1.2	3.7	11.7
2001 .....	0.4	1.4	*	*	*	*	*	0.2	0.5	1.1	3.5	12.0
2000 .....	0.4	1.4	*	*	*	*	*	0.2	0.4	1.1	4.0	12.6
1999 .....	0.4	1.3	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.2
<b>Female</b>												
2002 .....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.3
2001 .....	0.6	*	*	*	*	*	*	0.2	0.5	1.2	3.6	12.8
2000 .....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.3
1999 .....	0.6	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.6
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	4.3	12.5
2000 .....	0.2	*	*	*	*	*	*	*	*	*	4.5	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
Hispanic												
Female												
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.6	0.8	*	*	*	*	0.1	0.2	0.5	1.2	3.8	13.0
2001 .....	0.6	1.0	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.5
2000 .....	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.7	13.1
1999 .....	0.6	1.3	*	*	*	*	0.1	0.1	0.5	1.1	3.7	12.7
Male												
2002 .....	0.5	*	*	*	*	*	*	0.2	0.4	1.3	3.7	11.9
2001 .....	0.5	1.2	*	*	*	*	*	0.2	0.5	1.2	3.5	12.1
2000 .....	0.5	1.4	*	*	*	*	*	0.2	0.4	1.1	3.9	12.8
1999 .....	0.5	1.5	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.3
Female												
2002 .....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.4
2001 .....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.5	12.7
2000 .....	0.7	*	*	*	*	*	*	0.2	0.5	1.1	3.6	13.2
1999 .....	0.7	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.4
Non-Hispanic white												
Both sexes												
2002 .....	0.7	0.9	*	*	*	*	0.1	0.2	0.4	1.2	4.0	13.6
2001 .....	0.6	0.9	*	*	*	*	*	0.2	0.5	1.2	3.5	12.8
2000 .....	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.7	13.3
1999 .....	0.6	1.4	*	*	*	*	0.1	0.1	0.5	1.1	3.7	13.0
Male												
2002 .....	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.8	12.2
2001 .....	0.5	*	*	*	*	*	*	0.2	0.5	1.2	3.5	11.9
2000 .....	0.5	*	*	*	*	*	*	0.2	0.4	1.1	4.0	13.0
1999 .....	0.5	*	*	*	*	*	*	0.2	0.6	1.1	4.1	13.0
Female												
2002 .....	0.8	*	*	*	*	*	*	0.2	0.4	1.2	4.1	14.2
2001 .....	0.8	*	*	*	*	*	*	0.2	0.5	1.1	3.5	13.1
2000 .....	0.8	*	*	*	*	*	*	0.2	0.5	1.1	3.6	13.4
1999 .....	0.7	*	*	*	*	*	*	*	0.4	1.1	3.4	12.9
Non-Hispanic black												
Both sexes												
2002 .....	0.4	*	*	*	*	*	*	*	0.8	1.6	3.2	7.9
2001 .....	0.4	*	*	*	*	*	*	*	0.9	1.5	3.6	11.2
2000 .....	0.4	*	*	*	*	*	*	*	*	1.3	4.0	12.7
1999 .....	0.4	*	*	*	*	*	*	*	*	1.5	4.2	10.0
Male												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Hernia (K40-K46)</b>												
Non-Hispanic black												
Female												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	9.0
2000 .....	0.5	*	*	*	*	*	*	*	*	*	4.3	12.6
1999 .....	0.5	*	*	*	*	*	*	*	*	*	4.2	*
<b>Chronic liver disease and cirrhosis (K70,K73-K74)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	9.5	*	*	*	0.1	0.9	7.0	18.0	22.9	29.4	31.4	21.4
2001 .....	9.5	*	*	*	0.1	1.0	7.4	18.5	22.7	30.0	30.2	22.2
2000 .....	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8	29.8	31.0	23.1
1999 .....	9.4	*	*	*	0.1	1.0	7.3	17.4	23.7	30.6	31.9	23.2
Male												
2002 .....	12.3	*	*	*	*	1.2	9.3	26.9	33.1	38.5	38.4	28.4
2001 .....	12.4	*	*	*	*	1.3	9.8	27.5	33.1	40.2	36.8	32.0
2000 .....	12.5	*	*	*	*	1.4	10.3	26.9	34.4	39.9	38.4	34.3
1999 .....	12.5	*	*	*	*	1.3	10.3	26.9	34.6	41.6	39.6	33.1
Female												
2002 .....	6.7	*	*	*	*	0.7	4.8	9.5	13.5	21.9	26.7	18.4
2001 .....	6.7	*	*	*	*	0.7	5.0	9.8	13.1	21.5	25.9	18.1
2000 .....	6.5	*	*	*	*	0.7	4.7	8.8	14.0	21.5	26.1	18.5
1999 .....	6.4	*	*	*	*	0.8	4.4	8.3	13.7	21.6	26.9	19.3
Hispanic												
Both sexes												
2002 .....	8.8	*	*	*	*	1.1	8.7	26.9	38.4	56.2	53.5	45.5
2001 .....	8.9	*	*	*	*	1.2	9.7	28.0	39.0	54.4	60.4	36.9
2000 .....	9.0	*	*	*	*	1.4	9.8	27.2	41.4	58.3	63.8	48.4
1999 .....	8.8	*	*	*	*	1.3	10.5	26.6	39.0	56.3	59.4	51.2
Male												
2002 .....	12.2	*	*	*	*	1.6	13.5	44.2	60.0	77.8	64.4	45.2
2001 .....	12.7	*	*	*	*	1.8	15.0	46.4	63.9	77.1	76.0	41.9
2000 .....	12.7	*	*	*	*	2.3	15.0	45.6	66.1	83.4	80.1	60.5
1999 .....	12.3	*	*	*	*	1.8	16.5	44.3	62.6	79.1	70.9	66.9
Female												
2002 .....	5.2	*	*	*	*	*	3.5	9.7	18.9	38.9	46.1	45.6
2001 .....	5.0	*	*	*	*	*	4.0	9.9	17.0	36.5	49.9	34.4
2000 .....	5.1	*	*	*	*	*	4.1	9.3	19.5	38.7	52.8	42.5
1999 .....	5.0	*	*	*	*	*	4.1	9.3	18.1	38.4	51.7	43.7
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	9.5	*	*	*	0.1	0.9	6.7	17.0	21.5	27.5	30.2	20.4
2001 .....	9.5	*	*	*	0.1	0.9	7.1	17.5	21.3	28.2	28.7	21.6
2000 .....	9.4	*	*	*	0.1	1.0	7.1	16.7	22.3	27.9	29.5	22.1
1999 .....	9.5	*	*	*	*	1.0	6.9	16.5	22.5	29.0	30.7	22.1
Male												
2002 .....	12.2	*	*	*	*	1.1	8.6	24.9	30.7	35.7	37.0	27.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Chronic liver disease and cirrhosis (K70,K73-K74)</b>												
<b>Non-Hispanic <sup>4</sup></b>												
<b>Male</b>												
2001 .....	12.3	*	*	*	*	1.2	9.1	25.6	30.6	37.6	34.9	31.4
2000 .....	12.4	*	*	*	*	1.2	9.6	25.0	31.8	37.1	36.5	33.0
1999 .....	12.5	*	*	*	*	1.2	9.5	25.2	32.4	39.3	38.2	31.6
<b>Female</b>												
2002 .....	6.9	*	*	*	*	0.7	4.9	9.4	13.0	20.6	25.8	17.3
2001 .....	6.9	*	*	*	*	0.7	5.1	9.8	12.7	20.4	24.7	17.5
2000 .....	6.7	*	*	*	*	0.7	4.7	8.7	13.6	20.3	24.9	17.7
1999 .....	6.6	*	*	*	*	0.8	4.4	8.1	13.3	20.6	25.8	18.3
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	10.2	*	*	*	*	0.8	6.9	16.8	21.4	28.5	31.5	21.2
2001 .....	10.1	*	*	*	*	0.9	7.0	17.0	21.0	28.9	29.9	22.5
2000 .....	10.0	*	*	*	*	0.8	7.0	16.2	22.1	28.4	30.6	22.8
1999 .....	9.9	*	*	*	*	0.9	6.6	15.8	21.8	29.5	31.7	22.6
<b>Male</b>												
2002 .....	13.2	*	*	*	*	1.0	8.8	24.7	30.5	36.6	38.7	28.7
2001 .....	13.0	*	*	*	*	1.1	8.9	24.8	29.7	38.1	36.3	32.9
2000 .....	13.1	*	*	*	*	1.1	9.4	24.4	31.5	37.7	37.8	34.1
1999 .....	13.1	*	*	*	*	1.1	9.1	24.2	31.5	39.7	39.2	32.7
<b>Female</b>												
2002 .....	7.5	*	*	*	*	0.6	5.0	9.0	12.9	21.5	26.8	18.0
2001 .....	7.4	*	*	*	*	0.7	5.1	9.3	12.8	21.1	25.7	18.2
2000 .....	7.0	*	*	*	*	0.6	4.5	8.0	13.2	20.6	25.9	18.2
1999 .....	6.9	*	*	*	*	0.7	4.1	7.5	12.7	21.0	26.7	18.6
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	7.0	*	*	*	*	0.9	5.6	21.2	25.3	21.8	18.3	12.1
2001 .....	7.7	*	*	*	*	0.9	7.5	22.9	26.3	25.3	18.2	10.6
2000 .....	7.7	*	*	*	*	1.1	8.0	22.6	26.1	26.5	18.1	14.0
1999 .....	8.0	*	*	*	*	1.0	8.2	23.0	29.7	26.9	21.7	16.8
<b>Male</b>												
2002 .....	9.3	*	*	*	*	1.1	6.9	30.7	38.2	32.4	22.0	*
2001 .....	10.7	*	*	*	*	1.0	10.4	35.1	42.0	39.7	22.5	*
2000 .....	10.3	*	*	*	*	1.1	10.7	33.4	38.5	38.2	25.0	*
1999 .....	11.0	*	*	*	*	1.1	11.3	35.8	43.8	41.0	30.0	*
<b>Female</b>												
2002 .....	5.0	*	*	*	*	0.8	4.5	13.2	15.0	14.2	16.2	10.5
2001 .....	4.9	*	*	*	*	0.7	5.0	12.5	13.7	15.1	15.8	9.8
2000 .....	5.4	*	*	*	*	1.1	5.7	13.4	16.2	18.3	14.2	11.8
1999 .....	5.4	*	*	*	*	1.0	5.5	12.1	18.5	16.9	17.0	15.5
<b>Alcoholic liver disease (K70)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	4.2	*	*	*	*	0.6	4.4	10.3	11.6	10.1	5.7	2.1
2001 .....	4.3	*	*	*	*	0.7	4.8	10.7	11.5	10.1	5.7	2.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
All origins <sup>3</sup>												
Both sexes												
2000 .....	4.3	*	*	*	*	0.7	4.9	10.3	12.2	10.3	6.1	2.2
1999 .....	4.3	*	*	*	*	0.6	4.8	10.4	12.0	10.5	6.5	2.8
Male												
2002 .....	6.2	*	*	*	*	0.8	5.8	15.6	18.3	16.5	9.7	4.7
2001 .....	6.4	*	*	*	*	0.9	6.5	16.1	18.4	16.6	10.1	5.5
2000 .....	6.5	*	*	*	*	1.0	6.8	16.0	19.0	17.3	11.5	5.3
1999 .....	6.5	*	*	*	*	0.8	6.8	16.3	19.2	18.1	11.2	6.1
Female												
2002 .....	2.3	*	*	*	*	0.4	3.0	5.2	5.4	4.8	3.1	1.0
2001 .....	2.3	*	*	*	*	0.4	3.1	5.5	5.1	4.7	2.8	1.0
2000 .....	2.2	*	*	*	*	0.4	3.1	4.8	6.0	4.5	2.5	1.0
1999 .....	2.1	*	*	*	*	0.5	2.9	4.7	5.4	4.3	3.4	1.5
Hispanic												
Both sexes												
2002 .....	4.3	*	*	*	*	0.7	5.5	16.1	19.9	20.8	10.5	*
2001 .....	4.7	*	*	*	*	0.9	6.3	17.1	21.4	21.1	17.0	*
2000 .....	4.8	*	*	*	*	1.0	6.9	16.8	23.4	22.9	15.4	*
1999 .....	4.6	*	*	*	*	0.8	6.9	16.8	21.7	20.6	14.4	*
Male												
2002 .....	7.1	*	*	*	*	1.2	9.0	27.0	36.3	40.3	18.1	*
2001 .....	7.7	*	*	*	*	1.5	10.0	29.3	38.9	41.9	30.4	*
2000 .....	7.9	*	*	*	*	1.8	10.6	29.3	42.8	46.6	33.9	*
1999 .....	7.4	*	*	*	*	1.2	10.8	29.1	40.0	39.5	27.1	*
Female												
2002 .....	1.4	*	*	*	*	*	1.8	5.1	5.1	5.3	*	*
2001 .....	1.5	*	*	*	*	*	2.2	5.1	5.9	4.7	7.9	*
2000 .....	1.4	*	*	*	*	*	2.9	4.5	6.2	4.3	*	*
1999 .....	1.5	*	*	*	*	*	2.8	4.9	5.6	5.7	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	4.2	*	*	*	*	0.6	4.2	9.7	10.8	9.3	5.5	1.9
2001 .....	4.2	*	*	*	*	0.6	4.6	10.1	10.7	9.3	5.2	2.2
2000 .....	4.2	*	*	*	*	0.6	4.7	9.7	11.3	9.5	5.7	2.2
1999 .....	4.2	*	*	*	*	0.6	4.6	9.8	11.3	9.9	6.1	2.5
Male												
2002 .....	6.0	*	*	*	*	0.7	5.3	14.4	16.7	14.8	9.2	4.3
2001 .....	6.2	*	*	*	*	0.8	5.9	14.8	16.7	14.9	9.2	5.1
2000 .....	6.2	*	*	*	*	0.8	6.3	14.7	17.1	15.4	10.5	5.0
1999 .....	6.4	*	*	*	*	0.7	6.2	15.1	17.7	16.7	10.5	5.5
Female												
2002 .....	2.4	*	*	*	*	0.5	3.1	5.2	5.4	4.8	3.0	0.9
2001 .....	2.4	*	*	*	*	0.5	3.2	5.5	5.0	4.7	2.5	1.0
2000 .....	2.3	*	*	*	*	0.4	3.1	4.8	5.9	4.5	2.5	1.0
1999 .....	2.2	*	*	*	*	0.5	2.9	4.7	5.3	4.2	3.3	1.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Alcoholic liver disease (K70)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	4.4	*	*	*	*	0.5	4.3	9.7	10.8	9.5	5.5	1.9
2001 .....	4.4	*	*	*	*	0.6	4.5	9.9	10.5	9.3	5.3	2.1
2000 .....	4.3	*	*	*	*	0.6	4.5	9.4	11.1	9.5	5.6	2.2
1999 .....	4.3	*	*	*	*	0.6	4.4	9.5	10.8	9.7	6.0	2.4
<b>Male</b>												
2002 .....	6.4	*	*	*	*	0.7	5.4	14.4	16.7	14.9	9.4	4.1
2001 .....	6.4	*	*	*	*	0.8	5.8	14.7	16.4	14.6	9.3	4.8
2000 .....	6.5	*	*	*	*	0.8	6.1	14.5	17.0	15.4	10.6	5.1
1999 .....	6.5	*	*	*	*	0.7	6.0	14.7	17.1	16.3	10.3	5.1
<b>Female</b>												
2002 .....	2.4	*	*	*	*	0.4	3.2	5.1	5.2	4.9	3.0	1.0
2001 .....	2.4	*	*	*	*	0.4	3.2	5.3	5.0	4.8	2.5	1.0
2000 .....	2.2	*	*	*	*	0.4	3.0	4.4	5.5	4.6	2.4	1.0
1999 .....	2.2	*	*	*	*	0.5	2.8	4.3	4.9	4.2	3.2	1.3
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	3.2	*	*	*	*	0.5	3.0	10.2	12.3	8.7	5.1	*
2001 .....	3.7	*	*	*	*	0.5	4.4	11.7	13.1	10.8	4.6	*
2000 .....	3.9	*	*	*	*	0.7	5.1	12.3	13.5	10.4	5.6	*
1999 .....	4.2	*	*	*	*	0.5	5.1	12.9	15.4	11.9	7.2	*
<b>Male</b>												
2002 .....	4.6	*	*	*	*	*	3.9	15.1	18.6	15.1	8.8	*
2001 .....	5.5	*	*	*	*	*	6.1	17.4	21.7	20.0	7.8	*
2000 .....	5.5	*	*	*	*	*	7.1	17.9	19.6	18.4	10.2	*
1999 .....	6.1	*	*	*	*	*	6.9	19.7	24.6	21.9	12.8	*
<b>Female</b>												
2002 .....	2.1	*	*	*	*	*	2.2	6.1	7.2	4.2	*	*
2001 .....	2.2	*	*	*	*	*	2.8	6.8	6.2	4.3	*	*
2000 .....	2.5	*	*	*	*	*	3.4	7.5	8.6	4.8	*	*
1999 .....	2.4	*	*	*	*	*	3.5	7.0	8.1	4.8	4.0	*
<b>Other chronic liver disease and cirrhosis (K73-K74)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	5.2	*	*	*	0.1	0.3	2.6	7.7	11.3	19.3	25.7	19.3
2001 .....	5.2	*	*	*	*	0.3	2.6	7.8	11.2	19.9	24.5	19.9
2000 .....	5.1	*	*	*	0.1	0.3	2.5	7.3	11.6	19.5	24.9	20.9
1999 .....	5.1	*	*	*	*	0.4	2.5	7.0	11.7	20.1	25.5	20.4
<b>Male</b>												
2002 .....	6.1	*	*	*	*	0.4	3.5	11.2	14.8	22.0	28.7	23.7
2001 .....	6.1	*	*	*	*	0.4	3.4	11.4	14.7	23.6	26.6	26.5
2000 .....	6.0	*	*	*	*	0.4	3.5	10.9	15.4	22.7	26.9	29.0
1999 .....	6.0	*	*	*	*	0.5	3.5	10.6	15.3	23.6	28.4	27.0
<b>Female</b>												
2002 .....	4.5	*	*	*	*	0.3	1.8	4.3	8.1	17.1	23.7	17.4
2001 .....	4.4	*	*	*	*	0.2	1.9	4.3	8.0	16.8	23.1	17.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
All origins <sup>3</sup>												
Female												
2002 .....	4.3	*	*	*	*	0.3	1.6	4.0	8.0	16.9	23.6	17.5
1999 .....	4.3	*	*	*	*	0.3	1.4	3.6	8.3	17.3	23.5	17.7
Hispanic												
Both sexes												
2002 .....	4.5	*	*	*	*	0.4	3.2	10.8	18.5	35.3	43.0	38.8
2001 .....	4.3	*	*	*	*	*	3.5	10.9	17.6	33.3	43.5	30.6
2000 .....	4.3	*	*	*	*	0.4	2.9	10.5	18.0	35.4	48.4	44.5
1999 .....	4.2	*	*	*	*	0.5	3.6	9.7	17.2	35.7	45.0	40.0
Male												
2002 .....	5.1	*	*	*	*	*	4.6	17.1	23.8	37.5	46.4	*
2001 .....	5.0	*	*	*	*	*	5.1	17.1	25.0	35.1	45.6	*
2000 .....	4.8	*	*	*	*	0.6	4.4	16.2	23.3	36.7	46.2	50.4
1999 .....	4.9	*	*	*	*	0.6	5.7	15.1	22.6	39.5	43.8	45.3
Female												
2002 .....	3.8	*	*	*	*	*	1.7	4.6	13.8	33.6	40.8	42.3
2001 .....	3.4	*	*	*	*	*	1.8	4.7	11.0	31.8	42.0	33.5
2000 .....	3.7	*	*	*	*	*	1.3	4.8	13.3	34.3	49.9	41.5
1999 .....	3.5	*	*	*	*	*	1.3	4.5	12.4	32.7	45.8	37.4
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	5.4	*	*	*	*	0.3	2.5	7.3	10.7	18.2	24.8	18.5
2001 .....	5.3	*	*	*	*	0.3	2.5	7.5	10.7	18.9	23.6	19.4
2000 .....	5.2	*	*	*	0.1	0.3	2.5	7.0	11.0	18.4	23.8	19.9
1999 .....	5.2	*	*	*	*	0.3	2.3	6.7	11.2	19.2	24.6	19.6
Male												
2002 .....	6.2	*	*	*	*	0.4	3.3	10.6	14.0	20.9	27.7	23.4
2001 .....	6.2	*	*	*	*	0.4	3.1	10.8	13.8	22.8	25.7	26.4
2000 .....	6.1	*	*	*	*	0.3	3.3	10.3	14.7	21.7	26.0	28.0
1999 .....	6.1	*	*	*	*	0.4	3.3	10.1	14.7	22.6	27.7	26.2
Female												
2002 .....	4.6	*	*	*	*	0.3	1.8	4.2	7.6	15.8	22.8	16.4
2001 .....	4.5	*	*	*	*	0.2	1.9	4.3	7.7	15.7	22.2	16.5
2000 .....	4.4	*	*	*	*	0.3	1.6	3.9	7.6	15.8	22.4	16.7
1999 .....	4.4	*	*	*	*	0.3	1.4	3.5	8.0	16.3	22.6	17.0
Non-Hispanic white												
Both sexes												
2002 .....	5.9	*	*	*	*	0.3	2.6	7.1	10.7	19.0	26.0	19.3
2001 .....	5.8	*	*	*	*	0.3	2.5	7.1	10.5	19.6	24.7	20.4
2000 .....	5.7	*	*	*	*	0.3	2.4	6.7	11.0	18.9	25.0	20.6
1999 .....	5.6	*	*	*	*	0.3	2.3	6.4	11.0	19.8	25.6	20.2
Male												
2002 .....	6.7	*	*	*	*	0.3	3.4	10.3	13.7	21.7	29.3	24.7
2001 .....	6.6	*	*	*	*	0.4	3.0	10.2	13.3	23.5	26.9	28.1
2000 .....	6.6	*	*	*	*	0.3	3.4	9.9	14.6	22.3	27.2	29.0
1999 .....	6.6	*	*	*	*	0.4	3.2	9.5	14.4	23.3	28.9	27.5

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
Non-Hispanic white												
Female												
2002 .....	5.0	*	*	*	*	0.2	1.8	3.9	7.7	16.6	23.8	17.0
2001 .....	5.0	*	*	*	*	0.3	1.9	4.1	7.9	16.3	23.2	17.2
2000 .....	4.8	*	*	*	*	0.3	1.5	3.6	7.6	16.0	23.5	17.2
1999 .....	4.8	*	*	*	*	0.2	1.4	3.2	7.7	16.8	23.5	17.3
Non-Hispanic black												
Both sexes												
2002 .....	3.8	*	*	*	*	0.4	2.6	11.0	13.0	13.1	13.2	9.7
2001 .....	3.9	*	*	*	*	*	3.2	11.2	13.2	14.5	13.6	7.8
2000 .....	3.8	*	*	*	*	0.4	2.9	10.3	12.7	16.1	12.6	12.4
1999 .....	3.9	*	*	*	*	0.5	3.1	10.1	14.3	15.0	14.5	12.3
Male												
2002 .....	4.7	*	*	*	*	*	2.9	15.5	19.6	17.3	13.2	*
2001 .....	5.3	*	*	*	*	*	4.3	17.6	20.3	19.7	14.7	*
2000 .....	4.8	*	*	*	*	*	3.6	15.5	18.9	19.8	14.8	*
1999 .....	4.9	*	*	*	*	*	4.5	16.0	19.3	19.1	17.2	*
Female												
2002 .....	3.0	*	*	*	*	*	2.3	7.1	7.7	10.1	13.3	9.6
2001 .....	2.7	*	*	*	*	*	2.1	5.7	7.5	10.8	13.0	*
2000 .....	2.9	*	*	*	*	*	2.3	5.9	7.7	13.5	11.3	11.3
1999 .....	2.9	*	*	*	*	*	1.9	5.0	10.4	12.1	13.0	12.9
Cholelithiasis and other disorders of gallbladder (K80-K82)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.0	*	*	*	*	0.1	0.1	0.3	0.8	2.8	7.8	22.9
2001 .....	1.0	*	*	*	*	*	0.1	0.3	1.0	2.9	7.5	23.7
2000 .....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	7.5	24.8
1999 .....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.6	7.8	23.9
Male												
2002 .....	0.9	*	*	*	*	*	0.1	0.4	1.0	3.1	9.8	24.1
2001 .....	1.0	*	*	*	*	*	0.1	0.4	1.2	3.2	9.3	26.9
2000 .....	0.9	*	*	*	*	*	0.1	0.3	0.9	2.9	9.4	27.6
1999 .....	0.9	*	*	*	*	*	0.1	0.3	1.1	3.0	9.3	27.7
Female												
2002 .....	1.1	*	*	*	*	*	0.1	0.2	0.7	2.5	6.4	22.4
2001 .....	1.1	*	*	*	*	*	*	0.3	0.8	2.6	6.3	22.4
2000 .....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	6.2	23.7
1999 .....	1.1	*	*	*	*	*	0.1	0.4	0.7	2.2	6.8	22.4
Hispanic												
Both sexes												
2002 .....	0.5	*	*	*	*	*	*	*	1.3	3.6	9.2	24.4
2001 .....	0.5	*	*	*	*	*	*	*	1.2	4.5	8.5	31.3
2000 .....	0.5	*	*	*	*	*	*	*	*	3.8	10.1	29.2
1999 .....	0.5	*	*	*	*	*	*	*	1.2	3.3	11.9	32.3
Male												
2002 .....	0.5	*	*	*	*	*	*	*	*	4.0	9.4	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Cholelithiasis and other disorders of gallbladder (K80-K82)</b>												
<b>Hispanic</b>												
<b>Male</b>												
2001 .....	0.5	*	*	*	*	*	*	*	*	5.0	11.2	*
2000 .....	0.5	*	*	*	*	*	*	*	*	5.3	15.7	*
1999 .....	0.5	*	*	*	*	*	*	*	*	*	15.1	*
<b>Female</b>												
2002 .....	0.5	*	*	*	*	*	*	*	*	3.2	9.0	23.2
2001 .....	0.6	*	*	*	*	*	*	*	*	4.1	6.6	30.7
2000 .....	0.5	*	*	*	*	*	*	*	*	*	*	31.7
1999 .....	0.6	*	*	*	*	*	*	*	*	*	9.7	32.2
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.7	7.7	22.8
2001 .....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.8	7.5	23.4
2000 .....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.3	7.4	24.6
1999 .....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	7.6	23.6
<b>Male</b>												
2002 .....	1.0	*	*	*	*	*	0.1	0.4	0.9	3.0	9.8	23.9
2001 .....	1.0	*	*	*	*	*	0.1	0.4	1.1	3.1	9.2	26.6
2000 .....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.7	9.2	27.7
1999 .....	1.0	*	*	*	*	*	0.1	0.3	1.0	3.0	9.0	27.5
<b>Female</b>												
2002 .....	1.2	*	*	*	*	*	0.1	0.2	0.6	2.4	6.3	22.4
2001 .....	1.2	*	*	*	*	*	*	0.3	0.8	2.5	6.3	22.1
2000 .....	1.2	*	*	*	*	*	0.1	0.3	0.6	2.0	6.2	23.4
1999 .....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.6	22.0
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.6	7.5	23.4
2001 .....	1.2	*	*	*	*	*	0.1	0.3	0.8	2.6	7.3	23.6
2000 .....	1.2	*	*	*	*	*	0.1	0.2	0.7	2.2	7.3	25.3
1999 .....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.4	7.6	24.0
<b>Male</b>												
2002 .....	1.1	*	*	*	*	*	*	0.3	0.9	3.0	9.8	24.5
2001 .....	1.1	*	*	*	*	*	*	0.4	1.0	2.9	9.2	26.4
2000 .....	1.0	*	*	*	*	*	*	0.2	0.8	2.6	9.3	28.9
1999 .....	1.1	*	*	*	*	*	*	0.3	0.9	2.9	9.3	27.8
<b>Female</b>												
2002 .....	1.3	*	*	*	*	*	*	0.2	0.6	2.3	6.1	22.9
2001 .....	1.3	*	*	*	*	*	*	0.2	0.7	2.2	6.1	22.4
2000 .....	1.3	*	*	*	*	*	*	0.2	0.6	1.8	6.0	23.9
1999 .....	1.3	*	*	*	*	*	*	0.3	0.6	1.9	6.5	22.5
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	0.7	*	*	*	*	*	*	0.6	1.1	3.3	9.6	15.1
2001 .....	0.9	*	*	*	*	*	*	0.7	1.7	4.8	8.3	20.0
2000 .....	0.8	*	*	*	*	*	*	0.7	1.2	3.8	8.3	18.1
1999 .....	0.7	*	*	*	*	*	0.4	0.5	1.5	3.7	7.3	15.5

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Cholelithiasis and other disorders of gallbladder (K80-K82)</b>												
<b>Non-Hispanic black</b>												
<b>Male</b>												
2002 .....	0.6	*	*	*	*	*	*	*	*	*	10.8	*
2001 .....	0.8	*	*	*	*	*	*	*	2.2	5.0	6.3	27.7
2000 .....	0.6	*	*	*	*	*	*	*	*	3.4	8.0	*
1999 .....	0.6	*	*	*	*	*	*	*	*	3.8	7.2	*
<b>Female</b>												
2002 .....	0.8	*	*	*	*	*	*	*	*	3.8	9.0	16.3
2001 .....	1.0	*	*	*	*	*	*	*	*	4.6	9.4	17.1
2000 .....	1.0	*	*	*	*	*	*	*	*	4.0	8.5	19.6
1999 .....	0.8	*	*	*	*	*	*	*	*	3.6	7.4	15.5
<b>Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	14.2	4.3	*	0.1	0.2	0.7	1.7	4.7	13.0	39.2	109.1	288.6
2001 .....	13.9	3.3	*	*	0.2	0.6	1.7	4.6	13.0	40.2	104.2	287.7
2000 .....	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8	38.0	100.8	277.8
1999 .....	12.7	4.4	*	0.1	0.2	0.6	1.6	4.0	12.0	37.1	97.6	268.9
<b>Male</b>												
2002 .....	13.9	5.3	*	*	0.2	0.6	1.8	5.5	14.3	45.8	136.2	385.5
2001 .....	13.5	3.7	*	*	0.2	0.6	1.8	5.5	14.1	46.1	130.0	393.7
2000 .....	12.9	5.3	*	*	0.2	0.6	1.9	5.3	14.2	42.9	127.3	380.7
1999 .....	12.4	4.6	*	*	0.2	0.6	1.8	4.7	13.4	42.3	125.3	371.1
<b>Female</b>												
2002 .....	14.5	3.2	*	*	0.2	0.7	1.5	4.0	11.8	33.7	91.1	246.6
2001 .....	14.2	2.9	*	*	0.2	0.6	1.6	3.9	11.9	35.2	87.1	243.3
2000 .....	13.6	3.2	*	*	0.2	0.6	1.4	3.5	11.5	34.0	83.5	235.9
1999 .....	13.0	4.1	*	*	0.2	0.6	1.3	3.4	10.8	33.0	79.6	227.7
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	4.9	4.2	*	*	*	0.5	1.1	4.3	13.5	36.2	88.7	201.2
2001 .....	5.0	*	*	*	*	0.4	1.4	3.9	12.8	41.4	94.5	222.0
2000 .....	4.7	3.2	*	*	*	0.4	1.0	3.5	13.6	37.5	84.7	229.6
1999 .....	4.7	2.6	*	*	*	0.3	1.3	3.6	14.0	35.6	88.7	232.2
<b>Male</b>												
2002 .....	4.7	4.9	*	*	*	*	1.3	4.8	15.1	40.9	112.0	235.9
2001 .....	4.7	*	*	*	*	*	1.6	3.8	15.2	43.3	112.2	281.9
2000 .....	4.4	*	*	*	*	*	1.4	4.4	14.3	40.9	101.3	292.2
1999 .....	4.7	*	*	*	*	*	1.2	4.4	17.8	39.1	110.5	336.4
<b>Female</b>												
2002 .....	5.1	*	*	*	*	0.6	1.0	3.8	12.1	32.4	72.8	184.0
2001 .....	5.4	*	*	*	*	*	1.1	4.0	10.7	39.8	82.5	192.7
2000 .....	4.9	*	*	*	*	*	*	2.5	12.9	34.8	73.6	198.8
1999 .....	4.7	*	*	*	*	*	1.3	2.9	10.7	32.9	74.2	182.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	15.6	4.3	*	*	0.2	0.7	1.7	4.7	12.9	39.3	109.8	291.2
2001 .....	15.1	3.5	*	*	0.2	0.6	1.7	4.7	12.9	40.0	104.3	289.6
2000 .....	14.4	4.4	*	0.1	0.2	0.7	1.7	4.4	12.6	37.9	101.2	279.0
1999 .....	13.8	4.8	*	0.1	0.2	0.7	1.6	4.1	11.8	37.1	97.7	269.2
Male												
2002 .....	15.4	5.4	*	*	0.2	0.7	1.9	5.6	14.2	45.9	136.9	391.0
2001 .....	14.8	3.7	*	*	0.2	0.6	1.9	5.6	14.0	46.1	130.6	397.6
2000 .....	14.1	5.4	*	*	0.2	0.7	2.0	5.4	14.1	42.9	128.2	384.0
1999 .....	13.5	4.8	*	*	0.2	0.7	1.9	4.7	13.0	42.3	125.7	370.9
Female												
2002 .....	15.8	3.2	*	*	0.2	0.7	1.6	3.9	11.7	33.7	91.8	248.2
2001 .....	15.4	3.3	*	*	0.2	0.6	1.6	3.8	12.0	34.8	87.1	244.6
2000 .....	14.7	3.4	*	*	0.2	0.6	1.5	3.6	11.3	33.8	83.6	236.6
1999 .....	14.1	4.8	*	*	0.2	0.7	1.3	3.5	10.7	32.9	79.5	228.4
Non-Hispanic white												
Both sexes												
2002 .....	15.4	3.3	*	*	0.1	0.5	1.1	3.1	9.5	33.0	102.5	281.1
2001 .....	14.9	2.4	*	*	0.2	0.4	1.1	3.1	9.3	33.5	96.6	279.7
2000 .....	14.2	2.7	*	*	0.2	0.4	1.1	2.9	9.3	31.4	94.1	269.4
1999 .....	13.4	3.5	*	*	0.1	0.4	1.0	2.6	8.7	31.1	89.9	255.3
Male												
2002 .....	15.3	4.1	*	*	*	0.5	1.2	3.6	10.3	39.7	130.8	382.6
2001 .....	14.8	2.5	*	*	0.2	0.4	1.3	3.7	10.5	39.9	123.7	392.6
2000 .....	14.1	3.5	*	*	0.2	0.5	1.3	3.5	10.6	37.3	121.7	378.9
1999 .....	13.3	3.9	*	*	*	0.5	1.2	2.9	9.9	36.6	117.3	359.6
Female												
2002 .....	15.6	2.5	*	*	*	0.4	1.0	2.6	8.7	27.3	83.6	237.3
2001 .....	15.0	2.3	*	*	0.2	0.4	0.9	2.4	8.3	28.0	78.6	232.7
2000 .....	14.2	1.8	*	*	0.2	0.4	1.0	2.3	8.0	26.5	76.0	225.2
1999 .....	13.5	3.1	*	*	*	0.4	0.8	2.2	7.5	26.4	71.9	213.8
Non-Hispanic black												
Both sexes												
2002 .....	20.5	8.5	*	*	0.6	2.1	5.5	16.5	41.8	101.5	205.5	444.0
2001 .....	20.2	8.5	*	*	0.5	1.8	5.5	16.2	42.5	104.2	204.0	432.9
2000 .....	19.5	12.0	*	*	0.4	1.9	5.5	16.2	41.2	101.2	197.3	410.2
1999 .....	19.2	10.2	*	*	0.5	2.0	5.6	15.6	37.8	96.6	196.4	436.8
Male												
2002 .....	19.7	10.6	*	*	*	2.0	6.2	20.6	50.1	116.3	237.1	568.9
2001 .....	18.6	8.9	*	*	*	2.2	5.3	19.9	46.3	116.0	235.3	517.4
2000 .....	18.3	13.6	*	*	*	2.0	6.7	20.8	45.6	105.7	234.2	496.7
1999 .....	18.1	8.7	*	*	*	1.9	6.7	19.4	41.2	103.9	251.1	532.4
Female												
2002 .....	21.2	6.3	*	*	0.7	2.2	4.9	13.1	35.1	91.0	187.7	396.3
2001 .....	21.7	8.1	*	*	*	1.4	5.6	13.0	39.4	95.8	186.5	401.7
2000 .....	20.7	10.2	*	*	*	1.8	4.4	12.2	37.6	98.1	176.6	378.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Non-Hispanic black												
Female												
1999 .....	20.2	11.8	*	*	*	2.0	4.7	12.3	35.1	91.5	165.8	401.1
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2001 .....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
Male												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.8
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.7	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.9
1999 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2001 .....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.8
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
Non-Hispanic <sup>4</sup>												
Male												
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.8
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.7	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	0.9
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
Non-Hispanic white												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.8
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.8
2000 .....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.2
1999 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.5	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.6	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.8
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	1.0
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Non-Hispanic black												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07,N26)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.4	2.8
2001 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999 .....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.6	3.2
Male												
2002 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	3.8
2001 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000 .....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.5
1999 .....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.2	5.0
Female												
2002 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	2.3
2001 .....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000 .....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.1
1999 .....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	2.4
Hispanic												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	2.8
2001 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.7	3.2
Male												
2002 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	3.8
2001 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000 .....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.2
1999 .....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.3	5.1
Female												
2002 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	2.3
2001 .....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07,N26)												
Non-Hispanic <sup>4</sup>												
Female												
2000 .....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.2
1999 .....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.4
Non-Hispanic white												
Both sexes												
2002 .....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.6
2001 .....	0.2	*	*	*	*	*	*	*	0.1	0.6	1.1	2.8
2000 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.4
1999 .....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.6	2.9
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.7	1.7	3.8
2001 .....	0.2	*	*	*	*	*	*	*	*	0.8	1.7	4.0
2000 .....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.7	3.7
1999 .....	0.2	*	*	*	*	*	*	*	*	0.5	2.2	4.6
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	2.0
2001 .....	0.2	*	*	*	*	*	*	*	*	0.4	0.8	2.4
2000 .....	0.2	*	*	*	*	*	*	*	*	0.5	1.1	1.9
1999 .....	0.2	*	*	*	*	*	*	*	*	0.5	1.2	2.2
Non-Hispanic black												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	2.1	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Renal failure (N17-N19)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	13.9	4.1	*	*	0.2	0.6	1.6	4.6	12.7	38.6	107.2	284.8
2001 .....	13.6	3.1	*	*	0.2	0.5	1.6	4.5	12.7	39.4	102.6	284.0
2000 .....	13.0	4.0	*	*	0.2	0.6	1.6	4.2	12.5	37.2	98.9	273.7
1999 .....	12.4	4.3	*	0.1	0.2	0.6	1.5	3.9	11.7	36.4	95.3	264.8
Male												
2002 .....	13.7	5.1	*	*	0.2	0.6	1.8	5.4	14.0	45.0	134.0	380.7

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Renal failure (N17-N19)</b>												
<b>All origins <sup>3</sup></b>												
<b>Male</b>												
2001 .....	13.2	3.5	*	*	0.2	0.5	1.8	5.3	13.8	45.2	128.0	388.7
2000 .....	12.6	5.1	*	*	0.2	0.6	1.8	5.1	13.8	42.1	125.1	374.2
1999 .....	12.1	4.5	*	*	0.1	0.5	1.7	4.5	13.1	41.5	122.3	364.6
<b>Female</b>												
2002 .....	14.2	3.0	*	*	0.2	0.7	1.4	3.8	11.6	33.2	89.5	243.2
2001 .....	14.0	2.7	*	*	0.2	0.5	1.5	3.7	11.8	34.5	85.9	240.1
2000 .....	13.3	2.8	*	*	0.2	0.5	1.3	3.4	11.2	33.3	81.8	232.8
1999 .....	12.7	4.1	*	*	0.2	0.6	1.3	3.3	10.4	32.2	77.8	224.5
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	4.8	4.2	*	*	*	0.4	1.1	4.2	13.2	35.7	86.9	198.4
2001 .....	4.9	*	*	*	*	0.3	1.2	3.7	12.5	40.5	93.2	217.6
2000 .....	4.5	2.9	*	*	*	0.4	1.0	3.3	13.3	36.9	83.0	225.6
1999 .....	4.5	*	*	*	*	*	1.2	3.5	13.6	34.4	86.9	228.0
<b>Male</b>												
2002 .....	4.6	4.9	*	*	*	*	1.3	4.8	14.4	40.7	109.2	232.6
2001 .....	4.6	*	*	*	*	*	1.4	3.7	14.6	42.5	110.4	278.1
2000 .....	4.3	*	*	*	*	*	1.3	4.2	13.8	40.5	99.3	280.1
1999 .....	4.5	*	*	*	*	*	1.1	4.3	17.2	38.2	109.0	330.0
<b>Female</b>												
2002 .....	5.0	*	*	*	*	0.6	0.8	3.6	12.0	31.8	71.7	181.5
2001 .....	5.2	*	*	*	*	*	1.0	3.7	10.5	38.9	81.6	188.0
2000 .....	4.8	*	*	*	*	*	*	2.4	12.8	34.0	72.0	198.8
1999 .....	4.5	*	*	*	*	*	1.3	2.7	10.4	31.3	72.2	178.8
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	15.3	4.1	*	*	0.2	0.7	1.7	4.6	12.6	38.6	107.9	287.4
2001 .....	14.9	3.3	*	*	0.2	0.6	1.7	4.6	12.7	39.2	102.7	285.9
2000 .....	14.1	4.2	*	*	0.2	0.6	1.6	4.3	12.3	37.1	99.3	274.9
1999 .....	13.5	4.7	*	*	0.2	0.6	1.5	3.9	11.5	36.4	95.4	265.0
<b>Male</b>												
2002 .....	15.1	5.1	*	*	0.2	0.6	1.8	5.4	13.9	45.1	134.7	386.2
2001 .....	14.6	3.5	*	*	0.2	0.6	1.8	5.4	13.7	45.3	128.5	392.7
2000 .....	13.8	5.2	*	*	0.2	0.6	1.9	5.1	13.7	42.0	125.9	377.7
1999 .....	13.2	4.6	*	*	0.1	0.6	1.8	4.5	12.7	41.5	122.6	364.4
<b>Female</b>												
2002 .....	15.5	2.9	*	*	0.2	0.7	1.5	3.8	11.4	33.2	90.2	244.8
2001 .....	15.2	3.1	*	*	0.2	0.5	1.6	3.7	11.8	34.1	85.8	241.5
2000 .....	14.4	3.1	*	*	0.2	0.5	1.4	3.5	11.0	33.2	82.0	233.4
1999 .....	13.8	4.8	*	*	0.2	0.6	1.3	3.3	10.4	32.1	77.8	225.2
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	15.2	3.1	*	*	0.1	0.4	1.1	3.0	9.2	32.3	100.8	277.6
2001 .....	14.6	2.2	*	*	0.2	0.4	1.1	3.0	9.2	32.7	95.1	276.1
2000 .....	13.9	2.5	*	*	0.2	0.4	1.1	2.8	9.0	30.7	92.3	265.7
1999 .....	13.1	3.5	*	*	0.1	0.4	0.9	2.4	8.4	30.4	87.8	251.6

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Renal failure (N17-N19)</b>												
Non-Hispanic white												
Male												
2002 .....	15.0	3.9	*	*	*	0.4	1.2	3.5	10.0	38.8	128.8	377.8
2001 .....	14.5	2.3	*	*	*	0.3	1.3	3.6	10.3	39.1	121.7	387.6
2000 .....	13.7	3.3	*	*	0.2	0.4	1.2	3.3	10.3	36.5	119.4	373.4
1999 .....	13.0	3.8	*	*	*	0.4	1.1	2.8	9.6	35.9	114.4	353.8
Female												
2002 .....	15.3	2.2	*	*	*	0.4	1.0	2.5	8.5	26.7	82.0	234.3
2001 .....	14.7	2.1	*	*	0.2	0.4	0.9	2.4	8.1	27.3	77.5	229.7
2000 .....	14.0	*	*	*	*	0.3	0.9	2.3	7.7	25.8	74.4	222.2
1999 .....	13.2	3.1	*	*	*	0.4	0.8	2.1	7.3	25.8	70.3	210.8
Non-Hispanic black												
Both sexes												
2002 .....	20.2	8.0	*	*	0.6	2.0	5.4	16.0	41.4	100.9	202.7	436.5
2001 .....	20.0	8.5	*	*	0.4	1.7	5.3	15.8	42.1	102.9	201.9	428.9
2000 .....	19.2	11.4	*	*	0.4	1.8	5.3	15.6	40.7	99.9	194.6	404.2
1999 .....	18.8	10.2	*	*	0.5	1.9	5.4	15.2	37.0	95.3	193.0	428.7
Male												
2002 .....	19.4	10.0	*	*	*	1.8	6.1	20.1	49.8	115.4	233.3	562.3
2001 .....	18.4	8.9	*	*	*	2.1	5.1	19.3	45.7	114.3	233.5	514.0
2000 .....	17.9	13.6	*	*	*	1.9	6.5	19.9	45.0	104.2	231.7	487.3
1999 .....	17.7	8.7	*	*	*	1.8	6.4	19.0	40.5	102.5	246.7	519.4
Female												
2002 .....	20.9	*	*	*	*	2.1	4.8	12.5	34.7	90.6	185.4	388.4
2001 .....	21.4	8.1	*	*	*	1.4	5.5	12.8	39.2	94.9	184.1	397.4
2000 .....	20.3	9.2	*	*	*	1.7	4.1	11.8	37.3	96.9	173.7	373.3
1999 .....	19.8	11.8	*	*	*	2.0	4.5	12.0	34.3	90.2	163.0	394.9
<b>Other disorders of kidney (N25,N27)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
Hispanic												
Both sexes												
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
Non-Hispanic black												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.3
2001 .....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.0
2000 .....	0.3	*	*	*	*	*	0.0	0.1	0.2	0.6	2.0	7.5
1999 .....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.7	2.0	7.5
Male												
2002 .....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	5.3
2001 .....	0.2	*	*	*	*	*	*	*	*	0.5	1.5	4.3
2000 .....	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.6
1999 .....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.7	6.8
Female												
2002 .....	0.4	*	*	*	*	*	*	0.1	0.3	0.6	2.2	6.7
2001 .....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.5	2.3	6.7
2000 .....	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999 .....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.9	2.1	7.8
Hispanic												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
Hispanic												
Female												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	6.4
2001 .....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.1
2000 .....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.0	7.6
1999 .....	0.3	*	*	*	*	*	0.0	0.1	0.2	0.7	1.9	7.5
Male												
2002 .....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	5.3
2001 .....	0.2	*	*	*	*	*	*	*	*	0.5	1.6	4.3
2000 .....	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.8
1999 .....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	6.9
Female												
2002 .....	0.4	*	*	*	*	*	*	0.1	0.3	0.6	2.2	6.8
2001 .....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.4	2.3	6.9
2000 .....	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999 .....	0.4	*	*	*	*	*	*	0.2	0.2	0.9	2.1	7.7
Non-Hispanic white												
Both sexes												
2002 .....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.5
2001 .....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.4	2.0	6.2
2000 .....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.9	7.6
1999 .....	0.3	*	*	*	*	*	*	0.1	0.2	0.7	1.9	7.7
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.5	1.3	5.4
2001 .....	0.2	*	*	*	*	*	*	*	*	0.4	1.5	4.2
2000 .....	0.2	*	*	*	*	*	*	*	*	*	1.5	6.0
1999 .....	0.2	*	*	*	*	*	*	*	*	0.4	1.7	7.3
Female												
2002 .....	0.4	*	*	*	*	*	*	*	0.3	0.7	2.2	6.9
2001 .....	0.4	*	*	*	*	*	*	0.2	0.3	0.5	2.3	7.1
2000 .....	0.5	*	*	*	*	*	*	*	0.3	0.7	2.1	8.2
1999 .....	0.5	*	*	*	*	*	*	0.2	0.2	0.9	2.1	7.9
Non-Hispanic black												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	6.4
2001 .....	0.2	*	*	*	*	*	*	*	*	*	2.4	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	3.1	7.9
1999 .....	0.2	*	*	*	*	*	*	*	*	1.2	*	*
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Infections of kidney (N10-N12,N13.6,N15.1)</b>												
Non-Hispanic black												
Female												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.4	*	*	*	*	*	*	*	*	*	3.6	9.2
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
<b>Hyperplasia of prostate (N40)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	4.8
2001 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.5
2000 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999 .....	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.0
Male												
2002 .....	0.3	*	*	*	*	*	*	*	*	0.5	3.2	15.8
2001 .....	0.3	*	*	*	*	*	*	*	*	0.5	2.8	18.6
2000 .....	0.3	*	*	*	*	*	*	*	*	0.5	3.3	18.1
1999 .....	0.3	*	*	*	*	*	*	*	*	0.6	3.3	17.5
Female												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	4.9
2001 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.7
2000 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.1
Male												
2002 .....	0.3	*	*	*	*	*	*	*	*	0.5	3.2	16.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Hyperplasia of prostate (N40)</b>												
Non-Hispanic <sup>4</sup>												
Male												
2001 .....	0.4	*	*	*	*	*	*	*	*	0.5	2.8	19.3
2000 .....	0.3	*	*	*	*	*	*	*	*	0.4	3.3	18.0
1999 .....	0.3	*	*	*	*	*	*	*	*	0.5	3.4	18.0
Female												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white												
Both sexes												
2002 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.0
2001 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.9
2000 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999 .....	0.2	*	*	*	*	*	*	*	*	0.2	1.4	5.2
Male												
2002 .....	0.4	*	*	*	*	*	*	*	*	0.4	3.2	16.6
2001 .....	0.4	*	*	*	*	*	*	*	*	0.5	2.8	20.1
2000 .....	0.4	*	*	*	*	*	*	*	*	0.4	3.4	18.0
1999 .....	0.4	*	*	*	*	*	*	*	*	0.5	3.4	18.4
Female												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Inflammatory diseases of female pelvic organs (N70-N76)</b>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	0.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Inflammatory diseases of female pelvic organs (N70-N76)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2000 .....	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.6
1999 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
<b>Male</b>												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	1.1
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000 .....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.9
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	*
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	0.9
2000 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
<b>Male</b>												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	*

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Inflammatory diseases of female pelvic organs (N70-N76)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001 .....	0.0	*	*	*	*	*	*	*	*	*	0.3	0.9
2000 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999 .....	0.0	*	*	*	*	*	*	*	*	*	0.2	*
<b>Male</b>												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2001 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999 .....	0.1	*	*	*	*	*	*	*	*	*	0.4	*
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Female</b>												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
<b>Pregnancy, childbirth and the puerperium (O00-O99)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001 .....	0.1	...	...	*	0.3	0.5	0.2	*	*	*	*	*
2000 .....	0.1	...	...	*	0.2	0.5	0.3	*	*	*	*	*
1999 .....	0.1	...	...	*	0.3	0.5	0.3	*	*	*	*	*
<b>Male</b>												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Female</b>												
2002 .....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
2001 .....	0.3	...	...	*	0.6	1.0	0.5	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
All origins <sup>3</sup>												
Female												
2002 .....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
1999 .....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	0.2	...	...	*	*	0.4	0.4	*	*	*	*	*
2001 .....	0.2	...	...	*	0.4	0.6	*	*	*	*	*	*
2000 .....	0.2	...	...	*	0.4	0.6	0.4	*	*	*	*	*
1999 .....	0.2	...	...	*	*	0.4	0.5	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	0.4	...	...	*	*	0.8	0.8	*	*	*	*	*
2001 .....	0.5	...	...	*	0.8	1.2	*	*	*	*	*	*
2000 .....	0.5	...	...	*	0.8	1.2	0.9	*	*	*	*	*
1999 .....	0.4	...	...	*	*	0.9	1.0	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001 .....	0.1	...	...	*	0.3	0.5	0.2	*	*	*	*	*
2000 .....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999 .....	0.1	...	...	*	0.2	0.5	0.2	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	0.2	...	...	*	0.4	0.9	0.4	*	*	*	*	*
2001 .....	0.3	...	...	*	0.5	0.9	0.4	*	*	*	*	*
2000 .....	0.3	...	...	*	0.4	0.9	0.5	*	*	*	*	*
1999 .....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
2001 .....	0.1	...	...	*	0.2	0.3	0.1	*	*	*	*	*
2000 .....	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*
1999 .....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
Non-Hispanic white												
Female												
2002 .....	0.1	...	...	*	0.2	0.6	0.3	*	*	*	*	*
2001 .....	0.2	...	...	*	0.3	0.6	0.2	*	*	*	*	*
2000 .....	0.2	...	...	*	0.3	0.6	0.3	*	*	*	*	*
1999 .....	0.2	...	...	*	0.2	0.6	0.3	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	0.4	...	...	*	0.8	1.2	0.7	*	*	*	*	*
2001 .....	0.4	...	...	*	0.8	1.3	0.7	*	*	*	*	*
2000 .....	0.4	...	...	*	0.6	1.3	0.7	*	*	*	*	*
1999 .....	0.5	...	...	*	0.8	1.3	0.7	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	0.8	...	...	*	1.5	2.4	1.4	*	*	*	*	*
2001 .....	0.8	...	...	*	1.6	2.4	1.4	*	*	*	*	*
2000 .....	0.8	...	...	*	1.2	2.4	1.3	*	*	*	*	*
1999 .....	0.9	...	...	*	1.6	2.5	1.4	*	*	*	*	*
Pregnancy with abortive outcome (O00-O07)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
2001 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
2000 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
1999 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
2001 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
2000 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
1999 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	*	...	...	*	*	*	*	*	*	*	*	*
2001 .....	*	...	...	*	*	*	*	*	*	*	*	*
2000 .....	*	...	...	*	*	*	*	*	*	*	*	*
1999 .....	*	...	...	*	*	*	*	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
Hispanic												
Male												
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	*	...	...	*	*	*	*	*	*	*	*	*
2001 .....	*	...	...	*	*	*	*	*	*	*	*	*
2000 .....	*	...	...	*	*	*	*	*	*	*	*	*
1999 .....	*	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
2001 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
2000 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
1999 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
2001 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
2000 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
1999 .....	0.0	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	*	...	...	*	*	*	*	*	*	*	*	*
2001 .....	*	...	...	*	*	*	*	*	*	*	*	*
2000 .....	*	...	...	*	*	*	*	*	*	*	*	*
1999 .....	*	...	...	*	*	*	*	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	*	...	...	*	*	*	*	*	*	*	*	*
2001 .....	*	...	...	*	*	*	*	*	*	*	*	*
2000 .....	*	...	...	*	*	*	*	*	*	*	*	*
1999 .....	*	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	*	...	...	*	*	*	*	*	*	*	*	*
2001 .....	*	...	...	*	*	*	*	*	*	*	*	*
2000 .....	*	...	...	*	*	*	*	*	*	*	*	*
1999 .....	*	...	...	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
Non-Hispanic black												
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	*	...	...	*	*	*	*	*	*	*	*	*
2001 .....	*	...	...	*	*	*	*	*	*	*	*	*
2000 .....	*	...	...	*	*	*	*	*	*	*	*	*
1999 .....	*	...	...	*	*	*	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001 .....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2000 .....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999 .....	0.1	...	...	*	0.2	0.4	0.3	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2001 .....	0.3	...	...	*	0.5	0.9	0.4	*	*	*	*	*
2000 .....	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
1999 .....	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	0.2	...	...	*	*	0.4	0.4	*	*	*	*	*
2001 .....	0.2	...	...	*	0.4	0.6	*	*	*	*	*	*
2000 .....	0.2	...	...	*	0.3	0.5	0.4	*	*	*	*	*
1999 .....	0.2	...	...	*	*	0.4	0.5	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	0.4	...	...	*	*	0.8	0.8	*	*	*	*	*
2001 .....	0.5	...	...	*	0.8	1.2	*	*	*	*	*	*
2000 .....	0.5	...	...	*	0.7	1.1	0.9	*	*	*	*	*
1999 .....	0.4	...	...	*	*	0.9	1.0	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001 .....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2000 .....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999 .....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2001 .....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2000 .....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
1999 .....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
2001 .....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
2000 .....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
1999 .....	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	0.1	...	...	*	0.2	0.5	0.2	*	*	*	*	*
2001 .....	0.1	...	...	*	0.3	0.6	0.2	*	*	*	*	*
2000 .....	0.1	...	...	*	0.3	0.5	0.3	*	*	*	*	*
1999 .....	0.1	...	...	*	0.2	0.5	0.3	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	0.4	...	...	*	0.7	1.2	0.7	*	*	*	*	*
2001 .....	0.4	...	...	*	0.7	1.2	0.6	*	*	*	*	*
2000 .....	0.3	...	...	*	0.5	1.1	0.6	*	*	*	*	*
1999 .....	0.4	...	...	*	0.8	1.2	0.7	*	*	*	*	*
Male												
2002 .....	...	...	...	...	...	...	...	...	...	...	...	...
2001 .....	...	...	...	...	...	...	...	...	...	...	...	...
2000 .....	...	...	...	...	...	...	...	...	...	...	...	...
1999 .....	...	...	...	...	...	...	...	...	...	...	...	...
Female												
2002 .....	0.7	...	...	*	1.5	2.2	1.2	*	*	*	*	*
2001 .....	0.7	...	...	*	1.3	2.2	1.2	*	*	*	*	*
2000 .....	0.7	...	...	*	1.1	2.1	1.1	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
Non-Hispanic black												
Female												
1999 .....	0.8	...	...	*	1.5	2.3	1.3	*	*	*	*	*
Certain conditions originating in the perinatal period (P00-P96)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	4.9	349.7	0.4	0.1	0.1	*	*	*	*	*	*	*
2001 .....	4.9	340.5	0.5	0.1	*	*	*	*	*	*	*	*
2000 .....	5.0	365.7	0.5	0.1	*	*	*	*	*	*	*	*
1999 .....	5.1	371.0	0.6	0.1	*	*	*	*	*	*	*	*
Male												
2002 .....	5.7	385.7	0.5	*	*	*	*	*	*	*	*	*
2001 .....	5.6	376.6	0.6	0.1	*	*	*	*	*	*	*	*
2000 .....	5.8	403.9	0.7	0.1	*	*	*	*	*	*	*	*
1999 .....	5.9	408.6	0.7	0.1	*	*	*	*	*	*	*	*
Female												
2002 .....	4.2	312.0	0.3	*	*	*	*	*	*	*	*	*
2001 .....	4.2	302.6	0.4	*	*	*	*	*	*	*	*	*
2000 .....	4.3	325.6	0.4	*	*	*	*	*	*	*	*	*
1999 .....	4.4	331.7	0.5	0.1	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	6.2	284.8	*	*	*	*	*	*	*	*	*	*
2001 .....	6.0	269.6	*	*	*	*	*	*	*	*	*	*
2000 .....	6.1	274.9	*	*	*	*	*	*	*	*	*	*
1999 .....	6.3	281.4	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	6.8	313.1	*	*	*	*	*	*	*	*	*	*
2001 .....	6.7	302.0	*	*	*	*	*	*	*	*	*	*
2000 .....	6.6	299.0	*	*	*	*	*	*	*	*	*	*
1999 .....	6.8	305.4	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	5.6	255.2	*	*	*	*	*	*	*	*	*	*
2001 .....	5.3	235.9	*	*	*	*	*	*	*	*	*	*
2000 .....	5.5	249.7	*	*	*	*	*	*	*	*	*	*
1999 .....	5.8	256.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	4.7	361.1	0.4	0.1	*	*	*	*	*	*	*	*
2001 .....	4.6	353.5	0.5	0.1	*	*	*	*	*	*	*	*
2000 .....	4.7	380.5	0.5	0.1	*	*	*	*	*	*	*	*
1999 .....	4.8	383.8	0.6	0.1	*	*	*	*	*	*	*	*
Male												
2002 .....	5.4	398.6	0.5	*	*	*	*	*	*	*	*	*
2001 .....	5.4	390.1	0.6	0.1	*	*	*	*	*	*	*	*
2000 .....	5.5	421.7	0.7	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
Non-Hispanic <sup>4</sup>												
Male												
1999 .....	5.6	423.2	0.8	0.1	*	*	*	*	*	*	*	*
Female												
2002 .....	4.0	321.7	0.3	*	*	*	*	*	*	*	*	*
2001 .....	3.9	315.1	0.3	*	*	*	*	*	*	*	*	*
2000 .....	4.0	337.2	0.4	*	*	*	*	*	*	*	*	*
1999 .....	4.1	342.4	0.4	0.1	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	3.2	270.0	0.2	*	*	*	*	*	*	*	*	*
2001 .....	3.2	260.3	0.4	*	*	*	*	*	*	*	*	*
2000 .....	3.2	275.5	0.4	0.1	*	*	*	*	*	*	*	*
1999 .....	3.2	275.1	0.4	0.1	*	*	*	*	*	*	*	*
Male												
2002 .....	3.8	303.0	*	*	*	*	*	*	*	*	*	*
2001 .....	3.7	286.6	0.5	*	*	*	*	*	*	*	*	*
2000 .....	3.7	305.5	0.5	*	*	*	*	*	*	*	*	*
1999 .....	3.8	304.4	0.5	*	*	*	*	*	*	*	*	*
Female												
2002 .....	2.7	235.5	*	*	*	*	*	*	*	*	*	*
2001 .....	2.7	232.8	*	*	*	*	*	*	*	*	*	*
2000 .....	2.7	243.9	*	*	*	*	*	*	*	*	*	*
1999 .....	2.7	244.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	13.2	731.8	1.1	*	*	*	*	*	*	*	*	*
2001 .....	13.2	765.0	0.9	*	*	*	*	*	*	*	*	*
2000 .....	13.9	838.3	1.1	*	*	*	*	*	*	*	*	*
1999 .....	14.4	872.7	1.2	*	*	*	*	*	*	*	*	*
Male												
2002 .....	15.4	791.0	*	*	*	*	*	*	*	*	*	*
2001 .....	15.8	846.6	*	*	*	*	*	*	*	*	*	*
2000 .....	16.6	932.9	*	*	*	*	*	*	*	*	*	*
1999 .....	17.1	968.0	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	11.3	670.1	*	*	*	*	*	*	*	*	*	*
2001 .....	10.9	679.1	*	*	*	*	*	*	*	*	*	*
2000 .....	11.5	740.7	*	*	*	*	*	*	*	*	*	*
1999 .....	12.0	774.4	*	*	*	*	*	*	*	*	*	*
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	3.7	139.4	3.4	1.0	1.2	1.2	1.3	1.8	2.4	2.4	3.8	6.6
2001 .....	3.7	136.7	3.6	0.9	1.3	1.2	1.3	1.6	2.3	2.4	3.7	7.2
2000 .....	3.8	150.9	3.2	1.0	1.1	1.2	1.3	1.6	2.3	2.5	3.9	7.5

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
All origins <sup>3</sup>												
Both sexes												
1999 .....	3.7	144.2	3.6	1.0	1.1	1.2	1.2	1.8	2.3	2.5	4.0	7.8
Male												
2002 .....	3.9	142.6	3.5	1.0	1.4	1.4	1.3	1.9	2.5	2.6	4.0	7.0
2001 .....	3.9	143.1	3.6	1.0	1.5	1.2	1.4	1.7	2.3	2.7	4.1	7.7
2000 .....	4.0	155.4	3.3	1.0	1.3	1.4	1.4	1.9	2.3	2.7	4.0	10.6
1999 .....	3.9	147.9	4.0	1.0	1.3	1.2	1.3	1.9	2.5	2.5	4.2	8.5
Female												
2002 .....	3.5	136.1	3.3	1.0	1.1	1.0	1.2	1.6	2.3	2.3	3.8	6.5
2001 .....	3.4	129.9	3.7	0.8	1.0	1.1	1.2	1.6	2.3	2.2	3.5	7.0
2000 .....	3.5	146.2	3.1	1.0	0.9	1.0	1.1	1.4	2.4	2.4	3.9	6.3
1999 .....	3.5	140.3	3.2	1.1	0.9	1.1	1.1	1.6	2.2	2.6	3.9	7.6
Hispanic												
Both sexes												
2002 .....	4.4	150.7	3.5	1.0	1.0	0.7	0.7	1.1	1.3	*	3.8	*
2001 .....	4.4	146.2	3.4	0.8	1.1	0.7	0.8	1.1	1.3	1.9	4.3	*
2000 .....	4.6	153.3	3.8	0.8	0.8	0.7	0.8	1.1	1.8	2.0	4.7	*
1999 .....	4.4	141.8	3.5	0.9	0.9	0.9	0.7	1.4	1.6	2.9	*	*
Male												
2002 .....	4.5	153.1	3.8	0.9	1.1	0.9	*	1.3	*	*	*	*
2001 .....	4.7	153.4	3.5	1.0	1.1	0.7	1.0	1.5	*	*	*	*
2000 .....	4.7	153.3	4.0	0.8	1.1	0.7	1.0	1.4	*	*	*	*
1999 .....	4.3	138.1	4.3	0.9	0.9	0.7	*	1.6	*	*	*	*
Female												
2002 .....	4.3	148.2	3.3	1.0	0.9	*	0.8	*	*	*	*	*
2001 .....	4.1	138.8	3.3	0.6	1.1	*	*	*	*	*	*	*
2000 .....	4.4	153.3	3.5	0.7	*	0.7	*	*	*	*	*	*
1999 .....	4.5	145.8	2.7	0.9	0.9	1.0	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	3.6	134.9	3.3	1.0	1.2	1.3	1.4	1.8	2.5	2.5	3.8	6.7
2001 .....	3.5	132.8	3.7	0.9	1.3	1.3	1.4	1.7	2.4	2.5	3.7	7.2
2000 .....	3.6	148.8	3.1	1.0	1.2	1.3	1.3	1.7	2.4	2.6	3.9	7.5
1999 .....	3.6	143.1	3.6	1.1	1.2	1.2	1.3	1.8	2.4	2.5	4.0	7.9
Male												
2002 .....	3.8	138.1	3.5	1.0	1.4	1.5	1.4	2.0	2.6	2.6	3.9	7.0
2001 .....	3.7	139.4	3.6	1.0	1.6	1.4	1.5	1.7	2.3	2.7	4.0	7.6
2000 .....	3.9	154.4	3.1	1.0	1.4	1.5	1.5	1.9	2.2	2.7	3.9	10.4
1999 .....	3.8	148.4	3.9	1.1	1.4	1.3	1.4	1.9	2.6	2.4	4.2	8.5
Female												
2002 .....	3.4	131.5	3.2	1.0	1.1	1.1	1.3	1.7	2.4	2.3	3.8	6.6
2001 .....	3.3	125.8	3.7	0.9	1.0	1.2	1.3	1.7	2.4	2.2	3.5	7.0
2000 .....	3.4	142.8	3.0	1.0	1.0	1.1	1.2	1.4	2.5	2.4	3.9	6.4
1999 .....	3.4	137.4	3.3	1.1	0.9	1.1	1.2	1.6	2.2	2.6	3.9	7.6

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Non-Hispanic white												
Both sexes												
2002 .....	3.4	131.2	3.0	1.0	1.2	1.3	1.4	1.9	2.5	2.5	3.9	6.9
2001 .....	3.4	127.4	3.1	0.9	1.2	1.3	1.4	1.7	2.5	2.5	3.8	7.4
2000 .....	3.4	141.2	2.6	0.9	1.1	1.3	1.3	1.7	2.4	2.6	4.0	7.8
1999 .....	3.4	136.8	3.0	0.9	1.1	1.3	1.3	1.8	2.5	2.5	4.0	8.0
Male												
2002 .....	3.6	137.2	3.2	1.0	1.4	1.6	1.4	2.0	2.6	2.7	3.8	7.3
2001 .....	3.6	135.5	3.1	1.0	1.4	1.5	1.5	1.7	2.4	2.9	4.1	7.4
2000 .....	3.6	147.9	2.6	0.9	1.3	1.6	1.4	1.9	2.3	2.6	4.1	10.8
1999 .....	3.6	144.4	3.2	0.9	1.4	1.3	1.4	2.0	2.7	2.4	4.1	8.6
Female												
2002 .....	3.3	125.0	2.9	1.0	1.1	1.1	1.3	1.8	2.5	2.4	3.9	6.7
2001 .....	3.2	118.8	3.2	0.7	1.0	1.2	1.3	1.7	2.6	2.3	3.6	7.4
2000 .....	3.2	134.2	2.6	1.0	1.0	1.1	1.2	1.6	2.5	2.6	4.0	6.6
1999 .....	3.3	128.9	2.9	1.0	0.8	1.2	1.2	1.7	2.3	2.6	4.0	7.8
Non-Hispanic black												
Both sexes												
2002 .....	4.7	156.9	4.7	1.3	1.7	1.5	1.6	1.8	2.7	2.4	3.4	*
2001 .....	4.7	163.5	5.4	1.1	2.0	1.3	1.6	1.8	1.9	2.1	3.0	*
2000 .....	5.0	184.1	5.2	1.4	1.6	1.4	1.6	1.6	2.4	2.7	2.8	*
1999 .....	4.8	174.8	5.8	1.6	1.5	1.1	1.4	1.9	2.0	2.9	4.3	7.4
Male												
2002 .....	5.2	153.3	4.8	1.4	2.0	1.8	1.7	2.3	3.5	3.1	*	*
2001 .....	5.1	167.5	5.5	1.0	2.5	1.2	1.8	1.7	2.1	*	*	*
2000 .....	5.5	186.0	5.6	1.5	2.1	1.7	1.9	2.2	*	3.9	*	*
1999 .....	5.3	179.5	6.6	1.6	1.8	1.3	1.6	2.3	2.1	*	*	*
Female												
2002 .....	4.4	160.6	4.7	1.1	1.3	1.2	1.6	1.5	2.0	*	*	*
2001 .....	4.4	159.2	5.4	1.3	1.4	1.4	1.5	1.8	1.8	2.4	*	*
2000 .....	4.5	182.2	4.8	1.2	1.2	1.1	1.4	1.2	3.1	*	*	*
1999 .....	4.4	170.0	5.0	1.6	1.1	1.0	1.2	1.5	1.9	3.2	*	*
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	10.4	85.7	1.6	0.3	1.8	3.3	5.3	6.7	7.5	13.1	38.8	208.0
2001 .....	11.4	86.5	1.8	0.4	2.4	3.9	6.9	8.9	9.1	15.1	39.8	206.4
2000 .....	11.3	95.6	1.6	0.4	2.2	3.9	6.6	8.3	9.3	16.5	40.7	208.1
1999 .....	9.6	93.2	1.5	0.3	1.6	2.7	4.6	5.9	7.8	14.6	36.1	187.7
Male												
2002 .....	10.1	98.5	1.8	0.4	2.4	4.5	6.9	8.8	10.0	15.9	43.8	175.5
2001 .....	11.3	98.9	2.1	0.5	3.2	5.2	9.0	11.9	12.0	18.0	43.8	175.0
2000 .....	11.5	110.6	1.7	0.5	2.9	5.3	8.9	11.5	12.3	20.6	46.0	182.7
1999 .....	9.4	106.6	1.7	0.4	2.1	3.6	6.0	8.1	10.4	18.2	40.1	166.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
All origins <sup>3</sup>												
Female												
2002 .....	10.7	72.3	1.4	0.3	1.2	2.1	3.8	4.7	5.3	10.8	35.4	222.1
2001 .....	11.4	73.6	1.5	0.4	1.5	2.5	4.9	6.1	6.3	12.7	37.2	219.5
2000 .....	11.2	79.9	1.5	0.4	1.5	2.5	4.4	5.3	6.5	13.2	37.2	218.5
1999 .....	9.7	79.2	1.3	0.3	1.0	1.9	3.2	3.8	5.5	11.7	33.5	196.4
Hispanic												
Both sexes												
2002 .....	4.0	56.6	1.3	*	1.3	2.0	3.0	3.9	4.8	7.0	18.7	91.5
2001 .....	5.9	60.7	1.6	0.5	2.8	4.2	6.0	8.3	6.9	9.9	19.3	108.2
2000 .....	5.9	65.6	1.7	0.5	2.1	4.0	6.7	8.0	6.7	9.9	20.5	102.2
1999 .....	3.8	54.9	1.0	*	1.3	2.2	3.0	3.1	3.8	7.0	19.4	87.0
Male												
2002 .....	4.6	66.4	1.6	*	1.7	2.9	4.0	4.7	6.3	9.6	23.4	88.7
2001 .....	7.4	67.2	1.8	*	3.9	6.0	8.5	12.6	10.6	14.2	25.0	87.6
2000 .....	7.5	78.6	2.1	*	2.9	5.8	9.9	12.4	9.0	14.3	25.6	106.8
1999 .....	4.4	61.3	*	*	1.9	2.9	4.3	4.7	5.5	8.3	24.0	77.6
Female												
2002 .....	3.3	46.4	*	*	0.9	1.0	2.0	3.2	3.5	5.0	15.5	92.8
2001 .....	4.3	53.9	*	0.6	1.5	2.1	3.4	4.1	3.6	6.5	15.4	118.2
2000 .....	4.1	52.1	1.4	*	1.3	2.0	3.2	3.7	4.6	6.5	17.2	99.9
1999 .....	3.1	48.2	*	*	*	1.3	1.6	1.7	2.4	6.0	16.3	91.5
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	11.3	92.8	1.7	0.4	1.9	3.6	5.6	6.9	7.7	13.5	39.6	211.9
2001 .....	12.1	92.6	1.8	0.4	2.3	3.7	7.0	8.9	9.2	15.4	40.7	209.6
2000 .....	12.0	102.4	1.6	0.4	2.2	3.8	6.5	8.2	9.3	16.8	41.4	211.6
1999 .....	10.3	101.9	1.6	0.4	1.6	2.8	4.7	6.0	8.0	15.0	36.7	190.9
Male												
2002 .....	10.9	106.2	1.9	0.4	2.5	4.8	7.3	9.1	10.1	16.3	44.6	178.6
2001 .....	11.8	106.2	2.1	0.5	3.0	4.9	9.0	11.7	12.0	18.2	44.6	178.2
2000 .....	11.9	118.0	1.7	0.5	2.9	5.1	8.6	11.2	12.3	20.7	46.6	185.4
1999 .....	10.0	116.7	1.9	0.4	2.2	3.7	6.1	8.2	10.6	18.7	40.6	169.5
Female												
2002 .....	11.7	78.7	1.4	0.3	1.3	2.3	4.0	4.8	5.4	11.2	36.3	226.3
2001 .....	12.3	78.3	1.5	0.4	1.5	2.6	5.1	6.2	6.6	13.1	38.0	222.7
2000 .....	12.1	86.1	1.5	0.4	1.5	2.6	4.5	5.4	6.6	13.6	38.0	222.3
1999 .....	10.5	86.3	1.4	0.3	1.0	2.0	3.3	3.9	5.7	11.9	34.2	199.5
Non-Hispanic white												
Both sexes												
2002 .....	11.6	76.0	1.3	0.3	1.8	3.4	5.4	6.1	7.1	13.1	39.4	215.8
2001 .....	12.4	73.1	1.5	0.3	2.2	3.6	6.9	8.5	8.6	14.7	40.4	213.3
2000 .....	12.2	79.7	1.4	0.3	2.1	3.6	6.3	7.6	8.6	15.9	40.9	215.8
1999 .....	10.2	79.3	1.4	0.3	1.4	2.5	4.1	5.2	7.3	14.3	35.7	193.8
Male												
2002 .....	10.7	88.2	1.4	0.3	2.5	4.7	6.9	7.9	9.3	15.7	43.6	181.5
2001 .....	11.8	85.2	1.7	0.4	2.9	4.9	8.8	11.1	10.9	17.3	44.7	180.4

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Non-Hispanic white												
Male												
2002 .....	11.8	91.8	1.7	0.4	2.8	4.8	8.5	10.3	11.1	19.3	46.0	189.6
1999 .....	9.6	92.3	1.7	0.3	1.9	3.3	5.5	6.9	9.5	17.7	39.2	171.1
Female												
2002 .....	12.5	63.2	1.3	0.2	1.1	2.2	3.8	4.3	5.1	10.9	36.6	230.6
2001 .....	13.1	60.4	1.3	0.3	1.5	2.4	5.0	5.9	6.4	12.5	37.6	227.0
2000 .....	12.6	66.9	1.2	0.3	1.3	2.3	4.1	4.9	6.2	13.0	37.5	226.4
1999 .....	10.9	65.5	1.0	0.3	0.9	1.8	2.8	3.5	5.2	11.4	33.4	202.8
Non-Hispanic black												
Both sexes												
2002 .....	12.6	165.4	2.7	0.8	2.6	5.3	8.8	13.9	13.9	20.4	49.9	198.1
2001 .....	12.9	176.9	3.0	0.6	2.7	5.2	9.3	14.0	15.1	24.6	50.3	194.2
2000 .....	13.9	205.6	2.3	0.7	3.2	5.9	9.3	14.8	17.3	29.2	56.1	189.2
1999 .....	13.0	202.0	2.9	0.5	2.7	5.0	9.1	13.2	16.0	24.6	54.8	182.0
Male												
2002 .....	14.4	186.3	3.6	0.9	3.1	7.1	11.7	19.6	20.0	26.4	68.6	182.7
2001 .....	14.2	195.9	3.7	0.8	3.8	6.1	11.4	18.2	23.8	30.7	52.2	186.0
2000 .....	15.7	236.8	2.2	0.9	3.8	7.8	10.9	20.6	25.1	40.0	66.0	163.2
1999 .....	14.9	228.0	2.9	0.6	3.6	6.9	11.6	19.4	23.0	32.8	66.9	180.7
Female												
2002 .....	10.9	143.6	1.9	0.7	2.0	3.7	6.2	9.0	8.9	16.1	39.3	204.0
2001 .....	11.7	156.8	2.2	*	1.7	4.3	7.4	10.3	8.1	20.3	49.2	197.2
2000 .....	12.4	173.4	2.4	*	2.6	4.3	7.8	9.9	11.0	21.6	50.5	198.9
1999 .....	11.2	175.3	2.8	*	1.9	3.2	7.0	7.9	10.3	18.8	48.0	182.6
All other diseases (Residual)												
(D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	67.5	28.4	4.2	2.1	4.6	7.9	18.9	37.8	65.3	140.6	405.3	1,493.0
2001 .....	65.5	28.0	4.2	2.1	4.5	7.8	18.0	36.2	63.4	140.9	395.3	1,476.7
2000 .....	63.4	29.0	4.0	2.1	4.5	7.8	18.0	35.4	62.4	136.0	388.5	1,445.7
1999 .....	61.5	30.2	4.2	2.3	4.6	7.6	17.7	33.9	60.5	136.6	373.0	1,410.7
Male												
2002 .....	55.8	33.4	4.6	2.3	5.1	9.1	22.4	46.3	74.5	152.6	406.7	1,262.9
2001 .....	53.8	31.4	4.6	2.2	5.2	8.6	21.3	43.7	71.2	153.4	397.4	1,271.6
2000 .....	52.3	32.3	4.3	2.3	5.3	9.0	21.4	42.8	70.7	148.4	391.5	1,266.2
1999 .....	51.0	35.0	4.5	2.3	5.3	8.8	21.6	40.5	69.0	148.7	384.2	1,225.4
Female												
2002 .....	78.8	23.0	3.9	1.9	4.0	6.6	15.5	29.6	56.8	130.6	404.4	1,592.8
2001 .....	76.7	24.5	3.8	2.0	3.8	6.9	14.6	29.1	56.2	130.5	394.0	1,562.6
2000 .....	74.0	25.6	3.6	2.0	3.7	6.5	14.6	28.2	54.8	125.8	386.5	1,518.8
1999 .....	71.6	25.3	3.9	2.2	3.8	6.4	13.9	27.4	52.7	126.7	365.7	1,485.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual)												
(D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
Hispanic												
Both sexes												
2002 .....	21.6	25.1	3.8	2.1	3.7	5.6	13.0	27.5	48.5	111.2	272.2	870.8
2001 .....	21.2	22.9	3.7	1.7	3.7	5.2	12.3	26.3	49.7	107.2	277.5	991.0
2000 .....	20.7	25.4	3.7	1.7	3.3	5.9	13.8	27.5	47.5	103.4	284.0	913.0
1999 .....	21.5	26.5	3.7	1.6	3.5	6.5	14.0	28.2	51.4	109.3	263.5	1,039.7
Male												
2002 .....	21.1	27.2	4.1	2.4	4.0	6.6	17.1	35.5	56.4	126.0	290.1	808.2
2001 .....	20.2	25.4	4.3	1.8	4.1	6.0	15.9	34.7	59.1	117.2	273.7	939.0
2000 .....	20.2	25.3	3.6	1.8	3.9	7.0	18.2	35.6	57.7	119.9	273.9	870.7
1999 .....	21.0	32.1	3.4	1.6	3.8	8.1	18.8	37.0	62.6	123.9	276.3	923.1
Female												
2002 .....	22.1	22.8	3.5	1.7	3.2	4.4	8.6	19.6	41.4	99.5	260.0	901.9
2001 .....	22.3	20.2	3.1	1.7	3.2	4.2	8.4	18.0	41.4	99.4	280.2	1,016.5
2000 .....	21.4	25.5	3.8	1.5	2.5	4.5	9.0	19.6	38.5	90.4	290.9	933.8
1999 .....	22.0	20.6	4.0	1.7	3.1	4.6	8.8	19.7	41.6	97.9	255.0	1,096.0
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	74.4	29.0	4.3	2.1	4.7	8.3	19.7	38.6	66.4	142.2	410.7	1,514.9
2001 .....	71.8	29.0	4.3	2.2	4.6	8.3	18.6	37.0	64.2	142.7	399.8	1,491.3
2000 .....	69.3	29.6	4.0	2.2	4.7	8.1	18.4	35.8	63.3	137.5	391.9	1,462.0
1999 .....	66.8	30.8	4.3	2.4	4.8	7.8	18.1	34.1	60.9	137.8	376.6	1,420.1
Male												
2002 .....	61.2	34.9	4.6	2.3	5.4	9.7	23.1	47.1	75.6	153.8	411.5	1,280.6
2001 .....	58.8	32.7	4.6	2.2	5.4	9.2	21.9	44.2	71.7	155.2	402.2	1,282.3
2000 .....	56.9	33.6	4.4	2.3	5.5	9.4	21.7	43.0	71.3	149.4	395.5	1,279.8
1999 .....	55.2	35.1	4.7	2.4	5.6	9.0	21.7	40.5	69.1	149.6	387.8	1,234.0
Female												
2002 .....	86.8	22.7	4.0	1.9	4.1	7.0	16.4	30.4	57.9	132.5	410.1	1,615.9
2001 .....	84.2	25.2	4.0	2.1	3.9	7.4	15.4	30.0	57.3	132.2	398.3	1,578.3
2000 .....	81.0	25.4	3.5	2.1	3.9	6.8	15.2	28.9	55.9	127.6	389.5	1,535.7
1999 .....	77.9	26.2	3.9	2.3	4.0	6.7	14.4	28.0	53.3	128.2	369.3	1,494.7
Non-Hispanic white												
Both sexes												
2002 .....	80.5	24.7	3.8	2.0	4.3	7.4	17.8	35.3	62.7	139.1	410.5	1,540.5
2001 .....	77.3	23.2	3.9	2.1	4.3	7.2	16.4	33.2	60.1	139.7	399.5	1,512.9
2000 .....	74.3	24.0	3.4	2.1	4.5	7.0	16.2	32.0	59.0	133.9	390.1	1,481.3
1999 .....	71.5	24.2	4.0	2.2	4.5	7.0	15.5	30.1	56.9	133.5	375.1	1,443.8
Male												
2002 .....	64.8	31.4	4.0	2.2	5.2	8.9	21.8	43.5	71.1	148.8	408.2	1,293.4
2001 .....	61.9	25.3	4.4	2.2	5.3	8.2	19.8	40.0	66.9	150.8	399.4	1,293.1
2000 .....	59.6	26.7	3.8	2.3	5.4	8.5	19.4	39.2	66.0	144.1	390.9	1,288.3
1999 .....	57.6	27.4	4.5	2.2	5.4	8.4	19.4	36.2	64.5	144.2	382.8	1,239.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>All other diseases (Residual)</b> (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
<b>Non-Hispanic white</b>												
<b>Female</b>												
2002 .....	95.5	17.7	3.6	1.8	3.4	5.8	13.9	27.2	54.8	130.8	412.1	1,647.2
2001 .....	92.1	21.0	3.4	2.1	3.3	6.2	13.1	26.4	53.7	130.2	399.5	1,604.4
2000 .....	88.3	21.2	2.9	1.8	3.5	5.5	12.9	25.0	52.5	125.2	389.6	1,559.1
1999 .....	84.9	20.9	3.5	2.2	3.5	5.6	11.7	24.1	49.8	124.5	370.2	1,525.2
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	61.7	46.1	6.1	2.7	7.0	15.0	34.7	69.9	110.2	197.3	482.8	1,443.4
2001 .....	61.1	53.8	5.6	2.5	6.7	14.7	34.8	71.7	109.1	198.5	476.3	1,431.5
2000 .....	60.1	51.3	6.1	2.9	6.6	15.6	35.1	70.6	109.6	195.0	472.0	1,403.3
1999 .....	58.0	58.4	5.5	3.1	6.9	13.5	36.2	69.3	104.9	202.1	444.8	1,295.3
<b>Male</b>												
2002 .....	56.6	50.6	6.6	2.9	6.9	15.8	36.2	83.4	131.4	232.9	526.1	1,331.3
2001 .....	56.4	61.3	5.2	2.5	6.8	15.6	39.3	85.6	128.3	232.9	530.2	1,329.4
2000 .....	56.1	61.4	6.4	2.6	7.1	16.3	39.6	82.4	133.0	231.8	527.9	1,340.9
1999 .....	54.4	66.5	5.6	3.6	7.3	13.9	40.3	81.9	122.9	228.2	509.7	1,303.8
<b>Female</b>												
2002 .....	66.3	41.3	5.6	2.4	7.2	14.3	33.3	58.3	93.2	172.0	458.3	1,486.2
2001 .....	65.2	46.0	6.0	2.6	6.6	13.9	30.8	59.7	93.6	174.3	446.1	1,469.3
2000 .....	63.7	40.9	5.8	3.2	6.2	15.0	31.0	60.5	90.8	169.1	440.6	1,426.5
1999 .....	61.3	50.0	5.4	2.6	6.5	13.1	32.7	58.5	90.5	183.7	408.5	1,292.1
<b>Accidents (unintentional injuries) (V01-X59,Y85-Y86)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	37.0	23.5	10.5	6.6	38.0	31.5	37.2	36.6	31.4	44.2	101.3	275.4
2001 .....	35.7	24.2	11.2	6.9	36.1	29.9	35.4	34.1	30.3	42.8	100.9	276.4
2000 .....	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5
1999 .....	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4
<b>Male</b>												
2002 .....	48.9	26.8	12.9	8.2	54.9	47.7	53.7	53.3	45.2	60.6	131.6	346.6
2001 .....	47.2	26.2	13.3	8.6	52.9	45.4	51.8	50.6	43.0	57.3	133.4	358.5
2000 .....	46.2	25.7	14.0	8.8	52.1	45.0	51.0	49.0	44.1	55.9	124.3	359.7
1999 .....	46.4	25.4	14.4	9.4	50.7	45.0	50.2	47.9	44.0	60.2	132.4	375.4
<b>Female</b>												
2002 .....	25.6	20.0	8.1	5.0	20.1	14.9	20.8	20.5	18.6	30.6	81.2	244.6
2001 .....	24.5	22.1	8.9	5.2	18.4	14.0	19.2	18.0	18.5	30.8	79.6	242.0
2000 .....	23.8	20.5	9.7	5.6	19.1	13.7	17.5	16.8	18.8	30.3	76.1	238.4
1999 .....	24.1	19.0	10.3	5.6	19.2	14.0	17.6	16.3	18.3	31.8	79.7	244.8
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	26.1	15.2	10.9	5.4	35.5	27.6	31.7	32.0	28.7	39.5	77.4	153.0
2001 .....	25.8	18.8	10.3	5.4	32.6	28.2	32.3	31.7	30.5	40.0	73.8	166.3

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Accidents (unintentional injuries) (V01-X59,Y85-Y86)</b>												
<b>Hispanic</b>												
<b>Both sexes</b>												
2000 .....	25.0	15.3	10.4	5.6	31.8	27.3	31.5	30.4	28.8	38.8	72.1	184.5
1999 .....	25.5	17.7	11.9	5.6	30.8	28.4	32.6	31.2	28.5	42.5	69.6	182.4
<b>Male</b>												
2002 .....	38.5	19.5	13.5	6.6	53.5	43.0	48.5	48.7	43.5	58.7	105.9	224.2
2001 .....	37.6	18.0	12.7	6.2	49.2	43.8	50.1	48.8	45.0	54.9	105.1	198.1
2000 .....	36.9	17.0	11.6	6.1	48.2	42.9	49.6	48.4	43.2	57.2	93.9	251.9
1999 .....	37.7	20.7	13.5	6.8	46.2	44.1	51.3	50.2	43.5	63.3	94.9	258.8
<b>Female</b>												
2002 .....	12.8	10.8	8.2	4.1	14.5	9.5	13.4	15.3	15.3	24.2	57.9	117.7
2001 .....	13.2	19.7	7.8	4.5	13.3	10.1	13.0	14.9	17.7	28.2	52.6	150.8
2000 .....	12.4	13.5	9.2	5.2	12.5	9.2	12.2	12.8	16.0	24.4	57.4	151.3
1999 .....	12.6	14.6	10.3	4.2	12.8	10.1	12.6	12.8	15.2	26.2	52.7	145.6
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	38.5	25.5	10.4	6.9	38.3	32.2	37.8	36.8	31.4	44.4	102.1	279.8
2001 .....	36.9	25.4	11.3	7.2	36.6	30.1	35.6	34.0	30.0	42.8	101.9	279.8
2000 .....	36.0	25.0	12.2	7.5	36.7	29.8	34.2	32.5	30.9	41.9	95.8	276.1
1999 .....	36.2	23.4	12.4	7.9	36.0	29.6	33.7	31.6	30.5	44.5	101.3	285.0
<b>Male</b>												
2002 .....	50.3	28.6	12.7	8.4	55.0	48.6	54.2	53.4	44.9	60.3	132.3	351.2
2001 .....	48.5	28.0	13.4	9.0	53.5	45.6	51.7	50.4	42.5	57.1	134.3	364.5
2000 .....	47.3	27.7	14.5	9.4	52.7	45.2	50.8	48.6	43.8	55.5	125.2	363.1
1999 .....	47.4	26.6	14.5	9.9	51.4	44.8	49.7	47.3	43.5	59.7	133.2	378.7
<b>Female</b>												
2002 .....	27.3	22.3	8.1	5.2	21.1	15.9	21.8	20.9	18.8	31.0	82.2	249.0
2001 .....	26.0	22.6	9.2	5.3	19.3	14.7	19.9	18.3	18.5	30.8	80.6	244.6
2000 .....	25.2	22.2	9.7	5.6	20.3	14.5	18.1	17.0	18.9	30.6	76.7	240.9
1999 .....	25.5	20.0	10.2	5.9	20.2	14.6	18.0	16.5	18.5	32.0	80.6	247.5
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	40.6	19.8	9.4	6.3	41.6	33.7	38.7	36.3	30.6	43.9	104.4	290.9
2001 .....	38.5	20.9	10.9	6.6	39.0	31.1	35.9	32.8	29.0	42.4	103.7	289.3
2000 .....	37.3	20.1	11.1	7.2	39.2	30.0	34.2	31.0	30.0	41.0	97.2	285.6
1999 .....	37.2	19.0	11.1	7.4	38.1	29.9	33.5	30.0	28.9	43.3	101.7	293.3
<b>Male</b>												
2002 .....	52.0	23.3	11.1	7.7	59.2	50.3	55.0	52.1	43.0	58.8	133.5	361.6
2001 .....	49.7	22.9	13.2	8.1	56.6	46.7	51.8	48.0	40.3	56.1	135.2	374.7
2000 .....	48.1	22.9	13.4	8.9	55.9	45.2	50.4	45.8	41.6	53.6	125.2	373.8
1999 .....	47.8	22.0	12.9	9.1	53.6	44.9	49.0	44.2	40.7	57.3	132.7	387.1
<b>Female</b>												
2002 .....	29.6	16.0	7.5	4.8	23.2	16.8	22.4	20.6	18.8	31.2	84.9	260.3
2001 .....	27.8	18.7	8.5	5.0	20.7	15.2	19.9	17.9	18.3	30.7	82.8	253.8
2000 .....	26.9	17.1	8.7	5.3	21.9	14.6	18.0	16.3	19.1	30.4	78.8	250.0
1999 .....	27.0	15.9	9.3	5.6	22.0	14.7	17.9	15.9	17.7	31.5	81.5	255.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Non-Hispanic black												
Both sexes												
2002 .....	34.0	50.1	14.6	9.4	28.4	32.5	39.9	47.6	41.9	49.7	84.6	175.4
2001 .....	34.4	44.5	14.2	9.7	29.6	31.1	40.9	48.5	42.4	51.3	89.6	183.9
2000 .....	34.4	46.5	16.8	9.6	29.6	33.5	40.6	47.8	41.3	53.0	85.3	183.2
1999 .....	36.1	44.1	18.2	10.9	30.5	33.2	40.7	48.3	45.1	58.6	103.8	208.9
Male												
2002 .....	49.0	52.1	18.9	11.9	42.6	52.2	59.7	73.3	69.2	77.2	131.3	262.5
2001 .....	49.5	50.4	15.8	13.1	44.5	48.8	61.3	78.6	68.2	74.9	138.3	256.4
2000 .....	50.3	49.5	19.4	12.3	44.2	53.5	63.3	76.7	69.4	80.4	133.3	277.1
1999 .....	52.3	48.5	21.8	14.3	46.4	53.0	63.0	79.4	71.8	89.6	151.0	318.5
Female												
2002 .....	20.3	48.0	10.1	6.8	14.2	14.7	22.5	25.6	20.0	30.2	58.2	142.1
2001 .....	20.6	38.3	12.4	6.2	14.8	15.1	22.7	22.7	21.6	34.6	62.2	157.0
2000 .....	20.1	43.3	14.2	6.7	15.1	15.5	20.5	23.0	18.7	33.7	58.4	148.3
1999 .....	21.4	39.6	14.5	7.4	14.8	15.4	21.1	21.8	23.7	36.8	77.4	167.9
Transport accidents (V01-V99,Y85)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	16.8	3.1	4.1	4.2	29.1	18.7	17.1	16.4	15.7	18.3	27.0	28.8
2001 .....	16.6	3.7	4.4	4.4	27.9	18.8	17.4	16.0	15.3	17.9	27.9	28.9
2000 .....	16.6	4.6	4.5	4.7	28.1	18.6	17.0	15.8	16.0	18.0	27.0	31.5
1999 .....	16.6	5.0	4.6	4.8	27.6	18.5	16.8	15.4	16.0	19.4	28.4	32.0
Male												
2002 .....	23.7	3.5	4.8	4.9	40.7	28.1	24.6	24.0	22.5	25.3	37.7	53.0
2001 .....	23.5	3.3	5.0	5.3	39.6	28.3	25.1	23.3	21.6	24.3	39.8	52.7
2000 .....	23.3	4.8	5.0	5.4	39.3	27.7	24.8	23.0	22.5	24.3	38.1	59.8
1999 .....	23.1	5.2	5.1	5.8	38.1	27.2	24.0	22.4	22.5	26.1	40.7	63.5
Female												
2002 .....	10.1	2.8	3.4	3.5	16.9	9.2	9.7	9.0	9.3	12.5	19.8	18.3
2001 .....	10.0	4.2	3.8	3.5	15.6	9.2	9.8	8.9	9.4	12.7	20.1	18.9
2000 .....	10.2	4.3	4.0	3.9	16.4	9.3	9.3	8.9	10.0	12.8	19.8	19.9
1999 .....	10.4	4.8	4.2	3.9	16.6	9.6	9.6	8.7	10.1	14.0	20.4	19.3
Hispanic												
Both sexes												
2002 .....	15.3	3.1	5.8	3.8	27.6	18.3	16.0	15.2	15.2	18.4	26.2	25.5
2001 .....	15.6	5.3	5.3	3.8	26.0	18.8	17.3	16.4	17.9	20.2	30.1	23.8
2000 .....	14.8	5.4	5.2	4.1	24.6	17.3	16.2	16.2	15.7	20.4	23.9	25.2
1999 .....	14.4	5.3	5.5	3.8	23.1	17.8	15.4	15.0	15.5	22.0	25.0	25.3
Male												
2002 .....	22.4	*	7.1	4.6	40.4	27.5	23.9	21.6	22.1	28.5	37.3	45.2
2001 .....	22.8	*	6.0	4.3	38.4	28.8	25.9	23.2	24.6	28.2	51.0	43.8
2000 .....	21.6	6.1	5.6	4.5	36.1	26.4	24.1	24.1	23.5	29.5	36.4	44.3
1999 .....	20.8	5.7	5.8	4.5	33.3	26.7	23.2	22.6	21.0	31.6	40.1	*
Female												
2002 .....	7.7	*	4.5	2.9	12.6	7.3	7.5	8.7	8.9	10.3	18.5	*
2001 .....	8.0	6.7	4.6	3.2	11.6	7.1	8.0	9.7	12.0	13.9	16.0	*

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Hispanic												
Female												
2000 .....	7.6	*	4.8	3.7	11.0	6.8	7.8	8.5	8.7	13.3	15.5	*
1999 .....	7.6	*	5.2	3.1	11.0	7.6	7.0	7.7	10.6	14.5	14.9	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	16.9	3.2	3.7	4.3	29.3	18.8	17.2	16.4	15.6	18.3	26.9	28.9
2001 .....	16.7	3.4	4.2	4.6	28.2	18.8	17.3	15.8	15.0	17.7	27.7	29.0
2000 .....	16.8	4.3	4.3	4.8	28.8	18.7	17.0	15.7	15.9	17.8	27.1	31.7
1999 .....	16.9	4.9	4.4	5.0	28.4	18.5	16.9	15.4	16.0	19.2	28.3	32.2
Male												
2002 .....	23.8	3.4	4.2	5.0	40.6	28.1	24.6	24.1	22.4	24.9	37.5	53.2
2001 .....	23.5	3.2	4.7	5.5	39.7	28.1	24.8	23.2	21.3	24.0	39.2	52.9
2000 .....	23.4	4.5	4.8	5.6	39.8	27.8	24.7	22.8	22.2	23.9	38.0	60.4
1999 .....	23.3	5.1	4.9	6.0	39.0	27.2	24.0	22.2	22.4	25.6	40.4	64.3
Female												
2002 .....	10.4	2.9	3.1	3.6	17.7	9.5	10.0	9.0	9.3	12.7	19.8	18.4
2001 .....	10.2	3.6	3.6	3.6	16.3	9.5	10.0	8.8	9.1	12.5	20.2	19.1
2000 .....	10.5	4.2	3.8	3.9	17.4	9.7	9.4	8.9	10.1	12.8	20.0	20.1
1999 .....	10.7	4.7	3.9	4.0	17.5	10.0	9.9	8.8	10.0	13.9	20.5	19.3
Non-Hispanic white												
Both sexes												
2002 .....	17.5	2.5	3.3	4.3	31.8	19.0	17.3	16.4	15.5	18.3	27.4	29.7
2001 .....	17.0	3.1	3.8	4.4	29.9	19.0	17.4	15.6	14.7	17.6	28.3	29.9
2000 .....	17.1	3.5	3.8	4.8	30.9	18.5	17.0	15.4	15.8	17.6	27.6	32.4
1999 .....	17.1	4.3	3.7	5.0	30.3	18.6	16.7	15.0	15.6	19.1	28.6	32.5
Male												
2002 .....	24.3	2.9	3.6	5.0	43.6	28.1	24.6	24.0	22.0	24.5	37.5	54.0
2001 .....	23.8	2.9	4.4	5.2	41.9	28.2	24.6	22.7	20.8	23.7	39.6	54.3
2000 .....	23.7	3.8	4.3	5.7	42.5	27.1	24.5	22.2	21.6	23.2	38.1	62.3
1999 .....	23.5	4.9	4.0	5.9	41.2	26.9	23.6	21.6	21.7	25.2	40.3	64.6
Female												
2002 .....	10.9	2.1	2.9	3.5	19.4	9.8	10.1	8.9	9.4	13.0	20.7	19.2
2001 .....	10.6	3.3	3.2	3.5	17.4	9.7	10.2	8.6	9.1	12.5	20.8	19.8
2000 .....	10.9	3.2	3.4	4.0	18.9	9.7	9.5	8.6	10.4	12.8	20.7	20.4
1999 .....	11.0	3.6	3.4	4.0	19.1	10.2	9.9	8.6	9.9	13.9	20.9	19.7
Non-Hispanic black												
Both sexes												
2002 .....	15.7	5.6	5.0	4.7	21.6	21.1	18.5	18.0	17.3	17.3	21.8	18.8
2001 .....	16.3	*	5.4	5.5	22.8	20.5	19.2	19.1	18.4	18.8	21.4	17.8
2000 .....	16.4	6.9	5.9	4.8	22.4	22.2	19.1	19.7	17.5	19.9	19.9	21.9
1999 .....	16.7	7.7	7.1	5.6	22.7	20.6	19.5	19.2	18.5	19.5	26.0	27.8
Male												
2002 .....	23.6	*	6.5	5.4	31.0	34.1	28.0	28.0	28.6	28.7	40.7	47.0
2001 .....	24.5	*	5.8	7.1	33.4	32.6	29.7	30.7	30.2	27.4	36.9	35.8
2000 .....	24.8	7.2	6.4	5.8	32.5	35.6	30.5	30.1	29.6	32.6	36.4	45.8
1999 .....	24.9	*	8.0	7.0	32.9	33.0	29.8	30.7	29.3	29.5	43.5	68.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Non-Hispanic black												
Female												
2002 .....	8.5	*	3.6	3.9	12.1	9.3	10.1	9.4	8.2	9.1	11.1	*
2001 .....	8.9	*	5.0	3.8	12.3	9.6	9.9	9.2	9.0	12.8	12.6	11.1
2000 .....	8.8	*	5.4	3.7	12.5	10.1	8.9	10.8	7.8	11.0	10.6	13.1
1999 .....	9.4	9.3	6.2	4.1	12.6	9.4	10.4	9.5	9.8	12.5	16.3	12.4
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	15.7	3.0	3.9	3.9	28.2	17.8	15.8	14.8	14.1	17.0	25.7	28.0
2001 .....	15.4	3.6	4.1	4.1	26.8	17.5	15.7	14.2	13.7	16.3	26.3	27.8
2000 .....	15.4	4.4	4.2	4.3	26.9	17.3	15.3	14.2	14.4	16.5	25.7	30.4
1999 .....	15.2	4.8	4.2	4.3	26.2	16.9	14.9	13.6	14.2	17.8	26.5	30.4
Male												
2002 .....	21.9	3.3	4.5	4.6	39.3	26.5	22.3	21.3	19.9	22.7	35.3	51.7
2001 .....	21.5	3.2	4.5	4.9	38.0	26.2	22.4	20.5	19.0	21.5	37.3	51.1
2000 .....	21.3	4.6	4.7	5.0	37.4	25.5	22.0	20.4	19.8	21.7	35.6	57.5
1999 .....	20.9	5.0	4.7	5.1	36.0	24.7	21.1	19.4	19.3	23.3	38.0	59.8
Female												
2002 .....	9.8	2.8	3.3	3.3	16.6	8.8	9.3	8.6	8.9	12.3	19.3	17.7
2001 .....	9.5	4.0	3.6	3.3	15.1	8.7	9.1	8.1	8.8	12.0	19.2	18.0
2000 .....	9.7	4.2	3.8	3.6	15.9	8.8	8.8	8.2	9.5	12.3	19.2	19.3
1999 .....	9.7	4.7	3.8	3.6	15.9	8.9	8.8	8.0	9.5	13.3	19.1	18.5
Hispanic												
Both sexes												
2002 .....	14.6	3.1	5.6	3.6	26.6	17.6	15.2	13.9	14.3	17.9	24.4	24.9
2001 .....	14.3	4.9	4.7	3.5	25.1	17.4	15.3	13.8	15.1	17.7	26.3	20.0
2000 .....	13.9	5.2	4.9	3.8	23.7	16.4	14.7	15.1	14.4	18.4	22.3	22.6
1999 .....	13.3	5.3	5.2	3.6	21.9	16.4	13.9	13.7	14.5	19.7	23.5	23.2
Male												
2002 .....	21.3	*	6.8	4.4	38.9	26.4	22.6	19.6	20.5	27.5	34.5	43.5
2001 .....	21.0	*	5.3	4.0	36.9	26.8	23.1	20.2	21.2	24.6	44.3	*
2000 .....	20.1	6.1	5.2	4.1	34.7	24.9	21.6	22.0	21.2	26.0	33.4	*
1999 .....	19.1	5.7	5.6	4.1	31.6	24.4	20.8	20.1	19.0	27.2	37.0	*
Female												
2002 .....	7.4	*	4.4	2.7	12.4	7.2	7.3	8.2	8.7	10.2	17.4	*
2001 .....	7.2	6.2	4.1	3.0	11.2	6.5	6.9	7.6	9.7	12.3	14.2	*
2000 .....	7.2	*	4.6	3.6	10.6	6.5	7.3	8.3	8.4	12.4	14.9	*
1999 .....	7.2	*	4.7	3.0	10.6	7.1	6.5	7.4	10.6	13.8	14.6	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	15.9	3.0	3.5	4.0	28.5	17.7	15.8	14.8	14.1	16.9	25.7	28.1
2001 .....	15.5	3.2	3.9	4.2	27.1	17.5	15.7	14.2	13.5	16.2	26.3	28.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2000 .....	15.6	4.2	4.1	4.4	27.5	17.4	15.4	14.1	14.4	16.3	25.8	30.6
1999 .....	15.4	4.7	4.0	4.5	26.9	16.9	15.0	13.5	14.0	17.6	26.5	30.5
Male												
2002 .....	21.9	3.2	4.0	4.6	39.2	26.4	22.2	21.3	19.7	22.3	35.1	51.9
2001 .....	21.5	3.1	4.3	5.1	38.1	25.9	22.2	20.4	18.8	21.3	36.9	51.6
2000 .....	21.4	4.2	4.5	5.1	37.9	25.5	22.0	20.2	19.6	21.3	35.7	58.3
1999 .....	21.0	4.8	4.4	5.2	36.7	24.6	21.1	19.2	19.1	22.9	37.7	60.7
Female												
2002 .....	10.1	2.9	3.0	3.4	17.4	9.1	9.5	8.6	8.9	12.4	19.4	17.8
2001 .....	9.8	3.4	3.5	3.3	15.8	9.1	9.4	8.2	8.7	12.0	19.3	18.2
2000 .....	10.0	4.2	3.6	3.6	16.9	9.2	8.9	8.2	9.6	12.2	19.3	19.4
1999 .....	10.0	4.7	3.6	3.7	16.8	9.2	9.1	8.1	9.3	13.2	19.2	18.5
Non-Hispanic white												
Both sexes												
2002 .....	16.3	2.4	3.0	4.0	30.8	17.8	15.8	14.8	14.0	16.9	26.2	28.9
2001 .....	15.8	3.0	3.5	4.1	28.7	17.6	15.7	13.9	13.3	16.1	26.9	29.0
2000 .....	15.8	3.3	3.7	4.4	29.5	17.0	15.3	13.7	14.2	16.1	26.2	31.3
1999 .....	15.6	4.1	3.4	4.4	28.7	16.8	14.8	13.1	13.7	17.5	26.8	30.9
Male												
2002 .....	22.3	2.7	3.3	4.6	42.1	26.1	22.0	21.2	19.4	21.8	35.2	52.6
2001 .....	21.7	2.9	3.9	4.8	40.1	25.9	21.9	19.9	18.2	21.0	37.3	53.1
2000 .....	21.5	3.4	4.1	5.1	40.3	24.7	21.6	19.6	18.9	20.7	35.6	60.0
1999 .....	21.1	4.7	3.6	5.1	38.8	24.2	20.6	18.5	18.5	22.6	37.6	61.2
Female												
2002 .....	10.5	2.1	2.8	3.3	19.0	9.4	9.7	8.4	8.8	12.7	20.2	18.6
2001 .....	10.1	3.1	3.1	3.3	17.0	9.3	9.6	8.1	8.7	11.9	20.0	19.0
2000 .....	10.3	3.2	3.2	3.6	18.4	9.3	9.0	7.9	9.8	12.3	20.0	19.8
1999 .....	10.3	3.5	3.1	3.7	18.3	9.4	9.0	7.9	9.2	13.2	19.7	18.8
Non-Hispanic black												
Both sexes												
2002 .....	14.8	5.4	4.9	4.3	21.0	20.3	17.2	16.3	15.7	16.0	20.2	18.5
2001 .....	15.2	*	5.3	5.0	22.2	19.6	17.6	17.2	16.6	17.4	19.9	16.9
2000 .....	15.5	6.9	5.6	4.5	21.6	21.1	17.6	18.3	16.3	18.4	18.8	20.7
1999 .....	15.5	7.6	6.5	5.1	21.7	19.2	17.9	17.4	16.7	18.1	24.0	25.5
Male												
2002 .....	22.0	*	6.3	5.0	30.1	32.7	25.7	24.8	25.4	25.6	36.6	45.9
2001 .....	22.7	*	5.6	6.5	32.4	31.0	26.9	27.5	26.8	24.2	34.8	35.8
2000 .....	23.1	7.2	6.0	5.4	31.1	33.6	27.9	27.6	27.2	29.8	34.6	44.6
1999 .....	22.8	*	7.0	6.2	31.5	30.7	26.9	27.5	25.8	26.8	41.3	61.8
Female												
2002 .....	8.3	*	3.5	3.6	11.9	9.1	9.7	9.0	8.0	9.1	10.9	*
2001 .....	8.5	*	4.9	3.4	12.1	9.2	9.4	8.4	8.4	12.6	11.5	9.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
Non-Hispanic black												
Female												
2000 .....	8.5	*	5.2	3.6	12.3	9.8	8.5	10.2	7.5	10.4	9.9	11.8
1999 .....	8.9	9.3	5.9	4.0	12.1	8.9	9.9	8.8	9.3	12.0	14.4	12.0
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.4	*	*	0.2	0.4	0.3	0.5	0.5	0.4	0.4	0.5	*
2001 .....	0.5	*	0.2	0.2	0.5	0.4	0.6	0.6	0.5	0.6	0.8	0.7
2000 .....	0.5	*	0.2	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999 .....	0.8	*	0.3	0.3	0.9	0.8	0.9	0.8	0.8	0.8	1.3	1.3
Male												
2002 .....	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.7	1.0	*
2001 .....	0.8	*	0.3	0.2	0.8	0.8	1.0	1.0	0.9	1.0	1.1	*
2000 .....	0.8	*	*	0.3	1.0	0.8	1.1	0.9	0.8	0.9	1.2	*
1999 .....	1.1	*	0.3	0.4	1.4	1.2	1.4	1.3	1.2	1.2	1.8	2.9
Female												
2002 .....	0.1	*	*	0.1	0.1	0.2	0.2	0.1	*	*	*	*
2001 .....	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.6	*
2000 .....	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.5	*
1999 .....	0.4	*	0.3	0.2	0.4	0.4	0.4	0.4	0.4	0.5	1.0	*
Hispanic												
Both sexes												
2002 .....	0.4	*	*	*	0.6	0.3	0.5	0.6	*	*	*	*
2001 .....	0.5	*	*	*	0.6	0.6	0.8	0.9	*	*	*	*
2000 .....	0.6	*	*	*	0.5	0.6	0.9	0.8	*	*	*	*
1999 .....	0.8	*	*	*	0.9	1.0	1.2	1.0	*	2.0	*	*
Male												
2002 .....	0.6	*	*	*	1.0	*	0.8	*	*	*	*	*
2001 .....	0.9	*	*	*	1.0	0.9	1.4	1.3	*	*	*	*
2000 .....	0.9	*	*	*	0.9	0.9	1.4	1.4	*	*	*	*
1999 .....	1.3	*	*	*	1.3	1.6	1.8	1.8	*	*	*	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01.V05-V06.V09.1.V09.3-V09.9.V10-V11, V15-V18.V19.3.V19.8-V19.9.V80.0-V80.2, V80.6-V80.9.V81.2-V81.9.V82.2-V82.9.V87.9, V88.9.V89.1.V89.3.V89.9)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.4	*	*	0.2	0.3	0.4	0.5	0.5	0.4	0.4	0.5	*
2001 .....	0.5	*	*	0.2	0.5	0.4	0.6	0.6	0.5	0.7	0.7	0.6
2000 .....	0.5	*	*	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999 .....	0.7	*	0.3	0.3	0.9	0.7	0.9	0.8	0.8	0.7	1.3	1.3
Male												
2002 .....	0.6	*	*	0.2	0.6	0.6	0.8	0.9	0.7	0.7	0.9	*
2001 .....	0.7	*	*	0.3	0.7	0.7	1.0	0.9	0.8	1.0	1.0	*
2000 .....	0.8	*	*	0.3	1.0	0.8	1.1	0.8	0.8	0.8	1.1	*
1999 .....	1.1	*	0.4	0.4	1.4	1.1	1.4	1.2	1.2	1.0	1.8	3.0
Female												
2002 .....	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2001 .....	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.5	*
2000 .....	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.4	*
1999 .....	0.4	*	*	0.2	0.5	0.4	0.4	0.4	0.4	0.5	1.0	*
Non-Hispanic white												
Both sexes												
2002 .....	0.4	*	*	0.2	0.4	0.4	0.5	0.5	0.4	0.4	0.4	*
2001 .....	0.5	*	*	0.2	0.5	0.4	0.5	0.6	0.4	0.6	0.7	0.5
2000 .....	0.5	*	*	0.3	0.7	0.5	0.6	0.6	0.4	0.5	0.7	0.8
1999 .....	0.8	*	0.3	0.3	1.0	0.7	0.9	0.8	0.8	0.7	1.2	1.3
Male												
2002 .....	0.6	*	*	0.2	0.6	0.6	0.7	0.8	0.6	0.6	0.8	*
2001 .....	0.7	*	*	0.2	0.8	0.7	0.8	0.9	0.8	0.9	0.9	*
2000 .....	0.8	*	*	0.3	1.0	0.8	1.0	0.8	0.7	0.7	1.2	*
1999 .....	1.1	*	*	0.4	1.4	1.1	1.3	1.2	1.2	1.0	1.7	2.8
Female												
2002 .....	0.1	*	*	*	*	0.2	0.2	*	*	*	*	*
2001 .....	0.2	*	*	*	0.2	*	0.2	0.2	*	0.4	0.5	*
2000 .....	0.3	*	*	0.2	0.3	0.2	0.2	0.4	*	*	0.4	*
1999 .....	0.4	*	*	0.2	0.5	0.4	0.4	0.4	0.4	0.4	0.9	*
Non-Hispanic black												
Both sexes												
2002 .....	0.5	*	*	*	*	*	0.7	0.9	*	*	*	*
2001 .....	0.6	*	*	0.4	0.4	0.5	1.1	0.9	0.9	*	*	*
2000 .....	0.6	*	*	*	0.5	0.6	1.0	0.8	*	*	*	*
1999 .....	0.7	*	*	0.3	0.6	0.8	0.9	1.0	*	*	*	*
Male												
2002 .....	0.8	*	*	*	*	*	1.3	1.9	*	*	*	*
2001 .....	0.9	*	*	*	*	*	1.8	1.4	*	*	*	*
2000 .....	0.9	*	*	*	0.9	1.0	1.6	1.4	*	*	*	*
1999 .....	1.2	*	*	*	1.0	1.4	1.6	1.8	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01.V05-V06.V09.1.V09.3-V09.9.V10-V11, V15-V18.V19.3.V19.8-V19.9.V80.0-V80.2, V80.6-V80.9.V81.2-V81.9.V82.2-V82.9.V87.9, V88.9.V89.1.V89.3.V89.9)												
Non-Hispanic black												
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.7	*	*	0.1	0.5	0.6	0.9	1.0	1.1	0.9	0.8	0.4
2001 .....	0.7	*	0.2	0.1	0.5	0.9	1.0	1.1	1.0	1.0	0.8	0.5
2000 .....	0.7	*	*	0.1	0.6	0.8	0.9	1.0	1.1	1.0	0.6	*
1999 .....	0.7	*	*	0.2	0.5	0.8	0.9	1.0	1.1	0.9	0.5	*
Male												
2002 .....	1.1	*	*	0.1	0.8	1.1	1.5	1.8	1.9	1.9	1.5	*
2001 .....	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.4	*
2000 .....	1.2	*	*	0.2	0.9	1.3	1.6	1.7	1.8	1.7	1.3	*
1999 .....	1.1	*	*	0.3	0.8	1.4	1.4	1.7	2.0	1.6	0.9	*
Female												
2002 .....	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.3	*	0.3	*
2001 .....	0.3	*	*	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.4	*
2000 .....	0.2	*	*	*	0.2	0.2	0.2	0.3	0.3	0.3	*	*
1999 .....	0.3	*	*	0.1	0.2	0.3	0.4	0.3	0.3	0.2	0.3	*
Hispanic												
Both sexes												
2002 .....	0.3	*	*	*	0.3	0.3	*	0.6	*	*	*	*
2001 .....	0.8	*	*	*	0.4	0.8	1.2	1.7	2.1	2.0	*	*
2000 .....	0.3	*	*	*	0.4	0.4	0.6	*	*	*	*	*
1999 .....	0.3	*	*	*	*	0.5	*	*	*	*	*	*
Male												
2002 .....	0.5	*	*	*	0.5	0.6	*	*	*	*	*	*
2001 .....	0.9	*	*	*	*	1.1	1.4	1.7	*	*	*	*
2000 .....	0.5	*	*	*	*	0.6	1.1	*	*	*	*	*
1999 .....	0.4	*	*	*	*	0.7	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.7	*	*	*	*	*	1.0	1.8	2.0	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.7	*	*	0.1	0.5	0.7	0.9	1.1	1.1	0.9	0.8	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2001 .....	0.7	*	*	0.1	0.6	0.9	1.0	1.1	1.0	0.9	0.7	*
2000 .....	0.7	*	*	0.1	0.6	0.8	1.0	1.0	1.1	1.0	0.6	*
1999 .....	0.7	*	*	0.2	0.6	0.9	1.0	1.1	1.1	0.9	0.6	*
Male												
2002 .....	1.2	*	*	0.1	0.8	1.2	1.6	1.9	2.0	2.0	1.5	*
2001 .....	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.3	*
2000 .....	1.2	*	*	0.2	1.0	1.5	1.7	1.8	1.9	1.7	1.3	*
1999 .....	1.2	*	*	0.3	0.8	1.5	1.6	1.8	2.1	1.7	1.0	*
Female												
2002 .....	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.4	*	0.3	*
2001 .....	0.2	*	*	*	0.2	0.3	0.3	0.3	0.2	0.2	0.4	*
2000 .....	0.2	*	*	*	0.2	0.2	0.3	0.3	0.4	0.3	*	*
1999 .....	0.3	*	*	0.1	0.3	0.3	0.4	0.4	0.3	0.3	0.3	*
Non-Hispanic white												
Both sexes												
2002 .....	0.8	*	*	0.1	0.6	0.8	1.0	1.2	1.2	1.0	0.8	*
2001 .....	0.8	*	*	0.1	0.7	1.0	1.1	1.1	1.0	0.9	0.7	*
2000 .....	0.8	*	*	0.2	0.7	1.0	1.1	1.1	1.2	1.0	0.7	*
1999 .....	0.8	*	*	0.2	0.6	1.0	1.1	1.1	1.1	0.9	0.5	*
Male												
2002 .....	1.3	*	*	*	0.9	1.3	1.8	2.0	2.1	2.0	1.4	*
2001 .....	1.3	*	*	0.2	1.1	1.6	1.8	1.9	1.8	1.7	1.4	*
2000 .....	1.4	*	*	0.2	1.1	1.7	1.8	1.9	2.0	1.8	1.3	*
1999 .....	1.3	*	*	0.3	1.0	1.6	1.7	1.9	2.0	1.7	1.0	*
Female												
2002 .....	0.3	*	*	*	0.3	0.2	0.3	0.3	0.4	*	0.3	*
2001 .....	0.3	*	*	*	0.3	0.3	0.4	0.3	0.2	0.2	0.3	*
2000 .....	0.3	*	*	*	0.3	0.3	0.3	0.3	0.4	0.3	*	*
1999 .....	0.3	*	*	*	0.3	0.4	0.5	0.4	0.3	0.3	*	*
Non-Hispanic black												
Both sexes												
2002 .....	0.4	*	*	*	*	0.4	0.6	0.8	0.9	*	*	*
2001 .....	0.5	*	*	*	*	0.5	0.5	1.0	0.9	*	*	*
2000 .....	0.4	*	*	*	*	0.5	0.5	0.6	*	*	*	*
1999 .....	0.5	*	*	*	0.4	0.5	0.7	0.8	1.1	*	*	*
Male												
2002 .....	0.8	*	*	*	*	*	1.0	1.3	1.8	*	*	*
2001 .....	0.8	*	*	*	*	0.9	1.0	1.8	2.1	*	*	*
2000 .....	0.7	*	*	*	*	0.9	1.0	1.1	*	*	*	*
1999 .....	0.9	*	*	*	*	0.9	1.2	1.4	2.3	*	*	*
Female												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	20.2	20.3	6.4	2.4	8.9	12.7	20.1	20.2	15.7	25.9	74.4	246.7
2001 .....	19.0	20.5	6.8	2.5	8.2	11.1	18.0	18.1	15.0	24.9	73.0	247.5
2000 .....	18.2	18.6	7.4	2.6	7.9	11.0	17.2	16.8	14.9	23.8	68.1	242.0
1999 .....	18.4	17.3	7.7	2.7	7.7	11.1	17.0	16.4	14.6	25.1	72.1	250.3
Male												
2002 .....	25.2	23.3	8.1	3.2	14.2	19.6	29.1	29.3	22.7	35.3	93.9	293.6
2001 .....	23.8	22.9	8.3	3.3	13.3	17.1	26.8	27.3	21.4	33.0	93.6	305.8
2000 .....	22.9	20.9	9.0	3.4	12.8	17.3	26.2	26.0	21.6	31.5	86.2	299.8
1999 .....	23.3	20.2	9.3	3.7	12.5	17.7	26.2	25.6	21.5	34.1	91.7	311.9
Female												
2002 .....	15.4	17.2	4.7	1.5	3.2	5.7	11.1	11.5	9.2	18.1	61.4	226.3
2001 .....	14.5	17.9	5.1	1.6	2.8	4.9	9.4	9.2	9.1	18.1	59.5	223.1
2000 .....	13.6	16.2	5.7	1.7	2.7	4.5	8.2	7.9	8.8	17.5	56.3	218.4
1999 .....	13.7	14.2	6.1	1.7	2.6	4.4	8.0	7.6	8.2	17.8	59.4	225.5
Hispanic												
Both sexes												
2002 .....	10.8	12.1	5.1	1.5	7.9	9.4	15.7	16.8	13.5	21.1	51.2	127.5
2001 .....	10.1	13.6	5.0	1.6	6.6	9.4	15.0	15.3	12.6	19.8	43.6	142.6
2000 .....	10.2	9.9	5.2	1.5	7.3	10.0	15.3	14.2	13.1	18.4	48.2	159.2
1999 .....	11.1	12.4	6.4	1.7	7.7	10.5	17.2	16.2	13.0	20.5	44.6	157.2
Male												
2002 .....	16.1	15.7	6.4	1.9	13.1	15.5	24.6	27.1	21.4	30.2	68.5	179.0
2001 .....	14.8	14.2	6.7	1.9	10.8	14.9	24.3	25.6	20.4	26.8	54.1	154.3
2000 .....	15.3	10.9	5.9	1.6	12.1	16.5	25.6	24.3	19.7	27.6	57.5	207.6
1999 .....	16.9	15.0	7.7	2.3	12.9	17.4	28.2	27.6	22.5	31.6	54.7	217.8
Female												
2002 .....	5.2	8.3	3.7	1.2	1.9	2.1	5.9	6.6	6.3	13.8	39.3	102.0
2001 .....	5.2	13.0	3.3	1.3	1.7	2.9	5.0	5.2	5.7	14.3	36.6	136.8
2000 .....	4.9	8.8	4.4	1.5	1.6	2.4	4.4	4.3	7.3	11.1	41.9	135.5
1999 .....	5.0	9.8	5.2	1.1	1.7	2.6	5.5	5.1	4.6	11.7	37.8	127.9
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	21.6	22.3	6.7	2.6	9.0	13.4	20.6	20.4	15.8	26.1	75.2	250.9
2001 .....	20.2	22.0	7.2	2.6	8.5	11.3	18.3	18.2	15.0	25.1	74.2	250.8
2000 .....	19.2	20.7	7.9	2.8	8.0	11.1	17.3	16.8	14.9	24.1	68.7	244.4
1999 .....	19.3	18.5	8.0	2.9	7.6	11.1	16.8	16.2	14.5	25.3	72.9	252.8
Male												
2002 .....	26.5	25.2	8.5	3.5	14.4	20.5	29.6	29.3	22.6	35.4	94.7	298.0
2001 .....	25.0	24.9	8.7	3.5	13.7	17.5	26.9	27.2	21.2	33.2	95.1	311.6
2000 .....	23.9	23.2	9.7	3.8	12.9	17.4	26.1	25.8	21.5	31.6	87.2	302.7
1999 .....	24.0	21.5	9.6	3.9	12.4	17.6	25.7	25.0	21.1	34.1	92.8	314.3
Female												
2002 .....	16.9	19.4	4.9	1.6	3.4	6.4	11.8	11.9	9.5	18.4	62.3	230.6
2001 .....	15.7	19.1	5.6	1.7	3.0	5.2	9.9	9.5	9.4	18.3	60.4	225.5
2000 .....	14.7	18.0	6.0	1.7	2.9	4.8	8.6	8.1	8.8	17.8	56.7	220.8
1999 .....	14.8	15.2	6.3	1.9	2.7	4.6	8.2	7.7	8.4	18.1	60.1	228.2

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Nontransport accidents (W00-X59,Y86)</b>												
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	23.1	17.2	6.1	2.0	9.8	14.7	21.4	19.9	15.1	25.6	77.0	261.2
2001 .....	21.5	17.8	7.1	2.2	9.1	12.1	18.5	17.2	14.2	24.7	75.4	259.4
2000 .....	20.1	16.6	7.3	2.3	8.3	11.5	17.2	15.6	14.1	23.4	69.6	253.2
1999 .....	20.0	14.7	7.4	2.4	7.8	11.3	16.7	14.9	13.3	24.2	73.2	260.8
<b>Male</b>												
2002 .....	27.8	20.4	7.6	2.7	15.6	22.2	30.5	28.1	21.0	34.3	96.0	307.6
2001 .....	25.9	20.0	8.8	2.9	14.7	18.6	27.2	25.3	19.6	32.5	95.6	320.4
2000 .....	24.5	19.1	9.1	3.2	13.5	18.1	26.0	23.6	19.9	30.4	87.2	311.5
1999 .....	24.3	17.1	8.9	3.2	12.5	17.9	25.4	22.7	19.1	32.0	92.4	322.6
<b>Female</b>												
2002 .....	18.6	13.9	4.6	1.2	3.8	7.1	12.3	11.8	9.4	18.3	64.2	241.1
2001 .....	17.2	15.5	5.3	1.5	3.3	5.5	9.8	9.3	9.2	18.1	62.0	234.0
2000 .....	16.0	14.0	5.4	1.3	3.0	4.9	8.4	7.7	8.7	17.6	58.1	229.7
1999 .....	16.0	12.3	5.9	1.5	2.9	4.6	8.0	7.3	7.9	17.6	60.6	236.2
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	18.3	44.5	9.5	4.7	6.9	11.4	21.4	29.6	24.6	32.4	62.8	156.7
2001 .....	18.1	41.6	8.7	4.3	6.8	10.5	21.7	29.4	23.9	32.5	68.2	166.1
2000 .....	18.0	39.5	10.9	4.8	7.1	11.3	21.5	28.1	23.8	33.1	65.4	161.2
1999 .....	19.4	36.4	11.1	5.3	7.8	12.7	21.2	29.1	26.6	39.1	77.8	181.1
<b>Male</b>												
2002 .....	25.5	46.7	12.5	6.5	11.6	18.0	31.7	45.3	40.6	48.5	90.6	215.5
2001 .....	25.0	48.2	10.0	6.0	11.0	16.2	31.6	47.9	38.0	47.5	101.4	220.6
2000 .....	25.5	42.3	12.9	6.5	11.7	17.9	32.7	46.6	39.8	47.8	96.9	231.3
1999 .....	27.5	42.3	13.7	7.3	13.5	20.0	33.3	48.7	42.5	60.1	107.6	249.6
<b>Female</b>												
2002 .....	11.8	42.3	6.5	2.9	2.1	5.4	12.3	16.2	11.7	21.1	47.1	134.2
2001 .....	11.8	34.6	7.4	2.4	2.5	5.4	12.9	13.5	12.6	21.9	49.5	145.9
2000 .....	11.3	36.6	8.8	3.0	2.6	5.4	11.6	12.2	10.9	22.8	47.8	135.2
1999 .....	12.0	30.3	8.3	3.3	2.1	6.1	10.7	12.3	13.8	24.3	61.1	155.5
<b>Falls (W00-W19)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	5.6	*	0.2	0.1	0.6	0.8	1.5	2.5	4.1	10.8	38.3	130.4
2001 .....	5.3	0.6	0.2	0.2	0.6	0.9	1.4	2.6	4.0	10.0	35.3	121.5
2000 .....	4.7	*	0.2	0.1	0.6	0.8	1.3	2.3	3.9	9.0	31.1	112.6
1999 .....	4.7	*	0.4	0.1	0.6	0.9	1.4	2.3	3.7	9.0	31.5	110.2
<b>Male</b>												
2002 .....	6.0	*	0.4	0.1	1.0	1.3	2.3	3.9	6.0	14.5	48.1	158.9
2001 .....	5.8	*	0.3	0.2	1.1	1.5	2.4	4.3	5.9	13.1	45.7	155.7
2000 .....	5.2	*	0.3	0.1	1.0	1.4	2.2	3.7	5.8	11.9	39.9	145.4
1999 .....	5.2	*	0.5	0.2	1.1	1.5	2.4	3.7	5.7	12.4	40.5	142.5
<b>Female</b>												
2002 .....	5.3	*	*	*	0.2	0.2	0.6	1.2	2.3	7.7	31.8	118.1
2001 .....	4.8	*	*	0.1	0.1	0.2	0.5	1.0	2.2	7.4	28.5	107.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
All origins <sup>3</sup>												
Female												
2000 .....	4.3	*	*	*	0.2	0.2	0.5	0.9	2.1	6.6	25.3	99.2
1999 .....	4.3	*	*	*	0.2	0.2	0.5	0.9	1.9	6.3	25.7	97.2
Hispanic												
Both sexes												
2002 .....	2.1	*	*	*	0.7	0.9	1.6	2.5	4.4	10.0	29.2	71.5
2001 .....	2.0	*	*	*	0.7	1.1	1.8	2.6	4.2	9.1	23.6	70.0
2000 .....	2.0	*	*	*	0.7	1.1	1.3	2.5	4.3	8.0	26.5	81.0
1999 .....	1.9	*	*	*	0.7	1.1	1.4	2.2	3.8	8.1	24.2	86.3
Male												
2002 .....	2.8	*	*	*	1.3	1.6	2.7	3.8	7.0	13.8	36.9	97.0
2001 .....	2.8	*	*	*	1.2	1.9	3.3	4.5	7.1	11.4	27.7	72.4
2000 .....	2.7	*	*	*	1.2	2.0	2.3	4.4	6.6	12.7	30.5	104.8
1999 .....	2.7	*	*	*	1.3	1.9	2.4	3.8	7.2	11.4	31.3	120.8
Female												
2002 .....	1.4	*	*	*	*	*	*	1.1	2.1	7.0	23.9	58.9
2001 .....	1.3	*	*	*	*	*	*	*	*	7.2	20.9	68.9
2000 .....	1.2	*	*	*	*	*	*	*	2.3	4.3	23.8	69.2
1999 .....	1.1	*	*	*	*	*	*	*	*	5.5	19.4	69.7
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	6.2	*	0.2	0.1	0.6	0.7	1.5	2.5	4.0	10.8	38.6	132.6
2001 .....	5.7	*	0.2	0.1	0.6	0.8	1.4	2.6	3.9	10.0	35.8	123.1
2000 .....	5.1	*	0.2	0.1	0.6	0.7	1.3	2.3	3.9	9.1	31.2	113.5
1999 .....	5.1	*	0.4	0.1	0.6	0.8	1.4	2.2	3.7	9.0	31.7	110.7
Male												
2002 .....	6.5	*	0.3	0.1	0.9	1.3	2.3	3.9	5.9	14.4	48.4	161.3
2001 .....	6.2	*	0.3	0.2	1.1	1.4	2.3	4.2	5.8	13.1	46.4	159.0
2000 .....	5.5	*	*	0.2	1.0	1.2	2.2	3.6	5.7	11.9	40.2	146.5
1999 .....	5.5	*	0.5	0.2	1.0	1.3	2.3	3.6	5.5	12.4	40.6	142.9
Female												
2002 .....	5.9	*	*	*	0.2	0.2	0.7	1.2	2.3	7.7	32.1	120.2
2001 .....	5.3	*	*	*	0.1	0.3	0.5	1.0	2.2	7.4	28.8	108.1
2000 .....	4.7	*	*	*	0.2	0.2	0.5	1.0	2.1	6.8	25.3	100.1
1999 .....	4.6	*	*	*	0.2	0.3	0.5	0.9	2.0	6.3	25.9	97.8
Non-Hispanic white												
Both sexes												
2002 .....	7.1	*	*	0.1	0.7	0.8	1.5	2.5	4.1	11.1	40.8	140.4
2001 .....	6.6	*	0.2	0.2	0.8	0.9	1.4	2.6	3.9	10.3	37.6	129.9
2000 .....	5.8	*	*	0.1	0.7	0.7	1.4	2.3	3.8	9.2	32.7	120.2
1999 .....	5.7	*	0.4	0.2	0.7	0.8	1.5	2.1	3.5	9.1	33.1	116.6
Male												
2002 .....	7.3	*	*	*	1.0	1.4	2.3	3.9	5.9	14.8	50.6	169.2
2001 .....	7.0	*	*	0.2	1.3	1.5	2.3	4.1	5.7	13.5	48.2	166.0
2000 .....	6.2	*	*	0.2	1.1	1.3	2.2	3.6	5.6	12.0	41.7	154.1
1999 .....	6.1	*	0.5	0.2	1.1	1.4	2.3	3.4	5.1	12.2	42.0	149.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
Non-Hispanic white												
Female												
2002 .....	7.0	*	*	*	0.2	0.2	0.7	1.2	2.4	8.1	34.2	128.0
2001 .....	6.2	*	*	*	0.2	0.3	0.5	1.1	2.2	7.7	30.6	114.8
2000 .....	5.5	*	*	*	0.2	0.2	0.5	1.0	2.1	6.9	26.8	106.5
1999 .....	5.4	*	*	*	0.2	0.3	0.6	0.8	1.9	6.4	27.2	103.4
Non-Hispanic black												
Both sexes												
2002 .....	2.3	*	*	*	*	0.4	1.3	2.8	4.3	8.2	16.3	49.3
2001 .....	2.3	*	*	*	*	0.5	1.6	3.0	4.4	8.8	18.7	44.0
2000 .....	2.1	*	*	*	*	0.4	1.3	2.7	4.8	8.3	15.2	39.1
1999 .....	2.5	*	*	*	0.4	0.8	1.3	3.5	5.5	9.3	17.2	45.3
Male												
2002 .....	3.0	*	*	*	*	0.8	2.2	4.7	6.8	11.9	24.9	76.6
2001 .....	3.2	*	*	*	*	*	2.7	5.4	6.9	12.6	29.1	68.1
2000 .....	2.8	*	*	*	*	*	2.1	4.6	8.2	11.4	21.6	61.1
1999 .....	3.4	*	*	*	*	1.4	2.5	5.9	8.7	15.0	24.1	58.2
Female												
2002 .....	1.6	*	*	*	*	*	*	1.2	2.2	5.6	11.4	38.9
2001 .....	1.6	*	*	*	*	*	*	*	2.3	6.1	12.8	35.1
2000 .....	1.5	*	*	*	*	*	*	1.0	2.0	6.1	11.6	31.0
1999 .....	1.7	*	*	*	*	*	*	1.4	2.9	5.3	13.3	40.4
Accidental discharge of firearms (W32-W34)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.3	*	*	0.1	0.5	0.4	0.3	0.2	0.2	0.2	0.3	*
2001 .....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000 .....	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999 .....	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*
Male												
2002 .....	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2001 .....	0.5	*	*	0.2	0.9	0.5	0.5	0.5	0.5	0.3	0.7	*
2000 .....	0.5	*	*	0.3	0.9	0.5	0.6	0.4	0.4	0.4	0.5	*
1999 .....	0.5	*	*	0.3	1.2	0.6	0.5	0.4	0.4	0.4	0.4	*
Female												
2002 .....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000 .....	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999 .....	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	0.2	*	*	*	0.4	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	0.5	0.3	*	*	*	*	*	*
2000 .....	0.2	*	*	*	0.4	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	0.5	*	*	*	*	*	*	*
Male												
2002 .....	0.3	*	*	*	0.6	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
Hispanic												
Male												
2001 .....	0.4	*	*	*	0.9	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	0.7	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	0.9	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.3	*	*	0.1	0.5	0.4	0.3	0.3	0.2	0.2	0.2	*
2001 .....	0.3	*	*	0.2	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000 .....	0.3	*	*	0.2	0.5	0.4	0.4	0.2	0.2	0.2	0.2	*
1999 .....	0.3	*	*	0.2	0.7	0.4	0.3	0.3	0.2	0.2	0.2	*
Male												
2002 .....	0.5	*	*	0.2	1.0	0.7	0.5	0.5	0.4	0.4	0.5	*
2001 .....	0.5	*	*	0.3	0.9	0.5	0.6	0.5	0.5	0.3	0.7	*
2000 .....	0.5	*	*	0.3	1.0	0.6	0.6	0.4	0.4	0.4	0.5	*
1999 .....	0.5	*	*	0.3	1.2	0.6	0.5	0.5	0.4	0.4	0.4	*
Female												
2002 .....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000 .....	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999 .....	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.2	0.2	0.2	*
2001 .....	0.3	*	*	0.2	0.5	0.3	0.4	0.3	0.3	0.2	0.3	*
2000 .....	0.3	*	*	0.2	0.5	0.3	0.4	0.2	0.2	0.2	0.2	*
1999 .....	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*
Male												
2002 .....	0.5	*	*	0.2	0.8	0.6	0.5	0.5	0.4	0.5	0.5	*
2001 .....	0.5	*	*	0.3	0.8	0.6	0.6	0.5	0.5	0.3	0.7	*
2000 .....	0.5	*	*	0.3	0.8	0.5	0.6	0.4	0.4	0.4	0.5	*
1999 .....	0.5	*	*	0.3	1.0	0.6	0.5	0.5	0.4	0.4	*	*
Female												
2002 .....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	0.4	*	*	*	1.0	0.8	0.4	*	*	*	*	*
2001 .....	0.3	*	*	*	0.8	*	*	*	*	*	*	*
2000 .....	0.4	*	*	*	0.9	0.6	*	*	*	*	*	*
1999 .....	0.4	*	*	*	1.2	0.4	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
Non-Hispanic black												
Male												
2002 .....	0.8	*	*	*	1.9	1.6	*	*	*	*	*	*
2001 .....	0.6	*	*	*	1.6	*	*	*	*	*	*	*
2000 .....	0.7	*	*	*	1.6	1.2	*	*	*	*	*	*
1999 .....	0.8	*	*	*	2.2	0.9	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Accidental drowning and submersion (W65-W74)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.2	1.6	2.9	0.8	1.5	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001 .....	1.2	1.7	3.0	0.8	1.5	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000 .....	1.2	2.0	3.2	0.9	1.6	1.1	1.1	0.9	0.9	1.0	1.3	1.5
1999 .....	1.3	1.8	3.2	0.9	1.7	1.1	1.1	1.0	0.9	1.0	1.3	1.6
Male												
2002 .....	1.9	1.9	3.9	1.2	2.8	1.9	2.0	1.5	1.3	1.6	2.1	3.6
2001 .....	1.8	1.5	3.7	1.1	2.7	1.6	1.7	1.6	1.3	1.5	2.0	3.5
2000 .....	2.0	2.2	4.0	1.3	2.9	1.8	1.8	1.6	1.5	1.6	1.8	2.8
1999 .....	2.0	2.0	4.0	1.3	3.0	2.0	1.8	1.7	1.5	1.6	2.3	2.3
Female												
2002 .....	0.5	1.2	1.9	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001 .....	0.5	1.9	2.2	0.5	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000 .....	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999 .....	0.5	1.6	2.4	0.5	0.3	0.2	0.3	0.3	0.4	0.6	0.6	1.4
Hispanic												
Both sexes												
2002 .....	1.2	*	2.1	0.4	2.1	1.1	0.9	0.8	*	*	*	*
2001 .....	1.1	*	2.6	0.6	1.5	1.1	1.0	0.7	*	*	*	*
2000 .....	1.2	*	2.5	0.6	2.2	1.0	0.8	0.6	*	*	*	*
1999 .....	1.4	*	2.7	0.7	2.3	1.3	0.9	0.8	*	*	*	*
Male												
2002 .....	2.0	*	3.0	0.6	3.7	2.0	1.6	1.5	*	*	*	*
2001 .....	1.8	*	3.5	0.8	2.7	1.8	1.7	1.2	*	*	*	*
2000 .....	1.9	*	2.9	0.6	3.8	1.7	1.4	*	*	*	*	*
1999 .....	2.3	*	2.9	1.1	4.1	2.2	1.5	1.6	*	*	*	*
Female												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	1.7	*	*	*	*	*	*	*	*	*
2000 .....	0.5	*	2.2	0.6	*	*	*	*	*	*	*	*
1999 .....	0.5	*	2.5	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.2	1.4	3.1	0.9	1.4	1.1	1.2	0.9	0.8	0.9	1.3	1.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2001 .....	1.1	1.6	3.1	0.9	1.5	0.9	1.0	0.9	0.8	0.9	1.3	1.7
2000 .....	1.2	2.1	3.4	1.0	1.5	1.1	1.1	1.0	0.9	1.0	1.3	1.5
1999 .....	1.2	1.7	3.3	0.9	1.5	1.1	1.1	1.0	0.9	1.0	1.3	1.7
Male												
2002 .....	1.9	1.6	4.1	1.3	2.6	1.8	2.0	1.5	1.3	1.6	2.0	3.6
2001 .....	1.8	1.3	3.8	1.2	2.6	1.6	1.7	1.6	1.3	1.5	2.1	3.4
2000 .....	2.0	2.4	4.3	1.4	2.7	1.8	1.9	1.6	1.6	1.6	1.8	2.8
1999 .....	2.0	1.7	4.2	1.4	2.7	1.9	1.8	1.7	1.5	1.6	2.3	2.4
Female												
2002 .....	0.5	*	2.1	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001 .....	0.5	1.9	2.4	0.5	0.3	0.2	0.4	0.3	0.4	0.4	0.7	1.0
2000 .....	0.5	1.9	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999 .....	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.4	0.6	0.6	1.4
Non-Hispanic white												
Both sexes												
2002 .....	1.1	1.4	3.3	0.5	1.2	1.0	1.2	0.9	0.8	0.9	1.2	1.6
2001 .....	1.1	1.5	3.5	0.5	1.3	0.8	1.0	0.9	0.8	0.9	1.2	1.7
2000 .....	1.1	1.7	3.7	0.7	1.3	1.0	1.1	0.9	0.9	1.0	1.3	1.4
1999 .....	1.1	1.3	3.6	0.5	1.3	1.0	1.1	1.0	0.9	1.0	1.2	1.6
Male												
2002 .....	1.7	*	4.3	0.7	2.2	1.7	1.9	1.4	1.2	1.5	2.0	3.7
2001 .....	1.6	*	4.3	0.7	2.2	1.4	1.5	1.5	1.2	1.5	2.1	3.5
2000 .....	1.8	1.9	4.7	1.0	2.2	1.7	1.8	1.5	1.6	1.5	1.8	2.7
1999 .....	1.8	*	4.5	0.7	2.2	1.6	1.8	1.7	1.5	1.4	2.2	2.4
Female												
2002 .....	0.5	*	2.2	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.7	0.7
2001 .....	0.5	1.7	2.6	0.4	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000 .....	0.5	*	2.6	0.4	0.3	0.3	0.4	0.3	0.3	0.5	0.9	0.9
1999 .....	0.5	*	2.6	0.3	0.3	0.3	0.4	0.4	0.4	0.6	0.6	1.3
Non-Hispanic black												
Both sexes												
2002 .....	1.7	*	2.6	2.1	2.1	1.4	1.3	1.3	0.8	*	*	*
2001 .....	1.6	*	2.0	2.1	2.3	1.2	1.3	1.2	1.3	*	*	*
2000 .....	1.7	4.3	2.3	2.1	2.3	1.1	1.4	1.3	*	*	*	*
1999 .....	1.8	3.7	2.5	2.5	2.5	1.6	1.1	0.9	1.2	1.3	*	*
Male												
2002 .....	2.9	*	3.5	3.3	4.1	2.7	2.5	2.3	*	*	*	*
2001 .....	2.8	*	2.4	3.1	4.4	2.5	2.2	2.1	2.4	*	*	*
2000 .....	2.8	*	2.8	3.1	4.4	2.1	2.6	2.4	*	*	*	*
1999 .....	3.2	*	3.4	3.8	4.6	3.2	2.1	1.9	2.1	*	*	*
Female												
2002 .....	0.5	*	*	0.8	*	*	*	*	*	*	*	*
2001 .....	0.5	*	*	1.0	*	*	*	*	*	*	*	*
2000 .....	0.6	*	1.8	1.0	*	*	*	*	*	*	*	*
1999 .....	0.6	*	*	1.1	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.1	0.9	1.4	0.6	0.5	0.7	0.8	1.1	1.3	2.1	3.4	4.8
2001 .....	1.2	1.2	1.5	0.6	0.5	0.6	1.0	1.1	1.5	2.2	3.4	5.7
2000 .....	1.2	1.0	1.9	0.6	0.5	0.6	0.9	1.2	1.5	2.2	3.8	6.2
1999 .....	1.2	1.1	2.0	0.6	0.5	0.7	0.9	1.0	1.3	2.3	4.0	5.9
Male												
2002 .....	1.4	1.0	1.6	0.7	0.6	0.8	1.1	1.5	1.7	2.9	4.5	7.0
2001 .....	1.4	1.5	1.8	0.7	0.7	0.8	1.4	1.5	1.9	2.8	4.1	8.3
2000 .....	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.6	2.1	2.8	5.0	9.4
1999 .....	1.5	1.1	2.2	0.7	0.7	1.0	1.3	1.4	1.7	2.9	5.1	9.3
Female												
2002 .....	0.8	*	1.2	0.5	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.9
2001 .....	0.9	1.0	1.1	0.5	0.4	0.4	0.6	0.7	1.1	1.8	2.9	4.6
2000 .....	0.9	*	1.5	0.6	0.4	0.4	0.6	0.7	1.0	1.7	3.0	4.9
1999 .....	0.9	1.1	1.7	0.6	0.4	0.4	0.5	0.6	1.0	1.7	3.2	4.5
Hispanic												
Both sexes												
2002 .....	0.6	*	1.2	0.4	0.4	*	0.5	0.6	*	1.7	*	*
2001 .....	0.5	*	*	0.3	*	0.3	0.5	*	*	*	*	*
2000 .....	0.5	*	0.7	*	0.3	*	0.5	*	*	*	*	*
1999 .....	0.6	*	1.4	0.4	0.4	*	0.4	*	*	*	*	*
Male												
2002 .....	0.7	*	*	*	*	*	0.7	*	*	*	*	*
2001 .....	0.6	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.6	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.7	*	1.6	*	*	*	*	*	*	*	*	*
Female												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.2	1.0	1.5	0.7	0.5	0.8	0.9	1.2	1.3	2.1	3.4	4.8
2001 .....	1.3	1.3	1.7	0.7	0.6	0.7	1.0	1.1	1.5	2.3	3.4	5.7
2000 .....	1.3	1.0	2.2	0.7	0.5	0.7	0.9	1.2	1.5	2.2	3.8	6.3
1999 .....	1.3	1.2	2.1	0.7	0.5	0.8	1.0	1.0	1.4	2.3	4.0	6.0
Male												
2002 .....	1.5	*	1.7	0.8	0.7	1.0	1.2	1.5	1.7	2.9	4.6	7.0
2001 .....	1.6	1.5	2.0	0.8	0.8	0.9	1.5	1.6	2.0	2.8	4.2	8.3
2000 .....	1.6	*	2.6	0.8	0.7	0.9	1.3	1.7	2.1	2.8	5.0	9.3
1999 .....	1.6	*	2.3	0.8	0.7	1.1	1.4	1.4	1.7	3.0	5.1	9.2
Female												
2002 .....	0.9	*	1.2	0.6	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.8
2001 .....	1.0	*	1.3	0.5	0.4	0.5	0.6	0.7	1.1	1.9	2.9	4.6
2000 .....	1.0	*	1.7	0.7	0.4	0.5	0.6	0.7	1.0	1.7	3.1	5.0
1999 .....	1.0	*	1.8	0.6	0.4	0.5	0.6	0.6	1.1	1.8	3.3	4.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
Non-Hispanic white												
Both sexes												
2002 .....	1.0	*	1.0	0.4	0.4	0.7	0.8	1.1	1.1	1.8	3.0	3.9
2001 .....	1.1	1.1	1.3	0.5	0.5	0.6	0.9	1.0	1.3	2.0	2.9	4.7
2000 .....	1.1	*	1.5	0.5	0.5	0.6	0.8	1.1	1.3	1.8	3.1	5.1
1999 .....	1.1	0.9	1.5	0.5	0.5	0.8	0.9	0.9	1.2	1.9	3.2	4.7
Male												
2002 .....	1.3	*	1.1	0.6	0.6	0.9	1.1	1.4	1.4	2.4	4.2	5.7
2001 .....	1.4	*	1.7	0.6	0.7	0.8	1.3	1.3	1.7	2.3	3.5	6.9
2000 .....	1.4	*	1.8	0.6	0.6	0.8	1.1	1.6	1.7	2.3	4.1	8.1
1999 .....	1.4	*	1.7	0.6	0.7	1.1	1.3	1.2	1.5	2.3	4.2	7.8
Female												
2002 .....	0.8	*	0.8	0.3	0.3	0.5	0.5	0.7	0.8	1.3	2.2	3.2
2001 .....	0.9	*	0.9	0.4	0.3	0.4	0.6	0.6	1.0	1.7	2.5	3.8
2000 .....	0.9	*	1.1	0.5	0.3	0.4	0.5	0.7	0.8	1.5	2.4	3.9
1999 .....	0.8	*	1.3	0.5	0.3	0.4	0.5	0.6	0.8	1.5	2.6	3.5
Non-Hispanic black												
Both sexes												
2002 .....	2.1	*	3.7	1.5	0.8	1.2	1.1	2.2	3.3	5.0	8.7	16.0
2001 .....	2.3	*	3.4	1.3	0.7	1.2	1.8	2.4	3.6	6.1	9.9	17.8
2000 .....	2.6	*	5.1	1.6	0.7	1.2	1.7	2.4	4.2	6.1	13.0	20.7
1999 .....	2.6	*	4.7	1.6	0.7	1.0	1.8	2.2	3.4	7.0	13.5	22.0
Male												
2002 .....	2.6	*	4.3	1.6	1.0	1.4	1.7	3.1	4.6	7.8	11.1	26.3
2001 .....	3.0	*	3.7	1.6	0.8	1.4	2.8	3.9	5.1	9.0	13.2	26.6
2000 .....	3.2	*	5.8	1.7	0.9	1.6	2.4	3.5	6.3	9.0	17.9	28.2
1999 .....	3.1	*	5.3	1.8	0.9	1.2	2.7	3.3	4.3	10.4	18.4	29.7
Female												
2002 .....	1.6	*	3.0	1.3	*	1.0	*	1.4	2.2	3.1	7.3	12.1
2001 .....	1.8	*	3.1	1.0	*	1.1	1.0	1.1	2.4	4.0	8.1	14.5
2000 .....	2.0	*	4.3	1.4	*	0.9	1.2	1.4	2.5	4.0	10.2	17.9
1999 .....	2.1	*	4.1	1.4	*	0.8	1.1	1.3	2.8	4.7	10.7	19.1
Accidental poisoning and exposure to noxious substances (X40-X49)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	6.1	0.6	0.2	0.1	4.1	7.8	13.4	11.7	4.1	2.1	2.3	3.9
2001 .....	4.9	*	0.2	0.1	3.4	6.3	11.2	9.1	3.2	1.6	2.2	3.5
2000 .....	4.5	*	0.2	0.1	3.0	6.0	10.3	8.1	2.8	1.5	2.0	4.3
1999 .....	4.4	*	0.2	0.1	2.5	5.9	10.1	7.8	2.8	1.6	2.2	3.4
Male												
2002 .....	8.5	1.0	*	0.1	6.3	11.5	18.4	15.9	5.4	2.5	2.7	5.1
2001 .....	7.1	*	0.3	0.1	5.3	9.3	15.7	13.0	4.1	2.0	2.7	3.8
2000 .....	6.6	*	0.3	0.1	4.6	9.0	15.1	12.0	3.4	1.8	2.1	6.2
1999 .....	6.5	*	*	0.1	3.8	9.0	14.9	11.7	3.9	2.1	2.4	4.4
Female												
2002 .....	3.7	*	*	*	1.9	4.0	8.4	7.6	3.0	1.7	2.1	3.4

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
All origins <sup>3</sup>												
Female												
2001 .....	2.9	*	*	0.1	1.4	3.3	6.7	5.2	2.3	1.2	1.9	3.3
2000 .....	2.5	*	*	*	1.3	2.8	5.6	4.4	2.3	1.3	1.9	3.6
1999 .....	2.3	*	*	*	1.1	2.6	5.3	4.0	1.8	1.3	2.0	3.0
Hispanic												
Both sexes												
2002 .....	4.3	*	*	*	2.9	5.0	10.1	10.1	4.0	2.2	*	*
2001 .....	3.6	*	*	*	2.2	4.9	8.7	7.7	3.5	*	*	*
2000 .....	3.7	*	*	*	2.0	5.3	9.6	8.0	3.1	*	*	*
1999 .....	4.4	*	*	*	2.3	6.1	11.9	9.8	3.5	*	*	*
Male												
2002 .....	6.6	*	*	*	4.4	7.8	15.4	15.8	6.4	*	*	*
2001 .....	5.6	*	*	*	3.1	7.4	13.6	12.9	5.8	*	*	*
2000 .....	6.0	*	*	*	3.2	8.4	15.6	13.6	4.4	*	*	*
1999 .....	7.3	*	*	*	3.6	9.8	19.4	16.4	6.6	*	*	*
Female												
2002 .....	1.7	*	*	*	1.1	1.6	4.3	4.4	*	*	*	*
2001 .....	1.5	*	*	*	1.1	1.9	3.4	2.6	*	*	*	*
2000 .....	1.3	*	*	*	*	1.6	3.1	2.5	*	*	*	*
1999 .....	1.4	*	*	*	0.8	1.7	3.9	3.3	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	6.3	0.6	0.2	0.1	4.4	8.4	13.8	11.8	4.1	2.1	2.3	4.1
2001 .....	5.1	*	0.2	0.1	3.6	6.6	11.4	9.1	3.1	1.6	2.3	3.6
2000 .....	4.6	*	0.2	0.1	3.1	6.0	10.3	8.0	2.8	1.5	2.0	4.4
1999 .....	4.3	*	0.2	0.1	2.5	5.8	9.7	7.5	2.7	1.6	2.2	3.5
Male												
2002 .....	8.8	*	*	0.1	6.6	12.3	18.7	15.9	5.2	2.4	2.6	5.3
2001 .....	7.2	*	*	0.1	5.7	9.6	15.9	12.9	3.9	2.0	2.8	3.9
2000 .....	6.6	*	*	0.1	4.9	9.1	14.8	11.7	3.3	1.7	2.1	6.2
1999 .....	6.3	*	*	0.1	3.8	8.8	14.2	11.1	3.6	2.0	2.5	4.4
Female												
2002 .....	4.0	*	*	*	2.0	4.5	8.9	7.9	3.0	1.8	2.1	3.5
2001 .....	3.1	*	*	*	1.5	3.5	7.1	5.4	2.3	1.2	1.9	3.4
2000 .....	2.7	*	*	*	1.4	3.0	5.9	4.4	2.3	1.3	2.0	3.6
1999 .....	2.4	*	*	*	1.2	2.8	5.4	4.0	1.8	1.3	2.0	3.1
Non-Hispanic white												
Both sexes												
2002 .....	6.6	*	*	0.1	5.3	9.6	14.5	11.4	3.7	2.0	2.3	4.1
2001 .....	5.2	*	*	0.1	4.3	7.3	11.7	8.3	2.7	1.5	2.1	3.4
2000 .....	4.6	*	*	0.1	3.7	6.5	10.4	7.2	2.5	1.4	2.1	4.1
1999 .....	4.2	*	*	0.1	2.8	6.0	9.6	6.7	2.4	1.5	2.3	3.3
Male												
2002 .....	9.1	*	*	0.2	8.0	13.9	19.6	15.0	4.3	2.2	2.5	5.4
2001 .....	7.3	*	*	*	6.7	10.7	16.3	11.5	3.1	1.9	2.5	3.8
2000 .....	6.6	*	*	*	5.8	9.8	14.9	10.3	2.7	1.5	2.2	5.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
Non-Hispanic white												
Male												
1999 .....	6.1	*	*	*	4.3	9.2	14.0	9.6	2.9	1.8	2.5	4.6
Female												
2002 .....	4.3	*	*	*	2.5	5.2	9.3	7.8	3.1	1.9	2.2	3.5
2001 .....	3.1	*	*	*	1.7	3.8	7.1	5.2	2.4	1.2	1.9	3.2
2000 .....	2.7	*	*	*	1.5	3.2	5.8	4.2	2.3	1.4	2.0	3.4
1999 .....	2.4	*	*	*	1.3	2.8	5.1	3.8	1.9	1.2	2.1	2.8
Non-Hispanic black												
Both sexes												
2002 .....	6.4	*	*	*	1.5	5.8	14.1	18.3	8.6	2.5	2.8	*
2001 .....	6.0	*	*	*	1.6	5.3	13.4	17.0	6.4	2.4	3.7	6.6
2000 .....	5.7	*	*	*	1.4	5.7	13.2	15.7	5.5	2.5	*	8.3
1999 .....	5.8	*	*	*	1.7	6.3	13.2	15.6	6.0	3.0	*	*
Male												
2002 .....	9.3	*	*	*	2.2	8.7	19.8	27.1	15.2	4.5	*	*
2001 .....	8.6	*	*	*	2.3	7.6	18.2	27.0	11.0	3.7	6.0	*
2000 .....	8.4	*	*	*	1.8	8.5	19.0	25.3	9.5	4.3	*	*
1999 .....	8.8	*	*	*	2.8	9.3	19.5	25.9	11.0	4.4	*	*
Female												
2002 .....	3.8	*	*	*	0.7	3.2	9.1	10.8	3.3	*	*	*
2001 .....	3.6	*	*	*	0.9	3.2	9.2	8.4	2.6	*	*	*
2000 .....	3.2	*	*	*	1.0	3.1	8.0	7.5	2.4	*	*	*
1999 .....	3.1	*	*	*	*	3.5	7.7	6.7	1.9	2.1	*	*
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	6.0	16.8	1.6	0.7	1.6	2.1	3.0	3.7	5.2	9.8	28.8	105.6
2001 .....	6.2	16.6	1.8	0.6	1.6	2.0	3.1	4.2	5.3	10.0	30.5	115.0
2000 .....	6.2	15.0	1.7	0.6	1.7	2.2	3.2	4.0	5.6	9.9	29.8	117.1
1999 .....	6.6	13.8	1.9	0.8	1.7	2.2	3.2	4.1	5.6	11.0	32.9	128.9
Male												
2002 .....	6.9	18.8	2.0	0.9	2.6	3.4	4.8	6.0	7.9	13.4	36.0	118.6
2001 .....	7.2	18.9	2.1	0.9	2.6	3.4	5.0	6.6	7.7	13.2	38.4	134.0
2000 .....	7.2	16.8	2.1	0.9	2.8	3.8	5.4	6.6	8.3	13.0	36.9	135.2
1999 .....	7.6	16.2	2.3	1.1	2.9	3.7	5.3	6.6	8.3	14.8	40.9	152.7
Female												
2002 .....	5.0	14.6	1.2	0.4	0.5	0.6	1.2	1.5	2.7	6.8	24.0	100.0
2001 .....	5.3	14.2	1.5	0.4	0.6	0.6	1.1	1.9	3.1	7.3	25.4	107.0
2000 .....	5.2	13.2	1.4	0.4	0.5	0.6	1.1	1.5	3.0	7.4	25.1	109.8
1999 .....	5.6	11.3	1.5	0.5	0.5	0.7	1.2	1.7	3.1	8.0	27.7	119.3
Hispanic												
Both sexes												
2002 .....	2.5	8.2	1.3	0.5	1.5	1.9	2.5	2.8	3.4	6.2	15.3	48.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)												
Hispanic												
Both sexes												
2001 .....	2.7	9.5	1.4	0.3	1.5	1.8	2.7	3.6	3.8	7.7	16.0	63.8
2000 .....	2.7	7.5	1.4	0.4	1.6	2.2	2.9	2.7	4.0	6.2	16.6	70.3
1999 .....	2.6	8.6	1.9	0.5	1.6	1.6	2.4	2.8	4.2	8.7	15.0	65.9
Male												
2002 .....	3.6	10.1	1.8	0.6	2.5	3.3	4.2	4.8	5.6	8.4	21.3	70.3
2001 .....	3.7	9.6	1.8	*	2.6	3.0	4.6	6.1	5.6	10.8	20.6	68.6
2000 .....	3.8	7.9	1.5	*	2.9	3.7	5.2	4.5	6.6	8.7	19.7	88.7
1999 .....	3.6	9.0	2.5	*	2.6	2.7	4.0	4.8	6.2	13.2	15.6	84.1
Female												
2002 .....	1.2	6.1	*	*	*	*	*	*	*	4.4	11.2	37.3
2001 .....	1.6	9.5	*	*	*	*	*	1.2	2.1	5.4	13.0	61.4
2000 .....	1.4	7.2	*	*	*	*	*	*	*	4.3	14.5	61.3
1999 .....	1.6	8.1	*	*	*	*	*	*	2.4	5.2	14.6	57.2
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	6.5	18.9	1.6	0.7	1.6	2.1	3.1	3.8	5.3	10.0	29.4	107.6
2001 .....	6.7	18.3	1.9	0.7	1.6	2.0	3.1	4.2	5.4	10.1	31.1	116.6
2000 .....	6.7	16.9	1.8	0.7	1.7	2.2	3.2	4.1	5.7	10.1	30.2	118.5
1999 .....	7.1	15.0	1.9	0.8	1.7	2.3	3.3	4.2	5.6	11.1	33.6	130.7
Male												
2002 .....	7.5	21.0	2.0	0.9	2.6	3.4	4.9	6.1	8.1	13.7	36.6	120.4
2001 .....	7.7	21.1	2.1	1.0	2.6	3.5	5.1	6.6	7.8	13.3	39.0	136.5
2000 .....	7.7	19.0	2.2	1.0	2.7	3.8	5.3	6.7	8.4	13.2	37.5	136.9
1999 .....	8.1	17.9	2.3	1.1	2.9	3.9	5.4	6.6	8.4	14.8	41.8	154.8
Female												
2002 .....	5.6	16.8	1.3	0.4	0.5	0.7	1.2	1.6	2.8	7.0	24.6	102.2
2001 .....	5.8	15.4	1.6	0.4	0.6	0.6	1.2	2.0	3.2	7.4	25.9	108.4
2000 .....	5.7	14.7	1.4	0.4	0.5	0.7	1.1	1.6	3.1	7.5	25.4	111.1
1999 .....	6.2	12.0	1.5	0.5	0.5	0.7	1.2	1.8	3.1	8.1	28.2	121.1
Non-Hispanic white												
Both sexes												
2002 .....	7.0	14.5	1.5	0.7	1.7	2.3	3.1	3.8	5.2	9.6	29.4	110.9
2001 .....	7.2	14.5	1.7	0.7	1.8	2.1	3.1	4.1	5.2	9.8	31.2	119.5
2000 .....	7.1	13.7	1.7	0.7	1.7	2.3	3.2	3.9	5.4	9.7	30.3	122.0
1999 .....	7.6	12.1	1.7	0.9	1.9	2.4	3.4	3.9	5.2	10.6	33.1	134.3
Male												
2002 .....	7.9	17.0	1.8	0.9	2.9	3.7	5.1	6.0	7.7	13.0	36.3	123.1
2001 .....	8.1	16.8	2.0	1.0	2.9	3.6	5.1	6.3	7.4	13.0	38.6	139.7
2000 .....	8.0	16.0	2.1	1.1	2.9	4.0	5.3	6.3	7.9	12.6	36.8	139.7
1999 .....	8.4	14.2	2.0	1.2	3.1	4.0	5.5	6.2	7.7	13.9	41.0	157.3
Female												
2002 .....	6.1	11.9	1.2	0.4	0.5	0.8	1.2	1.5	2.7	6.7	24.8	105.7
2001 .....	6.4	12.1	1.5	0.4	0.7	0.6	1.1	1.9	3.1	7.1	26.3	111.2
2000 .....	6.3	11.2	1.3	0.4	0.5	0.7	1.1	1.6	3.1	7.3	26.0	114.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)												
Non-Hispanic white												
Female												
1999 .....	6.8	9.8	1.5	0.5	0.6	0.7	1.3	1.7	2.8	7.9	28.0	125.2
Non-Hispanic black												
Both sexes												
2002 .....	5.4	38.6	2.5	0.9	1.2	1.7	3.2	4.8	7.4	15.6	32.9	84.4
2001 .....	5.5	35.5	2.6	0.6	1.1	2.0	3.3	5.7	8.0	14.2	34.2	96.5
2000 .....	5.6	31.7	2.3	0.7	1.6	2.3	3.6	5.8	8.4	15.1	34.1	90.9
1999 .....	6.3	29.5	2.6	0.7	1.3	2.5	3.5	6.7	10.2	18.2	43.1	106.7
Male												
2002 .....	6.8	40.0	3.3	1.2	1.8	2.9	4.8	7.6	12.1	22.1	46.6	107.2
2001 .....	7.0	41.8	2.9	0.9	1.6	3.3	5.2	9.2	12.2	19.9	49.5	115.5
2000 .....	7.5	32.8	2.7	1.0	2.6	3.8	6.1	10.3	14.2	20.8	53.7	124.5
1999 .....	8.3	35.0	3.5	1.0	2.4	4.1	5.9	11.2	16.0	27.4	57.8	155.7
Female												
2002 .....	4.2	37.2	*	*	*	*	1.8	2.4	3.6	11.0	25.2	75.7
2001 .....	4.2	28.9	2.3	*	*	0.7	1.6	2.7	4.7	10.3	25.6	89.4
2000 .....	3.9	30.7	2.0	*	*	0.9	1.3	1.9	3.7	11.1	23.1	78.5
1999 .....	4.5	23.9	*	*	*	1.1	1.4	2.8	5.5	11.8	34.8	88.4
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	11.0	...	...	0.6	9.9	12.6	15.3	15.7	13.6	13.5	17.7	18.0
2001 <sup>5</sup> .....	10.8	...	...	0.7	9.9	12.8	14.7	15.2	13.1	13.3	17.4	17.5
2000 .....	10.4	...	...	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6
1999 .....	10.5	...	...	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3
Male												
2002 .....	17.9	...	...	0.9	16.5	20.5	23.7	24.4	22.2	24.7	38.1	50.7
2001 <sup>5</sup> .....	17.6	...	...	1.0	16.6	21.0	23.1	23.4	21.1	24.6	37.8	51.1
2000 .....	17.1	...	...	1.2	17.1	19.6	22.8	22.4	19.4	22.7	38.6	57.5
1999 .....	17.1	...	...	0.9	16.8	20.7	22.4	21.5	19.8	24.7	38.8	57.1
Female												
2002 .....	4.3	...	...	0.3	2.9	4.6	6.9	7.4	5.7	4.1	4.2	3.8
2001 <sup>5</sup> .....	4.1	...	...	0.3	2.9	4.4	6.4	7.2	5.7	3.9	4.0	3.4
2000 .....	4.0	...	...	0.3	3.0	4.3	6.4	6.7	5.4	4.0	4.0	4.2
1999 .....	4.0	...	...	0.3	3.0	4.6	6.4	6.6	5.2	4.1	4.7	4.1
Hispanic												
Both sexes												
2002 .....	5.0	...	...	0.5	6.7	6.6	7.0	7.1	7.2	7.5	9.2	12.7
2001 <sup>5</sup> .....	5.0	...	...	0.5	6.2	7.1	7.3	7.0	6.2	7.6	8.8	13.8
2000 .....	5.1	...	...	0.4	6.8	6.5	7.5	7.8	6.0	7.8	11.7	*
1999 .....	5.0	...	...	0.5	6.2	6.7	7.5	7.1	6.8	7.0	13.1	15.4
Male												
2002 .....	8.3	...	...	0.6	10.6	10.7	11.2	11.3	13.2	14.7	19.3	35.1
2001 <sup>5</sup> .....	8.3	...	...	0.7	9.5	11.3	12.4	11.4	11.5	15.4	20.1	41.9

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Hispanic												
Male												
2000 .....	8.4	...	...	0.6	10.9	10.6	12.1	12.5	11.1	15.6	26.1	*
1999 .....	8.2	...	...	0.6	9.9	10.7	12.1	11.6	12.2	13.8	28.2	*
Female												
2002 .....	1.6	...	...	*	2.1	1.7	2.4	2.9	*	*	*	*
2001 <sup>5</sup> .....	1.5	...	...	*	2.3	2.2	1.8	2.7	*	*	*	*
2000 .....	1.5	...	...	*	2.0	1.7	2.6	3.2	*	*	*	*
1999 .....	1.6	...	...	*	1.8	2.0	2.6	2.7	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	11.8	...	...	0.7	10.5	13.9	16.4	16.5	14.0	13.8	18.1	18.1
2001 <sup>5</sup> .....	11.6	...	...	0.7	10.7	13.9	15.7	15.9	13.6	13.6	17.8	17.5
2000 .....	11.1	...	...	0.8	10.8	13.0	15.4	15.0	12.5	12.7	17.9	19.9
1999 .....	11.2	...	...	0.6	10.8	13.8	15.1	14.4	12.5	13.7	18.3	19.3
Male												
2002 .....	19.4	...	...	1.0	17.6	22.7	25.5	25.6	22.8	25.2	39.0	50.9
2001 <sup>5</sup> .....	19.0	...	...	1.1	18.1	23.1	24.6	24.5	21.7	25.0	38.5	51.3
2000 .....	18.3	...	...	1.3	18.3	21.4	24.1	23.2	19.9	23.1	39.1	58.4
1999 .....	18.4	...	...	1.0	18.2	22.7	23.6	22.2	20.2	25.2	39.1	57.6
Female												
2002 .....	4.6	...	...	0.3	3.1	5.2	7.4	7.8	5.9	4.3	4.3	3.9
2001 <sup>5</sup> .....	4.5	...	...	0.3	3.0	4.9	7.0	7.6	6.1	4.1	4.2	3.5
2000 .....	4.3	...	...	0.3	3.1	4.8	6.8	7.0	5.7	4.1	4.1	4.3
1999 .....	4.3	...	...	0.2	3.2	5.0	6.8	6.9	5.4	4.3	4.8	4.0
Non-Hispanic white												
Both sexes												
2002 .....	13.4	...	...	0.7	11.5	15.4	18.7	18.7	15.6	15.1	19.4	19.2
2001 <sup>5</sup> .....	13.0	...	...	0.7	11.6	15.4	17.7	17.9	15.0	14.7	19.0	18.7
2000 .....	12.5	...	...	0.9	11.6	14.2	17.2	16.7	13.9	13.8	19.0	21.3
1999 .....	12.4	...	...	0.7	11.5	15.0	16.9	16.0	13.7	14.8	19.5	20.8
Male												
2002 .....	21.9	...	...	1.1	19.3	24.6	28.7	28.6	25.0	27.5	41.7	54.3
2001 <sup>5</sup> .....	21.4	...	...	1.1	19.6	25.2	27.4	27.4	23.6	26.9	41.1	55.2
2000 .....	20.4	...	...	1.4	19.5	23.1	26.6	25.5	21.8	24.7	41.7	62.7
1999 .....	20.3	...	...	1.1	19.2	24.3	26.0	24.4	21.9	27.0	41.6	62.2
Female												
2002 .....	5.3	...	...	0.3	3.4	6.1	8.6	9.0	6.6	4.6	4.5	4.1
2001 <sup>5</sup> .....	5.0	...	...	0.3	3.3	5.5	8.1	8.6	6.8	4.4	4.4	3.5
2000 .....	4.9	...	...	0.4	3.3	5.2	7.8	7.9	6.4	4.6	4.2	4.6
1999 .....	4.9	...	...	0.3	3.4	5.5	7.8	7.8	6.0	4.6	5.0	4.2
Non-Hispanic black												
Both sexes												
2002 .....	5.2	...	...	0.5	6.6	9.6	7.7	5.9	4.9	4.9	5.4	*
2001 <sup>5</sup> .....	5.3	...	...	0.6	7.3	9.2	7.5	6.2	5.1	5.5	5.6	*
2000 .....	5.5	...	...	0.6	8.4	9.0	7.7	6.5	4.2	5.3	5.9	*
1999 .....	5.5	...	...	0.5	8.3	9.5	7.8	5.7	5.0	5.8	5.8	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Non-Hispanic black												
Male												
2002 .....	9.3	...	...	0.9	11.6	18.0	13.2	10.2	8.5	9.7	13.5	*
2001 <sup>5</sup> .....	9.3	...	...	0.9	13.4	16.6	12.5	9.8	9.3	10.8	13.2	*
2000 .....	9.6	...	...	1.0	14.6	16.1	13.3	11.1	7.5	11.4	12.3	*
1999 .....	9.9	...	...	0.7	14.8	17.5	13.7	10.1	9.2	11.7	14.1	*
Female												
2002 .....	1.6	...	...	*	1.7	2.1	2.9	2.2	2.0	*	*	*
2001 <sup>5</sup> .....	1.7	...	...	*	1.3	2.4	3.0	3.1	1.8	*	*	*
2000 .....	1.7	...	...	*	2.3	2.6	2.7	2.5	1.6	*	*	*
1999 .....	1.6	...	...	*	2.0	2.4	2.6	2.0	1.6	*	*	*
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	5.9	...	...	0.2	5.1	6.0	7.0	7.8	8.4	9.7	13.2	12.0
2001 .....	5.9	...	...	0.2	5.3	6.5	6.7	7.7	8.2	9.6	13.1	12.2
2000 .....	5.9	...	...	0.3	5.8	6.1	7.1	7.6	7.6	9.2	13.0	13.2
1999 .....	5.9	...	...	0.3	6.0	6.5	6.9	7.3	7.8	9.7	13.1	12.6
Male												
2002 .....	10.6	...	...	0.3	9.1	10.3	11.9	13.6	15.1	19.5	31.0	37.8
2001 .....	10.6	...	...	0.3	9.4	11.2	11.4	13.3	14.6	19.2	30.7	38.9
2000 .....	10.5	...	...	0.4	10.2	10.6	11.8	13.0	13.5	18.4	30.8	42.9
1999 .....	10.6	...	...	0.4	10.5	11.2	11.5	12.5	14.0	19.5	31.1	41.6
Female												
2002 .....	1.4	...	...	*	1.0	1.6	2.2	2.2	2.2	1.6	1.4	0.8
2001 .....	1.5	...	...	0.1	1.0	1.6	2.1	2.4	2.4	1.7	1.5	1.0
2000 .....	1.5	...	...	0.1	1.2	1.7	2.4	2.3	2.1	1.7	1.4	1.1
1999 .....	1.5	...	...	0.1	1.2	1.8	2.3	2.3	2.1	1.7	1.4	0.9
Hispanic												
Both sexes												
2002 .....	2.2	...	...	*	3.3	2.7	2.5	2.7	3.4	4.4	4.7	*
2001 .....	2.2	...	...	*	2.9	3.3	2.5	3.0	3.2	4.0	4.7	*
2000 .....	2.3	...	...	*	3.4	2.8	2.8	4.0	2.7	4.6	5.3	*
1999 .....	2.3	...	...	*	3.5	2.9	3.0	3.2	2.9	3.6	8.1	*
Male												
2002 .....	3.8	...	...	*	5.7	4.6	4.2	4.8	6.6	9.4	11.5	*
2001 .....	3.8	...	...	*	4.8	5.5	4.5	5.3	6.6	8.8	11.6	*
2000 .....	4.1	...	...	*	5.8	4.7	4.7	6.8	5.7	9.7	12.8	*
1999 .....	4.0	...	...	*	5.8	4.8	5.0	5.5	5.9	7.7	19.8	*
Female												
2002 .....	0.4	...	...	*	*	*	*	*	*	*	*	*
2001 .....	0.4	...	...	*	0.7	0.7	*	*	*	*	*	*
2000 .....	0.4	...	...	*	*	*	*	*	*	*	*	*
1999 .....	0.5	...	...	*	0.8	*	0.9	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	6.5	...	...	0.2	5.5	6.7	7.7	8.3	8.8	10.0	13.6	12.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Non-Hispanic <sup>4</sup>												
Both sexes												
2001 .....	6.5	...	...	0.2	5.8	7.1	7.3	8.1	8.6	9.9	13.5	12.4
2000 .....	6.4	...	...	0.3	6.2	6.8	7.6	7.9	7.9	9.5	13.4	13.4
1999 .....	6.4	...	...	0.3	6.5	7.2	7.3	7.7	8.2	10.1	13.3	12.7
Male												
2002 .....	11.7	...	...	0.4	9.8	11.6	13.1	14.4	15.7	20.1	31.9	38.4
2001 .....	11.6	...	...	0.4	10.4	12.5	12.3	14.0	15.1	19.8	31.5	39.7
2000 .....	11.4	...	...	0.5	11.1	11.7	12.7	13.5	14.0	18.9	31.6	43.8
1999 .....	11.5	...	...	0.4	11.5	12.5	12.3	13.1	14.5	20.1	31.5	42.2
Female												
2002 .....	1.6	...	...	*	1.1	1.9	2.4	2.4	2.3	1.6	1.4	0.8
2001 .....	1.6	...	...	0.1	1.1	1.8	2.4	2.5	2.5	1.8	1.6	1.0
2000 .....	1.6	...	...	0.1	1.3	1.9	2.6	2.4	2.3	1.8	1.5	1.1
1999 .....	1.6	...	...	*	1.3	1.9	2.4	2.4	2.3	1.8	1.5	0.9
Non-Hispanic white												
Both sexes												
2002 .....	7.5	...	...	0.2	5.9	7.5	8.8	9.5	9.9	11.2	14.7	13.1
2001 .....	7.4	...	...	0.3	6.2	8.0	8.3	9.3	9.6	10.9	14.5	13.4
2000 .....	7.3	...	...	0.4	6.7	7.4	8.6	8.9	8.7	10.4	14.5	14.5
1999 .....	7.3	...	...	0.3	6.8	7.9	8.3	8.6	9.0	11.0	14.3	13.7
Male												
2002 .....	13.4	...	...	0.4	10.5	12.6	14.9	16.3	17.4	22.2	34.3	41.3
2001 .....	13.2	...	...	0.4	11.0	13.8	13.9	15.9	16.7	21.6	33.8	43.0
2000 .....	12.9	...	...	0.6	11.8	12.7	14.3	15.1	15.3	20.4	34.2	47.6
1999 .....	12.9	...	...	0.5	12.0	13.4	13.7	14.6	15.9	21.8	33.8	45.7
Female												
2002 .....	1.8	...	...	*	1.2	2.3	2.8	2.8	2.7	1.8	1.6	0.9
2001 .....	1.9	...	...	*	1.2	2.1	2.7	2.9	2.9	1.9	1.7	1.1
2000 .....	1.9	...	...	*	1.4	2.1	3.0	2.8	2.6	2.0	1.5	1.2
1999 .....	1.9	...	...	*	1.4	2.3	2.9	2.7	2.6	2.0	1.6	1.0
Non-Hispanic black												
Both sexes												
2002 .....	2.9	...	...	*	4.2	5.3	3.6	2.9	2.8	3.3	4.3	*
2001 .....	3.0	...	...	*	5.1	5.0	3.5	2.8	2.7	3.8	5.1	*
2000 .....	3.1	...	...	*	5.4	5.2	3.5	3.2	3.0	3.8	3.6	*
1999 .....	3.2	...	...	*	5.8	5.1	3.6	3.0	3.5	4.0	4.8	*
Male												
2002 .....	5.5	...	...	*	7.7	10.5	6.5	5.5	5.5	7.0	11.4	*
2001 .....	5.6	...	...	*	9.6	9.5	6.4	4.9	5.5	7.9	12.0	*
2000 .....	5.7	...	...	*	9.9	9.6	6.4	5.6	5.7	9.0	8.3	*
1999 .....	6.1	...	...	*	10.6	10.0	6.9	5.6	7.1	8.9	12.2	*
Female												
2002 .....	0.5	...	...	*	0.7	*	1.0	*	*	*	*	*
2001 .....	0.6	...	...	*	*	0.9	1.0	1.0	*	*	*	*
2000 .....	0.7	...	...	*	1.0	1.1	0.8	1.1	*	*	*	*
1999 .....	0.6	...	...	*	1.0	0.8	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	5.0	...	...	0.4	4.7	6.6	8.2	7.9	5.2	3.8	4.5	6.0
2001 <sup>5</sup> .....	4.8	...	...	0.5	4.6	6.3	8.0	7.4	4.9	3.7	4.3	5.3
2000 .....	4.5	...	...	0.5	4.4	5.9	7.5	6.8	4.6	3.2	4.6	6.5
1999 .....	4.5	...	...	0.3	4.1	6.2	7.5	6.6	4.4	3.7	5.0	6.7
Male												
2002 .....	7.3	...	...	0.6	7.4	10.2	11.8	10.8	7.1	5.2	7.1	12.9
2001 <sup>5</sup> .....	7.1	...	...	0.7	7.2	9.8	11.7	10.2	6.5	5.4	7.1	12.2
2000 .....	6.6	...	...	0.7	6.9	9.0	10.9	9.4	6.0	4.3	7.8	14.6
1999 .....	6.6	...	...	0.5	6.3	9.5	10.9	8.9	5.8	5.3	7.7	15.5
Female												
2002 .....	2.9	...	...	0.2	2.0	3.0	4.7	5.2	3.5	2.5	2.8	3.0
2001 <sup>5</sup> .....	2.6	...	...	0.2	1.9	2.8	4.3	4.8	3.4	2.3	2.5	2.4
2000 .....	2.5	...	...	0.2	1.8	2.7	4.0	4.4	3.3	2.3	2.6	3.2
1999 .....	2.5	...	...	0.1	1.8	2.8	4.1	4.3	3.0	2.4	3.3	3.1
Hispanic												
Both sexes												
2002 .....	2.9	...	...	0.4	3.4	3.9	4.5	4.4	3.8	3.1	4.5	*
2001 <sup>5</sup> .....	2.8	...	...	0.5	3.3	3.9	4.8	4.0	3.0	3.6	4.1	*
2000 .....	2.8	...	...	0.3	3.4	3.7	4.7	3.8	3.2	3.3	6.3	*
1999 .....	2.7	...	...	*	2.7	3.8	4.5	3.8	3.9	3.5	5.0	*
Male												
2002 .....	4.4	...	...	*	4.9	6.1	7.0	6.5	6.5	5.4	*	*
2001 <sup>5</sup> .....	4.4	...	...	0.6	4.7	5.8	7.9	6.1	4.9	6.6	*	*
2000 .....	4.3	...	...	*	5.1	5.9	7.3	5.7	5.4	5.9	13.3	*
1999 .....	4.2	...	...	*	4.2	5.9	7.1	6.0	6.4	6.2	*	*
Female												
2002 .....	1.2	...	...	*	1.6	1.3	1.8	2.4	*	*	*	*
2001 <sup>5</sup> .....	1.2	...	...	*	1.6	1.6	1.4	2.0	*	*	*	*
2000 .....	1.1	...	...	*	1.4	1.1	1.9	2.0	*	*	*	*
1999 .....	1.1	...	...	*	1.0	1.4	1.7	1.7	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	5.3	...	...	0.4	5.0	7.2	8.7	8.2	5.3	3.8	4.5	5.9
2001 <sup>5</sup> .....	5.1	...	...	0.5	4.8	6.8	8.4	7.7	5.0	3.7	4.3	5.1
2000 .....	4.8	...	...	0.5	4.6	6.2	7.8	7.1	4.6	3.2	4.5	6.5
1999 .....	4.7	...	...	0.4	4.4	6.6	7.8	6.8	4.4	3.7	5.0	6.6
Male												
2002 .....	7.7	...	...	0.7	7.8	11.1	12.4	11.1	7.0	5.1	7.1	12.5
2001 <sup>5</sup> .....	7.5	...	...	0.7	7.7	10.6	12.2	10.5	6.6	5.3	7.0	11.6
2000 .....	6.9	...	...	0.8	7.2	9.6	11.4	9.7	6.0	4.2	7.5	14.6
1999 .....	6.9	...	...	0.6	6.7	10.2	11.3	9.1	5.7	5.1	7.6	15.4
Female												
2002 .....	3.1	...	...	0.2	2.0	3.3	5.1	5.4	3.6	2.6	2.9	3.1
2001 <sup>5</sup> .....	2.9	...	...	0.2	1.9	3.1	4.7	5.1	3.5	2.3	2.6	2.5

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
Non-Hispanic <sup>4</sup>												
Female												
2000 .....	2.7	...	...	0.2	1.9	2.9	4.3	4.6	3.4	2.4	2.6	3.2
1999 .....	2.7	...	...	0.1	1.9	3.0	4.4	4.5	3.1	2.5	3.3	3.1
Non-Hispanic white												
Both sexes												
2002 .....	6.0	...	...	0.5	5.6	7.9	9.8	9.2	5.7	3.9	4.7	6.1
2001 <sup>5</sup> .....	5.6	...	...	0.4	5.4	7.5	9.4	8.6	5.3	3.8	4.5	5.3
2000 .....	5.2	...	...	0.5	4.9	6.8	8.6	7.8	5.1	3.3	4.5	6.8
1999 .....	5.2	...	...	0.4	4.6	7.1	8.6	7.4	4.6	3.8	5.1	7.0
Male												
2002 .....	8.5	...	...	0.7	8.8	12.0	13.8	12.3	7.6	5.3	7.3	13.0
2001 <sup>5</sup> .....	8.2	...	...	0.7	8.6	11.5	13.5	11.5	6.9	5.3	7.3	12.3
2000 .....	7.5	...	...	0.8	7.7	10.3	12.4	10.4	6.5	4.3	7.5	15.2
1999 .....	7.4	...	...	0.6	7.2	10.8	12.3	9.8	6.0	5.3	7.8	16.5
Female												
2002 .....	3.5	...	...	0.2	2.2	3.8	5.8	6.1	3.9	2.7	2.9	3.2
2001 <sup>5</sup> .....	3.2	...	...	0.2	2.0	3.4	5.3	5.7	3.9	2.4	2.7	2.4
2000 .....	3.0	...	...	0.2	1.9	3.1	4.8	5.2	3.8	2.5	2.6	3.4
1999 .....	3.0	...	...	*	2.0	3.3	4.9	5.1	3.4	2.6	3.4	3.2
Non-Hispanic black												
Both sexes												
2002 .....	2.4	...	...	0.4	2.4	4.4	4.2	3.0	2.1	1.6	*	*
2001 <sup>5</sup> .....	2.3	...	...	0.4	2.3	4.2	3.9	3.4	2.4	1.7	*	*
2000 .....	2.4	...	...	0.5	3.0	3.9	4.2	3.3	1.3	1.4	2.3	*
1999 .....	2.3	...	...	0.4	2.5	4.4	4.2	2.7	1.5	1.8	*	*
Male												
2002 .....	3.8	...	...	0.6	3.8	7.5	6.7	4.7	3.0	*	*	*
2001 <sup>5</sup> .....	3.7	...	...	0.7	3.8	7.1	6.1	4.9	3.7	3.0	*	*
2000 .....	3.9	...	...	0.9	4.7	6.5	6.8	5.4	*	*	*	*
1999 .....	3.8	...	...	*	4.1	7.5	6.8	4.5	2.1	*	*	*
Female												
2002 .....	1.1	...	...	*	1.0	1.6	2.0	1.6	1.4	*	*	*
2001 <sup>5</sup> .....	1.1	...	...	*	0.7	1.5	2.0	2.1	*	*	*	*
2000 .....	1.1	...	...	*	1.3	1.5	1.9	1.5	*	*	*	*
1999 .....	1.0	...	...	*	0.9	1.6	2.0	1.3	*	*	*	*
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	6.1	7.5	2.7	0.9	12.9	11.2	7.2	4.8	3.2	2.3	2.3	2.1
2001 <sup>5</sup> .....	7.1	8.2	2.7	0.8	13.3	13.1	9.5	6.3	4.0	2.9	2.5	2.4
2000 .....	6.0	9.2	2.3	0.9	12.6	10.4	7.1	4.7	3.0	2.4	2.4	2.4
1999 .....	6.1	8.7	2.5	1.1	12.9	10.5	7.1	4.6	3.0	2.6	2.5	2.4
Male												
2002 .....	9.6	7.9	2.9	0.9	21.5	18.2	10.7	7.1	4.8	3.3	3.1	3.0

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
All origins <sup>3</sup>												
Male												
2001 <sup>5</sup> .....	11.1	9.5	3.0	0.9	22.2	20.8	13.9	9.5	5.9	4.0	3.1	3.2
2000 .....	9.3	10.4	2.5	1.1	20.9	16.7	10.3	6.9	4.6	3.4	3.2	3.3
1999 .....	9.3	9.6	2.5	1.2	21.1	16.5	10.3	6.8	4.7	3.7	3.4	3.4
Female												
2002 .....	2.7	7.1	2.5	0.8	3.8	4.2	3.8	2.6	1.6	1.4	1.8	1.7
2001 <sup>5</sup> .....	3.3	6.9	2.4	0.7	3.9	5.3	5.1	3.2	2.2	2.0	2.1	2.0
2000 .....	2.8	7.9	2.1	0.7	3.9	4.1	4.0	2.5	1.6	1.6	2.0	2.0
1999 .....	2.9	7.8	2.4	0.9	4.3	4.4	4.0	2.4	1.5	1.7	1.9	2.0
Hispanic												
Both sexes												
2002 .....	8.1	6.2	2.6	0.9	17.6	12.3	8.0	5.9	4.4	2.5	*	*
2001 <sup>5</sup> .....	9.0	6.7	2.3	0.8	18.3	14.0	10.4	7.3	4.4	4.0	*	*
2000 .....	8.3	7.0	1.9	1.1	17.1	12.4	8.7	6.7	4.5	3.3	*	*
1999 .....	8.4	7.0	1.8	1.4	17.7	12.7	8.8	6.6	4.2	3.2	*	*
Male												
2002 .....	13.2	6.6	3.1	1.0	29.6	19.8	12.1	9.1	7.7	4.8	*	*
2001 <sup>5</sup> .....	14.5	7.9	2.5	1.1	30.5	21.6	15.9	11.3	6.8	6.6	*	*
2000 .....	13.4	6.6	2.0	1.6	28.5	19.9	13.5	10.2	7.1	4.6	*	*
1999 .....	13.8	6.7	1.7	1.8	29.1	20.6	13.9	10.3	7.3	5.7	*	*
Female												
2002 .....	2.6	5.9	2.1	0.8	3.8	3.4	3.5	2.7	*	*	*	*
2001 <sup>5</sup> .....	3.2	5.5	2.1	*	4.0	5.1	4.4	3.4	2.4	*	*	*
2000 .....	2.8	7.4	1.9	0.6	3.7	3.7	3.6	3.3	2.2	*	*	*
1999 .....	2.8	7.3	1.9	0.9	4.4	3.5	3.3	3.1	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	5.7	7.7	2.7	0.9	11.8	10.9	7.0	4.6	3.0	2.3	2.2	2.1
2001 <sup>5</sup> .....	6.7	8.4	2.8	0.8	12.1	12.8	9.2	6.1	3.9	2.8	2.5	2.4
2000 .....	5.6	9.6	2.4	0.9	11.6	10.0	6.9	4.4	2.9	2.3	2.4	2.3
1999 .....	5.7	8.9	2.6	1.0	11.9	10.0	6.8	4.3	2.9	2.5	2.5	2.4
Male												
2002 .....	8.9	8.0	2.9	0.9	19.6	17.6	10.4	6.8	4.5	3.2	3.1	2.9
2001 <sup>5</sup> .....	10.4	9.8	3.2	0.9	20.1	20.3	13.4	9.2	5.8	3.8	3.1	3.1
2000 .....	8.6	11.2	2.6	1.0	19.1	15.9	9.8	6.6	4.4	3.2	3.1	3.1
1999 .....	8.6	10.1	2.6	1.1	19.4	15.6	9.7	6.4	4.5	3.5	3.3	3.4
Female												
2002 .....	2.7	7.4	2.6	0.8	3.7	4.3	3.8	2.5	1.7	1.5	1.7	1.7
2001 <sup>5</sup> .....	3.2	6.9	2.4	0.7	3.8	5.3	5.1	3.1	2.2	2.0	2.1	2.1
2000 .....	2.7	7.9	2.2	0.7	3.8	4.1	4.0	2.3	1.5	1.6	1.9	2.0
1999 .....	2.9	7.6	2.5	0.9	4.2	4.6	4.0	2.3	1.4	1.7	2.0	2.0
Non-Hispanic white												
Both sexes												
2002 .....	2.8	5.0	1.8	0.5	3.7	3.9	4.1	3.0	2.4	1.9	1.8	1.8
2001 <sup>5</sup> .....	4.0	5.8	1.7	0.6	4.2	6.5	6.6	4.6	3.3	2.3	2.1	2.2
2000 .....	2.8	6.2	1.6	0.6	3.5	4.0	4.0	2.9	2.2	1.8	2.0	2.1
1999 .....	2.9	6.3	1.7	0.7	3.9	4.3	4.1	2.9	2.2	2.1	2.0	2.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Non-Hispanic white												
Male												
2002 .....	3.8	5.8	1.9	0.5	5.2	5.3	5.6	3.9	3.3	2.4	2.3	2.7
2001 <sup>5</sup> .....	5.6	6.8	2.0	0.6	5.7	9.3	9.5	6.7	4.7	2.9	2.6	2.8
2000 .....	3.6	8.3	1.8	0.7	4.7	5.2	5.2	3.9	3.1	2.2	2.4	2.6
1999 .....	3.9	8.0	1.7	0.7	5.1	5.7	5.4	4.2	3.2	2.7	2.6	2.6
Female												
2002 .....	1.9	4.1	1.8	0.6	2.2	2.6	2.5	2.1	1.5	1.4	1.5	1.5
2001 <sup>5</sup> .....	2.4	4.7	1.5	0.7	2.7	3.7	3.7	2.6	1.9	1.8	1.8	2.0
2000 .....	1.9	4.1	1.3	0.6	2.3	2.7	2.7	1.9	1.3	1.4	1.7	1.9
1999 .....	2.0	4.4	1.6	0.7	2.6	2.9	2.8	1.7	1.3	1.6	1.6	1.9
Non-Hispanic black												
Both sexes												
2002 .....	22.5	17.6	6.5	2.0	48.3	47.0	24.3	15.8	8.4	5.8	7.3	*
2001 <sup>5</sup> .....	22.7	19.5	7.1	1.4	48.7	45.3	25.5	16.2	8.9	7.2	6.8	*
2000 .....	22.1	24.1	6.1	1.9	49.3	42.2	24.5	15.0	8.4	7.6	7.1	*
1999 .....	21.7	20.0	6.4	2.3	49.5	40.6	23.7	14.1	8.4	6.4	7.5	*
Male												
2002 .....	39.4	16.4	7.3	2.6	85.8	85.3	39.8	27.5	15.8	10.5	13.8	*
2001 <sup>5</sup> .....	39.3	21.7	8.4	2.0	88.3	81.7	39.5	27.2	15.0	12.6	10.2	*
2000 .....	38.3	24.9	6.1	2.1	88.1	76.5	39.4	26.3	15.0	13.9	12.0	*
1999 .....	37.2	20.1	6.2	2.7	88.1	71.5	37.9	23.7	15.4	11.9	12.2	*
Female												
2002 .....	7.2	18.9	5.7	1.5	10.6	12.3	10.6	5.7	2.5	2.4	3.6	*
2001 <sup>5</sup> .....	7.6	17.1	5.7	0.8	9.2	12.5	13.2	6.9	4.0	3.3	4.9	*
2000 .....	7.4	23.3	6.0	1.6	11.0	11.4	11.3	5.3	3.2	3.2	4.3	*
1999 .....	7.7	20.0	6.7	1.8	11.4	12.8	11.1	5.9	2.7	2.5	4.9	*
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	4.1	*	0.3	0.5	10.6	8.7	4.5	2.6	1.5	0.9	0.9	*
2001 .....	4.0	*	0.4	0.4	10.5	8.4	4.4	2.4	1.4	1.0	0.8	*
2000 .....	3.8	*	0.2	0.5	10.1	7.7	4.2	2.5	1.5	1.0	0.7	0.7
1999 .....	3.9	*	0.3	0.5	10.4	7.6	4.1	2.4	1.6	1.0	0.8	0.6
Male												
2002 .....	7.0	*	0.4	0.6	18.5	14.9	7.3	3.9	2.4	1.2	1.3	*
2001 .....	6.8	*	0.3	0.6	18.5	14.3	6.8	3.8	2.2	1.5	1.1	*
2000 .....	6.5	*	*	0.7	17.7	13.2	6.6	3.9	2.3	1.3	0.8	*
1999 .....	6.5	*	0.3	0.7	18.0	12.9	6.3	3.8	2.6	1.5	1.0	*
Female												
2002 .....	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001 .....	1.3	*	0.4	0.3	2.1	2.3	2.0	1.0	0.8	0.7	0.6	*
2000 .....	1.3	*	*	0.2	2.2	2.1	1.8	1.2	0.8	0.8	0.6	*
1999 .....	1.3	*	0.3	0.3	2.5	2.2	2.0	1.2	0.6	0.7	0.6	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Hispanic												
Both sexes												
2002 .....	5.6	*	*	0.6	14.3	9.3	5.1	3.2	1.6	*	*	*
2001 .....	5.7	*	*	0.4	14.6	9.5	5.6	3.0	1.1	*	*	*
2000 .....	5.5	*	*	0.6	13.7	8.9	4.9	4.0	2.0	*	*	*
1999 .....	5.7	*	*	0.8	14.4	8.7	5.2	3.8	2.1	*	*	*
Male												
2002 .....	9.7	*	*	0.8	24.9	15.6	8.1	5.0	2.8	*	*	*
2001 .....	9.9	*	*	0.6	24.9	16.1	9.1	4.8	*	*	*	*
2000 .....	9.6	*	*	1.0	23.5	15.0	7.8	6.4	3.2	*	*	*
1999 .....	9.8	*	*	1.2	24.2	14.7	8.6	5.7	3.5	*	*	*
Female												
2002 .....	1.2	*	*	*	2.1	1.8	1.8	1.4	*	*	*	*
2001 .....	1.3	*	*	*	2.5	1.8	1.9	1.2	*	*	*	*
2000 .....	1.3	*	*	*	2.2	1.9	1.7	1.7	*	*	*	*
1999 .....	1.4	*	*	*	2.9	1.7	1.7	1.9	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	3.8	*	0.3	0.5	9.8	8.5	4.4	2.5	1.5	0.9	0.9	*
2001 .....	3.7	*	0.3	0.4	9.6	8.0	4.2	2.3	1.5	1.0	0.8	*
2000 .....	3.6	*	0.2	0.4	9.3	7.4	4.1	2.4	1.5	1.0	0.7	0.7
1999 .....	3.6	*	0.4	0.5	9.6	7.3	4.0	2.3	1.5	1.1	0.8	0.5
Male												
2002 .....	6.5	*	0.3	0.6	17.0	14.7	7.1	3.7	2.3	1.2	1.2	*
2001 .....	6.3	*	0.3	0.6	17.0	13.8	6.5	3.7	2.2	1.4	1.1	*
2000 .....	6.0	*	*	0.6	16.3	12.7	6.4	3.7	2.2	1.3	0.8	*
1999 .....	6.0	*	0.4	0.7	16.7	12.4	6.0	3.6	2.5	1.5	0.9	*
Female												
2002 .....	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001 .....	1.2	*	0.3	0.3	2.0	2.3	2.0	1.0	0.8	0.7	0.6	*
2000 .....	1.2	*	*	0.2	2.2	2.2	1.9	1.1	0.8	0.7	0.6	*
1999 .....	1.3	*	0.4	0.3	2.4	2.3	2.0	1.1	0.6	0.7	0.6	*
Non-Hispanic white												
Both sexes												
2002 .....	1.5	*	0.2	0.3	2.5	2.6	2.4	1.6	1.2	0.7	0.8	*
2001 .....	1.6	*	0.2	0.4	2.4	2.7	2.4	1.6	1.2	0.8	0.7	*
2000 .....	1.4	*	*	0.3	2.2	2.4	2.3	1.6	1.2	0.8	0.6	0.6
1999 .....	1.5	*	0.2	0.3	2.4	2.6	2.3	1.5	1.1	0.9	0.6	0.6
Male												
2002 .....	2.2	*	*	0.3	3.7	3.7	3.4	2.2	1.7	0.9	1.0	*
2001 .....	2.2	*	*	0.4	3.6	3.9	3.4	2.3	1.6	1.0	0.9	*
2000 .....	2.0	*	*	0.4	3.2	3.4	3.2	2.2	1.6	0.9	0.6	*
1999 .....	2.2	*	*	0.4	3.5	3.9	3.2	2.2	1.7	1.2	0.6	*
Female												
2002 .....	0.9	*	*	0.3	1.2	1.4	1.3	1.1	0.7	0.6	0.7	*
2001 .....	0.9	*	*	0.3	1.2	1.5	1.4	0.8	0.8	0.7	0.6	*
2000 .....	0.9	*	*	0.2	1.2	1.4	1.4	0.9	0.8	0.7	0.6	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Non-Hispanic white												
Female												
1999 .....	0.9	*	*	0.2	1.2	1.4	1.5	0.8	0.6	0.7	0.6	*
Non-Hispanic black												
Both sexes												
2002 .....	17.1	*	*	1.2	43.1	39.7	16.7	8.6	4.2	2.4	2.1	*
2001 .....	16.3	*	*	0.9	43.0	36.4	15.5	7.7	3.7	2.6	*	*
2000 .....	16.0	*	*	1.0	42.6	34.5	15.4	8.5	3.6	2.8	*	*
1999 .....	15.9	*	*	1.3	43.6	32.4	14.4	8.1	4.5	2.4	2.4	*
Male												
2002 .....	31.9	*	*	1.6	78.6	75.6	30.0	15.3	7.7	4.7	*	*
2001 .....	30.6	*	*	1.3	80.6	69.4	26.7	14.0	7.1	5.5	*	*
2000 .....	30.1	*	*	1.6	79.1	65.7	27.4	15.3	6.5	5.5	*	*
1999 .....	29.3	*	*	1.8	80.0	61.1	24.6	14.1	9.1	4.7	*	*
Female												
2002 .....	3.7	*	*	0.8	7.5	7.2	5.0	2.8	*	*	*	*
2001 .....	3.2	*	*	*	5.6	6.5	5.5	2.2	*	*	*	*
2000 .....	3.3	*	*	*	6.7	6.4	4.7	2.6	*	*	*	*
1999 .....	3.7	*	*	0.7	7.8	6.6	5.5	3.0	*	*	*	*
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	2.0	7.3	2.4	0.4	2.2	2.6	2.7	2.2	1.6	1.4	1.4	1.7
2001 <sup>5</sup> .....	3.1	8.0	2.3	0.4	2.7	4.8	5.1	3.9	2.6	1.9	1.7	2.0
2000 .....	2.1	8.9	2.1	0.4	2.5	2.7	2.9	2.1	1.5	1.4	1.8	1.7
1999 .....	2.2	8.5	2.1	0.5	2.5	2.9	3.0	2.1	1.5	1.5	1.7	1.8
Male												
2002 .....	2.6	7.8	2.6	0.3	3.0	3.2	3.4	3.2	2.4	2.1	1.9	2.3
2001 <sup>5</sup> .....	4.3	9.2	2.7	0.3	3.6	6.5	7.1	5.7	3.8	2.5	2.0	2.7
2000 .....	2.8	10.0	2.3	0.4	3.3	3.5	3.7	3.0	2.3	2.1	2.4	2.5
1999 .....	2.8	9.4	2.1	0.5	3.1	3.6	4.0	3.1	2.2	2.2	2.4	2.6
Female												
2002 .....	1.4	6.8	2.2	0.4	1.4	1.9	1.9	1.3	0.9	0.8	1.1	1.4
2001 <sup>5</sup> .....	2.0	6.7	2.0	0.4	1.8	3.1	3.1	2.1	1.5	1.3	1.5	1.7
2000 .....	1.5	7.7	2.0	0.5	1.7	1.9	2.1	1.3	0.8	0.9	1.4	1.4
1999 .....	1.6	7.6	2.1	0.6	1.8	2.2	2.0	1.2	0.8	1.0	1.3	1.5
Hispanic												
Both sexes												
2002 .....	2.5	6.0	2.2	0.3	3.3	3.1	2.9	2.7	2.8	1.7	*	*
2001 <sup>5</sup> .....	3.3	6.2	1.9	0.4	3.7	4.5	4.8	4.3	3.4	2.7	*	*
2000 .....	2.7	6.9	1.8	0.5	3.4	3.5	3.9	2.7	2.5	2.0	*	*
1999 .....	2.7	6.7	1.6	0.6	3.3	4.0	3.5	2.9	2.1	2.3	*	*
Male												
2002 .....	3.5	6.6	2.6	*	4.7	4.3	4.0	4.1	4.9	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
Hispanic												
Male												
2001 <sup>5</sup> .....	4.5	7.2	2.2	*	5.6	5.5	6.9	6.5	5.1	4.6	*	*
2000 .....	3.8	6.6	1.8	0.6	5.0	5.0	5.7	3.8	3.9	*	*	*
1999 .....	4.0	6.2	1.5	0.6	4.9	5.9	5.3	4.5	3.8	4.4	*	*
Female												
2002 .....	1.4	5.4	1.8	*	1.7	1.6	1.7	1.4	*	*	*	*
2001 <sup>5</sup> .....	1.9	5.2	1.6	*	1.5	3.2	2.6	2.2	*	*	*	*
2000 .....	1.6	7.2	1.7	*	1.5	1.9	1.9	1.6	*	*	*	*
1999 .....	1.4	7.3	1.8	*	1.5	1.8	1.6	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.9	7.5	2.4	0.4	2.0	2.4	2.6	2.1	1.5	1.4	1.3	1.7
2001 <sup>5</sup> .....	3.1	8.2	2.4	0.4	2.5	4.7	5.0	3.8	2.5	1.8	1.7	2.1
2000 .....	2.0	9.2	2.2	0.4	2.3	2.6	2.8	2.0	1.4	1.4	1.7	1.6
1999 .....	2.1	8.7	2.2	0.5	2.3	2.7	2.9	2.0	1.4	1.5	1.7	1.8
Male												
2002 .....	2.5	7.8	2.5	0.3	2.6	2.9	3.3	3.1	2.2	2.0	1.8	2.3
2001 <sup>5</sup> .....	4.2	9.5	2.8	0.3	3.2	6.5	7.0	5.5	3.6	2.4	2.0	2.7
2000 .....	2.6	10.6	2.4	0.4	2.9	3.2	3.4	2.9	2.2	2.0	2.3	2.4
1999 .....	2.6	10.0	2.3	0.4	2.8	3.2	3.8	2.9	2.0	2.0	2.4	2.6
Female												
2002 .....	1.4	7.1	2.3	0.4	1.4	1.9	1.9	1.2	0.9	0.9	1.0	1.4
2001 <sup>5</sup> .....	2.0	6.8	2.0	0.4	1.9	2.9	3.1	2.1	1.4	1.3	1.5	1.8
2000 .....	1.5	7.8	2.0	0.5	1.7	1.9	2.1	1.2	0.7	0.9	1.3	1.3
1999 .....	1.6	7.3	2.2	0.6	1.8	2.3	2.0	1.2	0.8	1.0	1.3	1.5
Non-Hispanic white												
Both sexes												
2002 .....	1.3	5.0	1.6	0.3	1.2	1.4	1.7	1.4	1.2	1.1	1.0	1.4
2001 <sup>5</sup> .....	2.4	5.6	1.5	0.3	1.7	3.8	4.2	3.1	2.1	1.5	1.4	1.9
2000 .....	1.3	6.0	1.4	0.4	1.3	1.6	1.7	1.4	1.0	1.0	1.4	1.5
1999 .....	1.4	6.2	1.4	0.4	1.5	1.7	1.8	1.4	1.1	1.2	1.4	1.5
Male												
2002 .....	1.6	5.8	1.6	0.2	1.5	1.6	2.2	1.8	1.6	1.6	1.3	2.0
2001 <sup>5</sup> .....	3.4	6.7	1.8	0.2	2.0	5.4	6.1	4.5	3.1	1.9	1.7	2.3
2000 .....	1.6	8.1	1.7	0.3	1.5	1.8	2.0	1.8	1.5	1.3	1.8	1.9
1999 .....	1.7	7.8	1.5	0.3	1.6	1.8	2.3	2.0	1.5	1.5	2.0	*
Female												
2002 .....	1.0	4.1	1.6	0.3	0.9	1.2	1.2	1.0	0.8	0.8	0.8	1.2
2001 <sup>5</sup> .....	1.5	4.5	1.2	0.4	1.4	2.2	2.3	1.7	1.2	1.1	1.2	1.7
2000 .....	1.0	3.9	1.2	0.4	1.2	1.3	1.4	1.0	0.5	0.7	1.1	1.3
1999 .....	1.1	4.4	1.4	0.4	1.4	1.6	1.3	0.9	0.7	0.9	1.0	1.4
Non-Hispanic black												
Both sexes												
2002 .....	5.4	16.7	5.9	0.8	5.2	7.3	7.6	7.2	4.2	3.4	5.2	*
2001 <sup>5</sup> .....	6.4	19.1	6.3	0.6	5.7	9.0	10.1	8.6	5.1	4.5	4.9	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
Non-Hispanic black												
Both sexes												
2000 .....	6.1	22.9	5.7	0.8	6.7	7.8	9.1	6.5	4.9	4.8	5.2	*
1999 .....	5.8	19.7	5.7	1.0	5.9	8.2	9.2	6.0	3.9	4.0	5.2	*
Male												
2002 .....	7.6	15.8	6.6	0.9	7.3	9.7	9.9	12.3	8.1	5.9	9.4	*
2001 <sup>5</sup> .....	8.7	21.4	7.7	0.7	7.7	12.3	12.8	13.2	7.9	7.1	*	*
2000 .....	8.2	22.5	5.7	*	9.1	10.8	12.0	11.0	8.5	8.4	8.3	*
1999 .....	7.9	19.7	5.6	0.9	8.1	10.5	13.3	9.6	6.3	7.2	7.2	*
Female												
2002 .....	3.5	17.7	5.2	0.7	3.1	5.1	5.7	2.9	*	*	*	*
2001 <sup>5</sup> .....	4.4	16.8	5.0	*	3.6	5.9	7.7	4.6	2.9	2.7	4.4	*
2000 .....	4.1	23.3	5.8	1.1	4.3	5.0	6.5	2.7	2.0	2.3	3.5	*
1999 .....	4.0	19.6	5.7	1.1	3.6	6.1	5.6	2.9	1.9	*	4.0	*
Legal intervention (Y35,Y89.0)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
2001 .....	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000 .....	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
1999 .....	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
Male												
2002 .....	0.3	*	*	*	0.4	0.5	0.5	0.3	*	*	*	*
2001 .....	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2000 .....	0.3	*	*	*	0.4	0.5	0.4	0.3	*	*	*	*
1999 .....	0.3	*	*	*	0.3	0.7	0.5	0.3	0.2	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	0.2	*	*	*	0.3	0.3	*	*	*	*	*	*
2001 .....	0.2	*	*	*	0.3	0.3	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	*	0.4	*	*	*	*	*	*
Male												
2002 .....	0.3	*	*	*	0.6	0.5	*	*	*	*	*	*
2001 .....	0.3	*	*	*	0.6	0.6	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	0.7	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Hispanic												
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2001 .....	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000 .....	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
1999 .....	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
Male												
2002 .....	0.2	*	*	*	0.4	0.5	0.5	0.3	*	*	*	*
2001 .....	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2000 .....	0.2	*	*	*	0.3	0.6	0.4	0.3	*	*	*	*
1999 .....	0.3	*	*	*	0.3	0.7	0.5	0.3	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2001 .....	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2000 .....	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
1999 .....	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
Male												
2002 .....	0.2	*	*	*	0.2	0.4	0.4	0.2	*	*	*	*
2001 .....	0.2	*	*	*	0.3	0.4	0.4	0.2	*	*	*	*
2000 .....	0.2	*	*	*	0.2	0.4	0.3	0.3	*	*	*	*
1999 .....	0.2	*	*	*	0.2	0.5	0.4	0.2	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	0.3	*	*	*	0.6	0.5	0.5	*	*	*	*	*
2001 .....	0.2	*	*	*	0.4	0.4	0.4	*	*	*	*	*
2000 .....	0.3	*	*	*	0.5	0.6	0.6	*	*	*	*	*
1999 .....	0.4	*	*	*	0.6	0.8	0.7	*	*	*	*	*
Male												
2002 .....	0.6	*	*	*	1.2	1.1	1.0	*	*	*	*	*
2001 .....	0.5	*	*	*	0.8	0.8	0.8	*	*	*	*	*
2000 .....	0.6	*	*	*	1.0	1.3	1.3	*	*	*	*	*
1999 .....	0.8	*	*	*	1.1	1.7	1.4	*	*	*	*	*

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Non-Hispanic black												
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.7	2.5	0.4	0.2	1.2	2.1	3.4	2.8	1.2	0.8	0.7	1.4
2001 .....	1.5	2.1	0.3	0.1	1.0	1.8	3.1	2.5	0.9	0.5	0.9	1.0
2000 .....	1.4	1.7	0.3	0.1	0.9	1.7	2.8	2.2	0.9	0.6	0.6	1.3
1999 .....	1.4	1.4	0.3	0.1	1.1	1.8	2.9	2.2	0.9	0.6	0.8	1.4
Male												
2002 .....	2.2	2.5	0.4	0.2	1.7	2.9	4.4	3.5	1.6	1.0	0.9	2.2
2001 .....	2.0	2.4	*	0.2	1.5	2.6	4.2	3.4	1.2	0.8	1.1	1.5
2000 .....	1.9	1.4	0.3	0.2	1.4	2.4	3.9	3.1	1.3	0.8	1.0	1.7
1999 .....	2.0	1.7	0.3	0.2	1.7	2.6	4.0	3.0	1.3	0.9	1.1	2.1
Female												
2002 .....	1.2	2.5	0.4	0.1	0.7	1.2	2.4	2.2	0.8	0.6	0.6	1.1
2001 .....	0.9	1.7	0.3	*	0.5	1.1	2.1	1.6	0.6	0.3	0.7	0.8
2000 .....	0.8	2.0	*	*	0.4	1.0	1.8	1.4	0.6	0.3	0.4	1.2
1999 .....	0.9	1.2	0.3	*	0.5	1.1	1.8	1.4	0.6	0.4	0.6	1.1
Hispanic												
Both sexes												
2002 .....	0.8	*	*	*	0.7	1.0	1.5	1.3	*	*	*	*
2001 .....	0.8	*	*	*	0.5	0.9	1.7	1.6	*	*	*	*
2000 .....	0.6	*	*	*	0.5	0.9	1.2	0.8	*	*	*	*
1999 .....	0.7	*	*	*	0.8	1.1	1.2	0.9	*	*	*	*
Male												
2002 .....	1.2	*	*	*	1.2	1.5	2.1	2.2	*	*	*	*
2001 .....	1.1	*	*	*	0.7	1.4	2.6	2.1	*	*	*	*
2000 .....	0.9	*	*	*	0.8	1.4	1.8	1.3	*	*	*	*
1999 .....	1.1	*	*	*	1.3	1.6	1.9	1.6	*	*	*	*
Female												
2002 .....	0.4	*	*	*	*	*	0.9	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.8	2.7	0.4	0.2	1.3	2.3	3.6	3.0	1.2	0.7	0.8	1.4
2001 .....	1.6	2.4	0.3	0.1	1.1	2.0	3.3	2.6	0.9	0.5	0.9	0.9
2000 .....	1.5	1.9	0.3	0.2	1.0	1.8	3.0	2.4	0.9	0.6	0.6	1.4
1999 .....	1.5	1.6	0.3	0.1	1.2	2.0	3.1	2.3	0.9	0.6	0.8	1.4
Male												
2002 .....	2.3	2.8	0.3	0.2	1.8	3.2	4.7	3.6	1.6	1.0	0.9	2.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Non-Hispanic <sup>4</sup>												
Male												
2001 .....	2.1	2.7	*	0.2	1.7	2.8	4.4	3.5	1.2	0.7	1.2	*
2000 .....	2.0	1.6	0.4	0.3	1.5	2.6	4.2	3.2	1.3	0.8	1.0	1.7
1999 .....	2.1	1.9	*	0.2	1.8	2.7	4.2	3.1	1.3	0.9	1.1	2.1
Female												
2002 .....	1.3	2.6	0.4	0.2	0.7	1.4	2.6	2.3	0.9	0.5	0.7	1.1
2001 .....	1.0	2.0	0.4	*	0.5	1.2	2.2	1.7	0.6	0.4	0.7	0.8
2000 .....	0.9	2.2	*	*	0.5	1.1	1.9	1.5	0.6	0.3	0.4	1.2
1999 .....	0.9	*	*	*	0.5	1.2	2.0	1.5	0.6	0.4	0.6	1.1
Non-Hispanic white												
Both sexes												
2002 .....	1.8	1.8	0.4	0.2	1.4	2.4	3.7	2.9	1.2	0.7	0.7	1.5
2001 .....	1.6	1.7	0.3	0.1	1.3	2.2	3.3	2.5	0.9	0.5	0.9	1.0
2000 .....	1.4	1.5	0.2	0.1	1.1	1.9	3.0	2.3	0.8	0.5	0.7	1.3
1999 .....	1.4	1.3	*	0.1	1.2	2.0	3.0	2.1	0.9	0.6	0.8	1.4
Male												
2002 .....	2.3	1.9	*	0.2	2.0	3.4	4.8	3.4	1.5	0.8	0.9	2.2
2001 .....	2.1	2.0	*	0.2	1.8	3.0	4.3	3.3	1.1	0.7	1.2	*
2000 .....	2.0	*	*	0.2	1.6	2.6	4.1	3.0	1.1	0.7	1.0	*
1999 .....	2.0	*	*	0.2	1.8	2.7	4.0	2.6	1.1	0.8	1.1	2.1
Female												
2002 .....	1.3	1.8	*	0.2	0.8	1.5	2.7	2.4	0.9	0.6	0.6	1.1
2001 .....	1.1	*	*	*	0.6	1.4	2.4	1.8	0.7	0.4	0.6	0.8
2000 .....	0.9	*	*	*	0.6	1.2	1.9	1.5	0.6	0.3	0.4	1.2
1999 .....	1.0	*	*	*	0.5	1.3	1.9	1.5	0.7	0.4	0.6	1.1
Non-Hispanic black												
Both sexes												
2002 .....	2.0	5.7	*	*	0.9	2.3	4.3	4.3	1.5	*	*	*
2001 .....	1.8	5.9	*	*	0.7	1.7	4.0	3.8	1.2	*	*	*
2000 .....	1.9	*	*	0.3	0.9	2.0	4.1	3.7	1.8	*	*	*
1999 .....	2.0	*	*	*	1.3	2.3	4.3	4.2	1.2	*	*	*
Male												
2002 .....	2.9	*	*	*	1.3	3.4	6.2	6.5	2.6	*	*	*
2001 .....	2.8	6.4	*	*	1.3	2.7	6.3	6.3	2.4	*	*	*
2000 .....	2.9	*	*	*	1.6	3.2	6.1	6.0	3.0	*	*	*
1999 .....	3.1	*	*	*	2.0	3.7	6.6	7.3	2.6	*	*	*
Female												
2002 .....	1.2	*	*	*	*	1.3	2.7	2.4	*	*	*	*
2001 .....	0.9	*	*	*	*	0.9	1.9	1.5	*	*	*	*
2000 .....	1.0	*	*	*	*	0.9	2.4	1.8	*	*	*	*
1999 .....	0.9	*	*	*	*	1.0	2.4	1.6	*	*	*	*
Discharge of firearms, undetermined intent (Y22-Y24)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001 .....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
All origins <sup>3</sup>												
Both sexes												
2000 .....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999 .....	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
Male												
2002 .....	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001 .....	0.1	*	*	*	0.3	0.2	0.1	0.1	*	*	*	*
2000 .....	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999 .....	0.2	*	*	*	0.6	0.2	0.2	0.1	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001 .....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2000 .....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999 .....	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
Male												
2002 .....	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001 .....	0.1	*	*	*	0.3	0.2	0.2	*	*	*	*	*
2000 .....	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999 .....	0.2	*	*	*	0.6	0.2	0.2	0.1	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
Non-Hispanic white												
Both sexes												
2002 .....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001 .....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000 .....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
1999 .....	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
Male												
2002 .....	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001 .....	0.1	*	*	*	0.3	0.2	0.2	*	*	*	*	*
2000 .....	0.1	*	*	*	0.2	0.2	0.1	*	*	*	*	*
1999 .....	0.2	*	*	*	0.5	0.2	0.2	*	*	*	*	*
Female												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2002 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.2	*	*	*	0.6	*	*	*	*	*	*	*
Male												
2002 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	1.1	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	1.6	2.5	0.4	0.2	1.0	2.0	3.3	2.8	1.2	0.7	0.7	1.4
2001 .....	1.4	2.1	0.3	0.1	0.9	1.7	3.0	2.5	0.8	0.5	0.8	1.0
2000 .....	1.3	1.7	0.3	0.1	0.8	1.6	2.8	2.2	0.9	0.5	0.6	1.3
1999 .....	1.3	1.4	0.3	0.1	0.8	1.7	2.7	2.1	0.9	0.5	0.7	1.3
Male												
2002 .....	2.1	2.5	0.3	0.2	1.4	2.7	4.2	3.4	1.5	0.9	0.8	2.2
2001 .....	1.9	2.4	*	0.2	1.3	2.4	4.0	3.3	1.1	0.7	1.0	*
2000 .....	1.8	1.4	0.3	0.2	1.1	2.2	3.8	3.0	1.2	0.7	0.9	*
1999 .....	1.8	1.7	*	0.1	1.2	2.4	3.8	2.8	1.2	0.7	0.9	1.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)												
All origins <sup>3</sup>												
Female												
2002 .....	1.1	2.5	0.4	0.1	0.6	1.2	2.3	2.1	0.8	0.5	0.6	1.1
2001 .....	0.9	1.7	0.3	*	0.5	1.1	2.0	1.6	0.6	0.3	0.6	0.8
2000 .....	0.8	2.0	*	*	0.4	0.9	1.7	1.4	0.6	0.3	0.4	1.2
1999 .....	0.8	1.2	0.3	*	0.4	1.1	1.7	1.4	0.6	0.4	0.5	1.1
Hispanic												
Both sexes												
2002 .....	0.8	*	*	*	0.5	0.9	1.5	1.3	*	*	*	*
2001 .....	0.7	*	*	*	0.4	0.9	1.6	1.5	*	*	*	*
2000 .....	0.6	*	*	*	0.3	0.9	1.2	0.8	*	*	*	*
1999 .....	0.6	*	*	*	0.5	1.0	1.2	0.9	*	*	*	*
Male												
2002 .....	1.1	*	*	*	0.9	1.3	2.0	2.2	*	*	*	*
2001 .....	1.1	*	*	*	*	1.3	2.5	2.0	*	*	*	*
2000 .....	0.8	*	*	*	*	1.3	1.7	1.3	*	*	*	*
1999 .....	1.0	*	*	*	0.9	1.5	1.8	1.5	*	*	*	*
Female												
2002 .....	0.4	*	*	*	*	*	0.9	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.7	2.7	0.3	0.2	1.1	2.2	3.5	2.9	1.2	0.7	0.7	1.4
2001 .....	1.5	2.4	0.3	0.1	1.0	1.9	3.2	2.5	0.8	0.5	0.8	0.9
2000 .....	1.4	1.9	0.3	0.1	0.9	1.7	2.9	2.3	0.9	0.5	0.6	1.3
1999 .....	1.4	1.6	0.3	0.1	0.8	1.8	2.9	2.2	0.9	0.5	0.7	1.3
Male												
2002 .....	2.2	2.8	0.3	0.2	1.5	3.0	4.5	3.5	1.5	0.9	0.8	2.1
2001 .....	2.0	2.7	*	0.2	1.4	2.6	4.2	3.4	1.1	0.6	1.0	*
2000 .....	1.9	1.6	0.4	0.2	1.2	2.4	4.1	3.1	1.2	0.7	0.9	*
1999 .....	1.9	1.9	*	0.1	1.2	2.5	4.0	2.9	1.2	0.7	0.9	1.7
Female												
2002 .....	1.2	2.6	0.4	0.2	0.7	1.3	2.5	2.3	0.9	0.5	0.6	1.1
2001 .....	1.0	2.0	0.4	*	0.5	1.2	2.2	1.6	0.6	0.3	0.7	0.8
2000 .....	0.9	2.2	*	*	0.4	1.0	1.8	1.5	0.6	0.3	0.4	1.2
1999 .....	0.9	*	*	*	0.4	1.2	1.9	1.4	0.6	0.4	0.5	1.1
Non-Hispanic white												
Both sexes												
2002 .....	1.7	1.8	0.4	0.2	1.3	2.3	3.6	2.8	1.2	0.6	0.7	1.5
2001 .....	1.5	1.7	0.3	0.1	1.1	2.0	3.2	2.5	0.8	0.4	0.8	0.9
2000 .....	1.4	1.5	0.2	0.1	0.9	1.8	2.9	2.2	0.8	0.5	0.6	1.2
1999 .....	1.3	1.3	*	0.1	0.9	1.9	2.8	2.0	0.9	0.5	0.7	1.3
Male												
2002 .....	2.2	1.9	*	0.2	1.7	3.2	4.6	3.3	1.4	0.7	0.8	2.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)												
Non-Hispanic white												
Male												
2001 .....	2.0	2.0	*	0.2	1.6	2.7	4.1	3.2	1.0	0.6	1.0	*
2000 .....	1.8	*	*	0.2	1.4	2.4	4.0	2.9	1.0	0.6	0.9	*
1999 .....	1.8	*	*	0.2	1.4	2.5	3.7	2.5	1.0	0.7	0.9	*
Female												
2002 .....	1.3	1.8	*	*	0.8	1.5	2.7	2.4	0.9	0.5	0.6	1.1
2001 .....	1.1	*	*	*	0.6	1.3	2.3	1.7	0.7	0.3	0.6	0.8
2000 .....	0.9	*	*	*	0.5	1.1	1.9	1.5	0.6	0.3	0.4	1.2
1999 .....	0.9	*	*	*	0.4	1.3	1.9	1.5	0.7	0.4	0.6	1.1
Non-Hispanic black												
Both sexes												
2002 .....	1.9	5.7	*	*	0.6	2.1	4.2	4.3	1.5	*	*	*
2001 .....	1.7	5.9	*	*	0.5	1.7	3.9	3.7	1.1	*	*	*
2000 .....	1.7	*	*	*	0.6	1.7	4.0	3.7	1.6	*	*	*
1999 .....	1.8	*	*	*	0.6	2.0	4.2	4.1	1.2	*	*	*
Male												
2002 .....	2.7	*	*	*	0.8	3.0	6.0	6.5	2.6	*	*	*
2001 .....	2.6	6.4	*	*	0.8	2.5	6.3	6.2	2.1	*	*	*
2000 .....	2.6	*	*	*	1.0	2.7	5.9	5.9	2.7	*	*	*
1999 .....	2.8	*	*	*	0.9	3.2	6.4	7.1	2.5	*	*	*
Female												
2002 .....	1.2	*	*	*	*	1.3	2.6	2.4	*	*	*	*
2001 .....	0.9	*	*	*	*	0.9	1.9	1.5	*	*	*	*
2000 .....	0.9	*	*	*	*	0.8	2.3	1.8	*	*	*	*
1999 .....	0.9	*	*	*	*	0.9	2.3	1.6	*	*	*	*
Operations of war and their sequelae (Y36,Y89.1)												
All origins <sup>3</sup>												
Both sexes												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
Hispanic												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Operations of war and their sequelae (Y36,Y89.1)</b>												
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	*	*	*	*	*	*	*	*	*	*	*	*
2001 .....	*	*	*	*	*	*	*	*	*	*	*	*
2000 .....	*	*	*	*	*	*	*	*	*	*	*	*
1999 .....	*	*	*	*	*	*	*	*	*	*	*	*
<b>Complications of medical and surgical care (Y40-Y84,Y88)</b>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	1.0	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.3
2001 .....	1.1	0.6	0.2	*	0.1	0.2	0.4	0.7	1.5	3.1	6.9	12.8
2000 .....	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.5	3.5	6.7	13.6
1999 .....	1.0	1.4	0.2	*	0.1	0.2	0.3	0.6	1.3	3.2	6.6	12.7
<b>Male</b>												
2002 .....	0.9	*	*	0.1	*	0.2	0.4	0.6	1.3	3.1	6.3	12.3
2001 .....	1.0	*	*	*	*	0.2	0.4	0.7	1.6	3.2	7.7	14.5
2000 .....	1.0	*	*	*	*	0.2	0.3	0.8	1.6	4.0	8.2	16.0
1999 .....	1.0	1.6	*	*	0.1	0.2	0.3	0.5	1.4	3.4	7.8	16.5
<b>Female</b>												
2002 .....	1.1	*	*	*	*	0.2	0.4	0.7	1.3	3.0	6.2	10.9
2001 .....	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.3	12.1
2000 .....	1.1	*	*	*	*	0.2	0.3	0.8	1.5	3.1	5.7	12.7
1999 .....	1.1	1.2	*	*	*	0.2	0.3	0.6	1.2	3.0	5.9	11.2
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	0.4	*	*	*	*	*	*	*	*	2.4	4.8	*
2001 .....	0.4	*	*	*	*	*	*	0.6	*	2.2	5.2	*
2000 .....	0.4	*	*	*	*	*	*	*	*	3.3	5.1	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
<b>Male</b>												
2002 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
<b>Female</b>												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001 .....	0.4	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.



Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Hispanic												
Female												
2002 .....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 .....	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	1.1	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.6
2001 .....	1.2	0.7	*	*	0.1	0.2	0.4	0.7	1.6	3.1	6.9	12.9
2000 .....	1.2	*	*	0.1	0.1	0.2	0.3	0.8	1.6	3.5	6.7	13.7
1999 .....	1.1	1.6	0.2	*	0.1	0.2	0.4	0.6	1.3	3.3	6.8	12.8
Male												
2002 .....	1.0	*	*	0.1	*	0.3	0.4	0.7	1.4	3.1	6.3	12.9
2001 .....	1.1	*	*	*	*	0.2	0.4	0.8	1.7	3.3	7.7	14.5
2000 .....	1.1	*	*	*	*	0.3	0.4	0.8	1.6	4.0	8.2	16.1
1999 .....	1.0	1.8	*	*	*	0.2	0.4	0.6	1.5	3.5	7.9	16.8
Female												
2002 .....	1.2	*	*	*	*	0.2	0.4	0.7	1.3	3.1	6.2	11.1
2001 .....	1.2	*	*	*	*	0.2	0.4	0.7	1.5	3.0	6.4	12.2
2000 .....	1.2	*	*	*	*	0.2	0.3	0.8	1.6	3.1	5.7	12.7
1999 .....	1.2	1.4	*	*	*	0.2	0.4	0.6	1.2	3.1	6.1	11.2
Non-Hispanic white												
Both sexes												
2002 .....	1.1	*	*	*	0.1	0.2	0.3	0.6	1.2	3.0	6.1	11.5
2001 .....	1.2	*	*	*	0.1	0.2	0.3	0.6	1.4	2.8	6.8	12.2
2000 .....	1.2	*	*	*	*	0.2	0.3	0.8	1.4	3.3	6.5	13.2
1999 .....	1.1	1.3	*	*	0.1	0.2	0.3	0.5	1.2	3.1	6.5	12.1
Male												
2002 .....	1.0	*	*	*	*	0.3	0.4	0.6	1.3	2.9	6.3	12.4
2001 .....	1.0	*	*	*	*	0.2	0.3	0.6	1.4	2.9	7.8	13.5
2000 .....	1.2	*	*	*	*	0.2	0.4	0.8	1.4	3.7	8.2	15.9
1999 .....	1.1	*	*	*	*	0.2	0.3	0.5	1.3	3.3	7.7	15.8
Female												
2002 .....	1.2	*	*	*	*	*	0.3	0.6	1.1	3.0	6.0	11.1
2001 .....	1.3	*	*	*	*	0.2	0.3	0.6	1.4	2.7	6.2	11.6
2000 .....	1.2	*	*	*	*	0.2	0.3	0.7	1.4	3.0	5.4	12.0
1999 .....	1.2	*	*	*	*	*	0.3	0.6	1.1	3.0	5.7	10.7
Non-Hispanic black												
Both sexes												
2002 .....	1.2	*	*	*	*	*	0.9	1.7	2.7	4.7	8.5	14.2
2001 .....	1.4	*	*	*	*	0.4	0.9	1.6	3.4	6.4	9.3	23.7
2000 .....	1.4	*	*	*	*	0.4	0.6	1.4	3.6	6.2	10.6	21.3
1999 .....	1.3	*	*	*	*	*	0.6	1.3	2.6	5.2	10.6	20.4
Male												
2002 .....	1.1	*	*	*	*	*	*	1.4	2.6	5.4	7.3	*
2001 .....	1.5	*	*	*	*	*	1.0	1.9	4.3	7.9	9.0	30.0
2000 .....	1.2	*	*	*	*	*	*	1.3	3.2	7.8	10.2	*
1999 .....	1.2	*	*	*	*	*	*	1.4	2.9	6.0	11.9	23.8

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Non-Hispanic black												
Female												
2002 .....	1.4	*	*	*	*	*	1.1	1.9	2.8	4.2	9.1	11.7
2001 .....	1.4	*	*	*	*	*	0.7	1.4	2.7	5.3	9.4	21.4
2000 .....	1.6	*	*	*	*	*	*	1.5	3.9	5.2	10.8	22.2
1999 .....	1.4	*	*	*	*	*	*	1.2	2.4	4.7	9.8	19.1
Alcohol-induced causes (F10,G31.2,G62.1,I42.6,K29.2,K70,R78.0,X45,X65,Y15) <sup>6</sup>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	6.9	*	*	*	0.2	1.5	8.0	16.6	18.1	15.4	9.3	4.3
2001 .....	7.0	*	*	*	0.3	1.6	8.2	16.9	17.9	15.3	9.5	5.0
2000 .....	6.9	*	*	*	0.2	1.6	8.3	16.1	18.5	15.7	9.8	5.3
1999 .....	6.9	*	*	*	0.3	1.5	8.3	16.2	18.4	15.7	10.5	5.5
Male												
2002 .....	10.6	*	*	*	0.4	2.2	11.3	25.9	29.6	25.7	16.6	9.6
2001 .....	10.7	*	*	*	0.4	2.2	11.7	26.2	29.2	26.0	17.1	11.5
2000 .....	10.7	*	*	*	0.4	2.4	12.2	25.6	29.7	26.8	18.5	11.7
1999 .....	10.7	*	*	*	0.4	2.2	12.3	25.8	30.1	27.1	19.2	11.8
Female												
2002 .....	3.3	*	*	*	0.7	4.7	7.6	7.5	6.7	4.5	2.0	
2001 .....	3.4	*	*	*	0.9	4.7	8.0	7.4	6.5	4.5	2.3	
2000 .....	3.2	*	*	*	0.7	4.5	7.0	8.2	6.5	4.0	2.7	
1999 .....	3.2	*	*	*	0.1	0.9	4.4	6.8	7.7	6.4	4.8	3.0
Hispanic												
Both sexes												
2002 .....	6.1	*	*	*	0.5	1.9	8.6	21.0	26.2	26.7	14.5	*
2001 .....	6.4	*	*	*	0.5	2.1	9.0	22.4	28.1	26.0	20.4	*
2000 .....	6.5	*	*	*	0.5	2.0	10.1	21.9	29.3	28.7	19.0	*
1999 .....	6.4	*	*	*	0.4	2.0	10.1	22.5	28.5	25.9	18.7	14.0
Male												
2002 .....	10.2	*	*	*	0.8	3.2	14.0	36.0	48.3	52.0	26.7	*
2001 .....	10.6	*	*	*	0.7	3.2	14.6	38.5	51.5	52.1	38.0	*
2000 .....	11.0	*	*	*	0.8	3.5	16.3	38.5	54.5	58.5	42.3	*
1999 .....	10.6	*	*	*	0.7	3.2	16.6	39.2	52.6	50.5	36.0	*
Female												
2002 .....	1.8	*	*	*	*	*	2.7	6.1	6.4	6.5	6.2	*
2001 .....	2.0	*	*	*	*	0.8	2.9	6.6	7.4	5.4	8.5	*
2000 .....	1.7	*	*	*	*	*	3.4	5.7	7.1	5.3	*	*
1999 .....	1.9	*	*	*	*	*	3.2	6.2	7.3	6.5	7.3	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	7.0	*	*	*	0.2	1.3	7.8	16.0	17.3	14.5	9.0	4.1
2001 .....	7.0	*	*	*	0.2	1.5	8.0	16.2	16.9	14.5	9.0	4.9
2000 .....	6.9	*	*	*	0.2	1.5	8.0	15.4	17.6	14.7	9.3	5.2
1999 .....	6.9	*	*	*	0.2	1.4	8.0	15.5	17.5	15.0	10.1	5.2

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcohol-induced causes (F10,G31.2,G62.1,I42.6,K29.2,K70, R78.0,X45,X65,Y15) <sup>6</sup>												
Non-Hispanic <sup>4</sup>												
Male												
2002 .....	10.6	*	*	*	0.3	1.9	10.8	24.6	27.8	23.8	15.9	9.0
2001 .....	10.6	*	*	*	0.4	2.0	11.2	24.7	27.3	24.1	16.1	11.2
2000 .....	10.6	*	*	*	0.3	2.2	11.5	24.1	27.6	24.7	17.4	11.5
1999 .....	10.6	*	*	*	0.3	1.9	11.6	24.4	28.2	25.5	18.4	11.1
Female												
2002 .....	3.5	*	*	*	*	0.8	4.9	7.7	7.6	6.8	4.4	1.9
2001 .....	3.6	*	*	*	*	0.9	4.9	8.1	7.4	6.6	4.3	2.3
2000 .....	3.4	*	*	*	*	0.8	4.6	7.0	8.3	6.5	4.1	2.6
1999 .....	3.3	*	*	*	*	0.9	4.5	6.9	7.7	6.3	4.7	2.9
Non-Hispanic white												
Both sexes												
2002 .....	7.2	*	*	*	0.2	1.3	7.9	15.9	16.6	14.2	8.9	4.0
2001 .....	7.1	*	*	*	0.2	1.4	7.9	15.6	16.2	14.1	8.9	4.7
2000 .....	6.9	*	*	*	0.2	1.4	7.8	14.7	16.6	14.3	9.2	5.1
1999 .....	6.8	*	*	*	0.2	1.4	7.6	14.6	16.4	14.3	9.7	5.0
Male												
2002 .....	10.9	*	*	*	0.3	1.8	10.9	24.4	26.5	22.8	15.7	8.6
2001 .....	10.7	*	*	*	0.3	2.0	10.8	23.8	25.6	22.9	15.9	10.7
2000 .....	10.6	*	*	*	0.3	2.0	11.2	23.1	26.0	23.7	17.1	11.1
1999 .....	10.6	*	*	*	0.3	1.9	11.0	23.0	26.2	23.9	17.8	10.5
Female												
2002 .....	3.6	*	*	*	*	0.7	4.9	7.6	7.2	6.9	4.3	2.0
2001 .....	3.6	*	*	*	*	0.8	4.9	7.6	7.2	6.7	4.3	2.2
2000 .....	3.3	*	*	*	*	0.7	4.4	6.4	7.7	6.5	4.0	2.7
1999 .....	3.2	*	*	*	*	0.9	4.1	6.2	7.1	6.3	4.5	2.8
Non-Hispanic black												
Both sexes												
2002 .....	6.5	*	*	*	*	1.2	6.8	18.4	24.9	18.0	10.9	*
2001 .....	7.3	*	*	*	*	1.3	8.6	22.0	25.5	20.1	10.9	*
2000 .....	7.4	*	*	*	*	1.7	8.9	22.5	26.3	20.9	11.0	6.4
1999 .....	7.8	*	*	*	*	1.3	9.8	23.1	28.9	23.0	14.5	7.4
Male												
2002 .....	10.0	*	*	*	*	1.9	9.4	29.1	42.7	35.1	21.4	*
2001 .....	11.6	*	*	*	*	1.8	13.2	35.0	45.5	39.6	21.9	*
2000 .....	11.4	*	*	*	*	2.4	12.9	35.6	43.9	39.5	23.4	*
1999 .....	12.2	*	*	*	*	1.6	14.2	37.0	50.2	45.1	27.8	*
Female												
2002 .....	3.3	*	*	*	*	*	4.4	9.2	10.6	5.9	5.0	*
2001 .....	3.4	*	*	*	*	0.9	4.5	10.9	9.4	6.4	4.7	*
2000 .....	3.8	*	*	*	*	1.1	5.3	11.2	12.1	7.8	4.0	*
1999 .....	3.9	*	*	*	*	0.9	6.0	11.1	11.9	7.4	7.0	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Drug-induced causes</b>												
(F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9, F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9, F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0, F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5, F19.7-F19.9,X40-X44,X60-X64,X85,Y10-Y14) <sup>6</sup>												
<b>All origins <sup>3</sup></b>												
<b>Both sexes</b>												
2002 .....	9.0	0.7	0.2	0.1	5.5	11.1	19.7	17.9	6.9	3.5	3.7	5.6
2001 .....	7.6	0.5	0.2	0.1	4.5	9.4	17.0	14.8	5.4	3.0	3.5	5.1
2000 .....	7.0	*	*	0.1	4.0	8.8	16.0	13.2	4.9	2.6	3.4	5.5
1999 .....	6.8	0.6	0.2	0.1	3.5	8.9	15.7	12.6	4.9	3.0	3.7	4.8
<b>Male</b>												
2002 .....	11.8	1.0	*	0.1	7.8	15.5	25.6	23.0	8.1	3.7	4.2	7.3
2001 .....	10.2	*	0.3	0.1	6.6	13.2	22.7	19.7	6.3	3.5	3.8	5.6
2000 .....	9.5	*	*	0.1	5.8	12.5	21.8	18.1	5.4	2.8	3.8	7.3
1999 .....	9.4	*	*	*	4.9	12.7	21.9	17.5	6.0	3.4	3.8	6.0
<b>Female</b>												
2002 .....	6.3	*	0.3	*	3.0	6.6	13.9	13.1	5.7	3.4	3.5	4.8
2001 .....	5.1	*	*	0.1	2.4	5.6	11.5	10.0	4.5	2.6	3.2	4.9
2000 .....	4.6	*	*	*	2.1	5.1	10.2	8.5	4.4	2.5	3.2	4.8
1999 .....	4.4	*	*	*	2.0	5.0	9.6	8.0	3.8	2.7	3.7	4.3
<b>Hispanic</b>												
<b>Both sexes</b>												
2002 .....	5.5	*	*	*	3.2	6.2	13.4	13.8	5.6	3.0	3.3	*
2001 .....	4.7	*	*	*	2.4	5.9	11.8	11.0	4.8	2.6	*	*
2000 .....	4.8	*	*	*	2.2	6.9	12.7	10.6	4.6	1.9	*	*
1999 .....	5.8	*	*	*	2.7	8.1	15.0	13.4	5.1	2.7	*	*
<b>Male</b>												
2002 .....	8.2	*	*	*	4.7	9.3	19.9	21.1	8.7	*	*	*
2001 .....	7.0	*	*	*	3.3	8.7	17.8	17.3	7.9	*	*	*
2000 .....	7.4	*	*	*	3.4	10.5	19.6	17.1	6.5	*	*	*
1999 .....	9.2	*	*	*	4.2	12.7	23.8	21.9	9.0	4.6	*	*
<b>Female</b>												
2002 .....	2.6	*	*	*	1.4	2.5	6.3	6.6	2.9	*	*	*
2001 .....	2.2	*	*	*	1.4	2.8	5.3	4.8	*	*	*	*
2000 .....	2.1	*	*	*	0.9	2.7	5.2	4.2	3.0	*	*	*
1999 .....	2.2	*	*	*	1.0	2.8	5.7	5.2	*	*	*	*
<b>Non-Hispanic <sup>4</sup></b>												
<b>Both sexes</b>												
2002 .....	9.5	0.8	0.2	0.1	5.9	12.2	20.5	18.2	6.9	3.5	3.7	5.7
2001 .....	8.0	0.7	0.2	0.1	4.9	10.1	17.6	15.0	5.3	3.0	3.5	5.2
2000 .....	7.2	*	*	0.1	4.3	9.1	16.3	13.2	4.8	2.6	3.5	5.6
1999 .....	6.9	*	0.2	0.1	3.6	9.0	15.6	12.4	4.7	3.0	3.8	4.8
<b>Male</b>												
2002 .....	12.3	*	*	0.1	8.4	17.0	26.3	23.0	8.0	3.6	4.1	7.4
2001 .....	10.6	*	*	0.1	7.3	14.1	23.1	19.7	6.0	3.5	3.8	5.7
2000 .....	9.7	*	*	0.1	6.3	12.8	21.9	17.9	5.2	2.8	3.9	7.2
1999 .....	9.3	*	*	*	5.1	12.7	21.4	16.8	5.6	3.3	3.9	6.1

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>Drug-induced causes</b>												
(F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9, F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9, F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0, F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5, F19.7-F19.9,X40-X44,X60-X64,X85,Y10-Y14) <sup>6</sup>												
<b>Non-Hispanic <sup>4</sup></b>												
<b>Female</b>												
2002 .....	6.9	*	*	*	3.3	7.5	14.9	13.6	5.9	3.4	3.5	5.0
2001 .....	5.5	*	*	*	2.5	6.1	12.2	10.5	4.7	2.6	3.3	5.0
2000 .....	4.9	*	*	*	2.3	5.5	10.8	8.7	4.5	2.5	3.3	5.0
1999 .....	4.6	*	*	*	2.1	5.4	10.0	8.2	3.9	2.7	3.7	4.3
<b>Non-Hispanic white</b>												
<b>Both sexes</b>												
2002 .....	10.0	*	*	0.1	7.1	13.8	21.6	17.8	6.6	3.6	3.9	5.8
2001 .....	8.3	*	*	0.1	5.9	11.3	18.1	14.2	5.1	3.0	3.6	5.1
2000 .....	7.4	*	*	0.1	5.1	9.9	16.4	12.3	4.7	2.7	3.7	5.5
1999 .....	6.9	*	*	0.1	4.1	9.5	15.4	11.4	4.4	3.0	4.0	4.8
<b>Male</b>												
2002 .....	12.8	*	*	*	10.1	19.2	27.5	21.7	7.0	3.5	4.0	7.7
2001 .....	10.8	*	*	*	8.6	15.8	23.7	17.8	5.1	3.4	3.8	5.6
2000 .....	9.8	*	*	0.2	7.5	13.8	21.9	16.1	4.6	2.7	4.1	7.0
1999 .....	9.1	*	*	*	5.8	13.2	21.0	14.7	4.7	3.2	4.0	6.4
<b>Female</b>												
2002 .....	7.4	*	*	*	3.9	8.4	15.7	13.9	6.3	3.7	3.8	5.0
2001 .....	5.8	*	*	*	3.0	6.8	12.5	10.6	5.0	2.8	3.5	4.9
2000 .....	5.1	*	*	*	2.6	6.0	10.8	8.6	4.8	2.7	3.4	5.0
1999 .....	4.8	*	*	*	2.4	5.7	9.9	8.1	4.2	2.9	4.0	4.2
<b>Non-Hispanic black</b>												
<b>Both sexes</b>												
2002 .....	9.4	*	*	*	2.1	8.7	21.2	27.3	11.7	2.9	2.5	*
2001 .....	8.7	*	*	*	1.9	7.8	20.4	25.6	8.6	3.1	2.7	6.2
2000 .....	8.5	*	*	*	2.0	8.5	20.6	24.1	7.5	2.9	2.4	7.6
1999 .....	8.7	*	*	*	2.2	9.3	21.2	23.8	8.1	3.4	*	*
<b>Male</b>												
2002 .....	13.2	*	*	*	3.0	12.2	28.3	40.1	20.2	5.0	*	*
2001 .....	12.5	*	*	*	2.9	11.0	27.5	40.1	14.8	5.2	*	*
2000 .....	12.3	*	*	*	2.7	12.3	28.7	38.1	12.3	4.9	*	*
1999 .....	12.9	*	*	*	3.4	13.8	30.4	38.3	14.9	5.1	*	*
<b>Female</b>												
2002 .....	6.0	*	*	*	1.3	5.6	14.9	16.3	4.8	*	*	*
2001 .....	5.3	*	*	*	0.9	5.0	14.1	13.1	3.7	*	*	*
2000 .....	5.0	*	*	*	1.3	5.0	13.4	12.1	3.6	*	*	*
1999 .....	4.9	*	*	*	1.1	5.3	13.1	11.5	2.7	2.1	*	*

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95, Y22-Y24,Y35.0) <sup>6</sup>												
All origins <sup>3</sup>												
Both sexes												
2002 .....	10.5	*	0.4	0.8	16.7	15.4	12.1	10.8	10.2	10.9	14.4	12.5
2001 .....	10.4	*	0.5	0.8	16.7	15.5	11.7	10.5	10.1	10.9	14.3	12.8
2000 .....	10.2	*	0.3	0.9	16.8	14.5	11.9	10.5	9.4	10.6	13.9	14.2
1999 .....	10.3	*	0.4	1.0	17.6	14.9	11.6	10.2	9.7	11.0	14.2	13.5
Male												
2002 .....	18.4	*	0.5	1.2	29.3	26.5	20.1	18.2	18.0	21.3	32.9	38.9
2001 .....	18.2	*	0.5	1.2	29.6	26.7	19.2	17.8	17.4	21.1	32.8	40.2
2000 .....	17.8	*	0.4	1.4	29.4	24.9	19.4	17.6	16.3	20.3	32.2	44.7
1999 .....	18.1	*	0.5	1.5	30.6	25.5	18.9	17.0	17.1	21.5	32.8	43.3
Female												
2002 .....	2.8	*	0.3	0.5	3.5	4.0	4.2	3.6	3.1	2.3	2.1	1.1
2001 .....	2.8	*	0.4	0.4	3.2	4.0	4.3	3.5	3.3	2.4	2.2	1.3
2000 .....	2.8	*	*	0.4	3.5	4.0	4.4	3.6	3.0	2.5	2.0	1.7
1999 .....	2.9	*	0.4	0.5	3.9	4.1	4.4	3.6	2.9	2.4	2.1	1.5
Hispanic												
Both sexes												
2002 .....	8.1	*	*	0.8	18.5	12.5	7.9	6.0	5.0	5.3	5.5	*
2001 .....	8.3	*	*	0.6	18.4	13.5	8.6	6.3	4.4	5.3	5.6	*
2000 .....	8.2	*	*	0.8	18.0	12.1	8.1	8.1	4.9	5.9	5.5	*
1999 .....	8.5	*	*	1.1	18.8	12.2	8.7	7.4	5.3	4.9	9.2	*
Male												
2002 .....	14.2	*	*	1.0	32.1	21.2	12.9	10.1	9.5	10.9	13.1	*
2001 .....	14.6	*	*	0.9	31.4	22.7	14.4	10.7	8.6	11.0	12.5	*
2000 .....	14.2	*	*	1.3	30.8	20.3	13.2	13.5	9.1	11.4	12.8	*
1999 .....	14.6	*	*	1.5	31.6	20.6	14.3	11.8	9.9	10.1	21.9	*
Female												
2002 .....	1.6	*	*	*	2.8	2.3	2.5	1.9	*	*	*	*
2001 .....	1.7	*	*	*	3.3	2.7	2.3	2.0	*	*	*	*
2000 .....	1.8	*	*	*	2.9	2.6	2.5	2.8	*	*	*	*
1999 .....	2.0	*	*	*	3.8	2.4	2.6	3.0	*	*	*	*
Non-Hispanic <sup>4</sup>												
Both sexes												
2002 .....	10.8	*	0.4	0.9	16.2	15.9	12.7	11.2	10.6	11.2	14.8	12.7
2001 .....	10.6	*	0.5	0.9	16.3	15.8	12.1	10.9	10.4	11.2	14.7	12.9
2000 .....	10.4	*	0.3	0.9	16.4	14.9	12.3	10.7	9.7	10.8	14.3	14.4
1999 .....	10.6	*	0.5	1.0	17.3	15.3	11.9	10.4	10.0	11.4	14.4	13.6
Male												
2002 .....	19.0	*	0.5	1.2	28.4	27.6	21.2	18.9	18.5	21.9	33.8	39.5
2001 .....	18.7	*	0.5	1.2	29.0	27.5	19.9	18.4	17.9	21.7	33.6	40.8
2000 .....	18.3	*	0.4	1.5	28.9	25.7	20.1	17.9	16.7	20.7	33.0	45.8
1999 .....	18.5	*	0.5	1.5	30.3	26.3	19.3	17.4	17.6	22.1	33.2	43.9
Female												
2002 .....	3.0	*	*	0.5	3.6	4.3	4.4	3.8	3.2	2.4	2.2	1.2
2001 .....	3.0	*	0.4	0.5	3.1	4.3	4.6	3.7	3.5	2.5	2.3	1.4
2000 .....	3.0	*	*	0.4	3.6	4.2	4.6	3.7	3.1	2.6	2.1	1.7

See footnotes at end of table.

Worktable Orig290R. Deaths rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages <sup>1</sup>	Under 1 year <sup>2</sup>	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95, Y22-Y24,Y35.0) <sup>6</sup>												
Non-Hispanic <sup>4</sup>												
Female												
1999 .....	3.0	*	0.4	0.5	3.9	4.4	4.6	3.6	3.0	2.5	2.2	1.5
Non-Hispanic white												
Both sexes												
2002 .....	9.4	*	0.3	0.6	9.2	10.7	11.8	11.5	11.4	12.3	15.8	13.7
2001 .....	9.4	*	0.3	0.8	9.4	11.4	11.4	11.4	11.2	12.1	15.7	14.0
2000 .....	9.1	*	0.2	0.8	9.6	10.5	11.5	10.8	10.2	11.5	15.4	15.5
1999 .....	9.3	*	0.3	0.9	10.1	11.2	11.2	10.5	10.5	12.2	15.3	14.7
Male												
2002 .....	16.3	*	*	0.9	15.6	17.4	19.3	19.2	19.7	23.7	35.9	42.5
2001 .....	16.3	*	*	1.2	16.0	18.9	18.4	19.0	19.0	23.1	35.7	44.0
2000 .....	15.7	*	*	1.3	16.2	17.2	18.4	18.0	17.5	21.9	35.5	49.6
1999 .....	15.9	*	*	1.3	17.2	18.6	17.8	17.6	18.1	23.5	35.1	47.5
Female												
2002 .....	2.8	*	*	0.4	2.5	3.8	4.3	4.0	3.5	2.6	2.3	1.2
2001 .....	2.9	*	*	0.4	2.5	3.7	4.4	3.8	3.9	2.7	2.4	1.5
2000 .....	2.9	*	*	0.4	2.7	3.7	4.5	3.8	3.4	2.8	2.2	1.8
1999 .....	2.9	*	*	0.5	2.8	3.8	4.5	3.6	3.3	2.7	2.3	1.6
Non-Hispanic black												
Both sexes												
2002 .....	20.7	*	*	1.6	49.2	46.4	21.0	11.9	7.2	5.8	6.9	*
2001 .....	19.8	*	1.0	1.2	49.6	42.2	19.5	10.7	6.9	6.6	7.4	*
2000 .....	19.8	*	*	1.4	49.8	40.9	19.7	12.1	6.8	7.0	5.6	*
1999 .....	20.0	*	1.1	1.6	51.8	38.9	18.8	11.6	8.4	6.8	7.6	*
Male												
2002 .....	38.9	*	*	2.3	89.8	89.0	37.9	21.7	13.7	12.0	17.0	*
2001 .....	37.3	*	*	1.8	93.1	80.4	34.1	19.3	13.6	13.8	17.7	*
2000 .....	37.1	*	*	2.2	92.2	77.6	35.3	21.7	12.8	15.4	12.3	*
1999 .....	37.1	*	*	2.4	95.0	73.8	33.1	20.7	16.8	14.3	18.4	*
Female												
2002 .....	4.3	*	*	0.9	8.4	7.8	6.1	3.5	2.1	*	*	*
2001 .....	4.0	*	*	*	6.3	7.7	6.7	3.3	1.5	*	*	*
2000 .....	4.2	*	*	*	7.9	7.7	5.9	3.9	2.0	*	*	*
1999 .....	4.5	*	*	0.9	9.3	7.6	6.2	3.8	1.6	*	*	*

\* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

<sup>1</sup> Figures for age not stated included in "All ages" but not distributed among age groups.

<sup>2</sup> Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

<sup>3</sup> Figures for origin not stated are included in "All origins" but not distributed among specified origins.

<sup>4</sup> Includes races other than white and black.

<sup>5</sup> Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

<sup>6</sup> Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.